

MATH 6101 Fall 2008

Euler and Trigonometric Series



Sums of Series

What do we know about sums of series?

How many series sums do we know? can we find?

Examples:

$$\sum_{n=0}^{\infty} a^n = \frac{1}{1-a}, \quad |a| < 1$$

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1$$

01-Oct-2008

MATH 6101

2

Sums of Series

Claim 1: $\sum_{n=1}^{\infty} \frac{n}{2^n} = 2$

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{n-1}{2^{n+1}} &= \sum_{n=1}^{\infty} \left(\frac{n}{2^n} - \frac{n+1}{2^{n+1}} \right) = \sum_{n=1}^{\infty} \frac{n}{2^n} - \sum_{n=1}^{\infty} \frac{n+1}{2^{n+1}} \\ &= \frac{1}{2} + \sum_{k=2}^{\infty} \frac{k}{2^k} - \sum_{k=2}^{\infty} \frac{k}{2^k} \end{aligned}$$

$$\sum_{n=1}^{\infty} \frac{n-1}{2^{n+1}} = \frac{1}{2}$$

01-Oct-2008

MATH 6101

3

Sums of Series

$$\frac{1}{2} = \sum_{n=1}^{\infty} \frac{n-1}{2^{n+1}} = \frac{1}{4} \sum_{n=1}^{\infty} \frac{n-1}{2^{n-1}} = \frac{1}{4} \sum_{k=1}^{\infty} \frac{k}{2^k}$$

$$\sum_{k=1}^{\infty} \frac{k}{2^k} = 2$$

01-Oct-2008 MATH 6101 4

Sums of Series

Claim 2: $\sum_{n=1}^{\infty} \frac{n^2}{2^n} = 6$

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{n^2-1}{2^{n+1}} &= \sum_{n=1}^{\infty} \left(\frac{n^2}{2^n} - \frac{n^2+1}{2^{n+1}} \right) = \sum_{n=1}^{\infty} \frac{n^2}{2^n} - \sum_{n=1}^{\infty} \frac{n^2+1}{2^{n+1}} + 2 \sum_{n=1}^{\infty} \frac{n}{2^n} - \sum_{n=1}^{\infty} \frac{2n}{2^n} \\ &= \sum_{n=1}^{\infty} \frac{n^2}{2^n} - \sum_{n=1}^{\infty} \frac{n^2+2n+1}{2^{n+1}} + 4 \\ &= 4 + \sum_{n=1}^{\infty} \frac{n^2}{2^n} - \sum_{n=1}^{\infty} \frac{(n+1)^2}{2^{n+1}} \\ \sum_{n=1}^{\infty} \frac{n^2-1}{2^{n+1}} &= 4 + \frac{1}{2} + \sum_{k=2}^{\infty} \frac{k^2}{2^k} - \sum_{k=2}^{\infty} \frac{k^2}{2^k} = \frac{5}{2} \end{aligned}$$

01-Oct-2008 MATH 6101 5

Sums of Series

$$\begin{aligned} \frac{5}{2} &= \sum_{n=1}^{\infty} \frac{n^2-1}{2^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{n^2}{2^{n+1}} - \sum_{n=1}^{\infty} \frac{1}{2^{n+1}} \\ &= \frac{1}{2} \sum_{n=1}^{\infty} \frac{n^2}{2^n} - \sum_{n=2}^{\infty} \frac{1}{2^n} \\ &= \frac{1}{2} \sum_{n=1}^{\infty} \frac{n^2}{2^n} - \frac{1}{2} \\ \sum_{n=1}^{\infty} \frac{n^2}{2^n} &= 6 \end{aligned}$$

01-Oct-2008 MATH 6101 6

Sums of Series

In a like manner we can show that

$$\sum_{n=1}^{\infty} \frac{n^3}{2^n} = 26$$

On the other hand we know that

$$\sum_{n=1}^{\infty} \frac{1}{n} \text{ diverges.}$$

01-Oct-2008

MATH 6101

7

Sums of Series

How about $\sum_{n=1}^{\infty} \frac{1}{n^2}$?

First, does it converge?

We know that $2n^2 \geq n(n+1)$ so $\frac{2}{n(n+1)} \geq \frac{1}{n^2}$

$$\sum_{n=1}^{\infty} \frac{1}{n^2} \leq \sum_{n=1}^{\infty} \frac{2}{n(n+1)} = 2$$

Therefore, it does converge. But to what?

01-Oct-2008

MATH 6101

8

Sums of Series

Jakob Bernoulli considered it and failed to find it. So did Mengoli and Leibniz. Finding the sum became known as the *Basel Problem*.

We will look at how Euler solved the problem.

01-Oct-2008

MATH 6101

9

Euler's Sum

Theorem [Euler, 1735]: $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$

This was a huge surprise, as no one expected π to appear in the sum at all!!

Euler's Proof: Let $p(x)$ be a polynomial of degree n with the following properties

1. $p(x)$ has non-zero roots a_1, a_2, \dots, a_n .
2. $p(0) = 1$

Then we can write:

01-Oct-2008 MATH 6101 10

Euler's Sum

$$p(x) = \left(1 - \frac{x}{a_1}\right) \left(1 - \frac{x}{a_2}\right) \dots \left(1 - \frac{x}{a_n}\right)$$

Euler claims that "what holds for a finite polynomial holds for an infinite polynomial".

He considers the infinite polynomial.

$$p(x) = 1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \frac{x^6}{7!} + \dots$$

which is an infinite polynomial with $p(0) = 1$.

01-Oct-2008 MATH 6101 11

Euler's Sum

Euler knew an infinite polynomial for $\sin(x)$.

$$\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots + (-1)^{k+1} \frac{x^{2k+1}}{(2k+1)!} + \dots$$

So Euler noticed that $x p(x)$

$$x p(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots = \sin(x)$$

From the properties of the sine function we know that this infinite polynomial has zeros at $x = \pm k\pi$ for $k = 1, 2, 3, \dots$

01-Oct-2008 MATH 6101 12

Euler's Sum

From his claim above, Euler could write this polynomial as an infinite product.

$$\begin{aligned} p(x) &= 1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \frac{x^6}{7!} + \dots \\ &= \left(1 - \frac{x}{\pi}\right) \left(1 + \frac{x}{\pi}\right) \left(1 - \frac{x}{2\pi}\right) \left(1 + \frac{x}{2\pi}\right) \times \dots \\ &= \left(1 - \frac{x^2}{\pi^2}\right) \left(1 - \frac{x^2}{4\pi^2}\right) \left(1 - \frac{x^2}{9\pi^2}\right) \left(1 - \frac{x^2}{16\pi^2}\right) \times \dots \end{aligned}$$

01-Oct-2008

MATH 6101

13

Euler's Sum

Now, we can multiply out the product and collect coefficients of like powers.

$$1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \frac{x^6}{7!} + \dots = 1 - \left(\frac{1}{\pi^2} + \frac{1}{4\pi^2} + \frac{1}{9\pi^2} + \frac{1}{16\pi^2} + \dots\right) x^2 + \dots$$

Equating the coefficients of x^2 , we get

$$\begin{aligned} -\frac{1}{3!} &= -\left(\frac{1}{\pi^2} + \frac{1}{4\pi^2} + \frac{1}{9\pi^2} + \frac{1}{16\pi^2} + \dots\right) \\ \frac{\pi^2}{6} &= 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \dots = \sum_{n=1}^{\infty} \frac{1}{n^2} \end{aligned}$$

01-Oct-2008

MATH 6101

14

Euler's Sum

Euler did not stop there. He had a good thing going.

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{n^4} &= \frac{\pi^4}{90} \\ \sum_{n=1}^{\infty} \frac{1}{n^6} &= \frac{\pi^6}{945} \end{aligned}$$

In 1744 he obtained

$$\sum_{n=1}^{\infty} \frac{1}{n^{26}} = \frac{2^{24} 76977927 \pi^{26}}{27!}$$

01-Oct-2008

MATH 6101

15

Euler's Trigonometric Sums

Euler took what we knew for real numbers and pushed it to complex numbers.

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

and replace x by the complex number

$$z = a(\cos \varphi + i \sin \varphi)$$

$$\frac{1}{1-z} = 1 + z + z^2 + z^3 + \dots$$

$$\frac{1}{1-a(\cos \varphi + i \sin \varphi)} = 1 + [a(\cos \varphi + i \sin \varphi)] + [a(\cos \varphi + i \sin \varphi)]^2 + [a(\cos \varphi + i \sin \varphi)]^3 + \dots$$

01-Oct-2008 MATH 6101 16

Euler's Trigonometric Sums

Euler chose to look at the case $a = -1$. Proceed by rationalizing the denominator

$$\frac{1}{1 + (\cos \varphi + i \sin \varphi)} = \frac{1}{1 + (\cos \varphi + i \sin \varphi)} \frac{1 + \cos \varphi - i \sin \varphi}{1 + \cos \varphi - i \sin \varphi}$$

$$= \frac{1 + \cos \varphi - i \sin \varphi}{(1 + \cos \varphi)^2 + \sin^2 \varphi} = \frac{1 + \cos \varphi - i \sin \varphi}{2(1 + \cos \varphi)}$$

$$= \frac{1}{2} - \frac{i \sin \varphi}{2(1 + \cos \varphi)}$$

01-Oct-2008 MATH 6101 17

Euler's Trigonometric Sums

On the right hand side, use De Moivre's Theorem: $(\cos \varphi + i \sin \varphi)^k = \cos k\varphi + i \sin k\varphi$

$$1 - (\cos \varphi + i \sin \varphi) + (\cos \varphi + i \sin \varphi)^2 - (\cos \varphi + i \sin \varphi)^3 + \dots$$

$$= 1 - (\cos \varphi + i \sin \varphi) + (\cos 2\varphi + i \sin 2\varphi) - (\cos 3\varphi + i \sin 3\varphi) + \dots$$

$$= (1 - \cos \varphi + \cos 2\varphi - \cos 3\varphi + \dots) - i(\sin \varphi - \sin 2\varphi + \sin 3\varphi + \dots)$$

Equating the real parts of the equation gives:

$$\frac{1}{2} = 1 - \cos \varphi + \cos 2\varphi - \cos 3\varphi + \cos 4\varphi - \cos 5\varphi + \dots$$

$$\cos \varphi - \cos 2\varphi + \cos 3\varphi - \cos 4\varphi + \cos 5\varphi - \dots = \frac{1}{2}$$

01-Oct-2008 MATH 6101 18

Euler's Trigonometric Sums

Integrate with respect to φ .

$$\sin \varphi - \frac{1}{2} \sin 2\varphi + \frac{1}{3} \sin 3\varphi - \frac{1}{4} \sin 4\varphi + \frac{1}{5} \sin 5\varphi - \dots = \frac{\varphi}{2} + C$$

Since $\sin(0) = 0$, we have $C = 0$ and

$$\sin \varphi - \frac{1}{2} \sin 2\varphi + \frac{1}{3} \sin 3\varphi - \frac{1}{4} \sin 4\varphi + \frac{1}{5} \sin 5\varphi - \dots = \frac{\varphi}{2}$$

Integrate again with respect to φ .

$$\cos \varphi - \frac{1}{4} \cos 2\varphi + \frac{1}{9} \cos 3\varphi - \frac{1}{16} \cos 4\varphi + \frac{1}{25} \cos 5\varphi - \dots = -\frac{\varphi^2}{4} + C$$

Since $\cos(0) = 1$, we have

01-Oct-2008 MATH 6101 19

Euler's Trigonometric Sums

$$C = 1 - \frac{1}{4} + \frac{1}{9} - \frac{1}{16} + \frac{1}{25} - \dots = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2}$$

What is C ? Is it related to the previous sum?

Use the fact that $\cos(2\pi + \alpha) = \cos \alpha$, make the substitution $\varphi = \pi/2$ and we get

$$C - \frac{(\pi/2)^2}{4} = \cos \frac{\pi}{2} - \frac{1}{4} \cos \pi + \frac{1}{9} \cos \frac{3\pi}{2} - \frac{1}{16} \cos 4\pi + \frac{1}{25} \cos \frac{5\pi}{2} - \dots$$

$$= 0 + \frac{1}{4} + 0 - \frac{1}{16} + 0 + \frac{1}{36} - \dots$$

$$= \frac{1}{4} \left(1 - \frac{1}{4} + \frac{1}{9} - \frac{1}{16} + \dots \right) = \frac{C}{4}$$

01-Oct-2008 MATH 6101 20

Euler's Trigonometric Sums

$$C - \frac{(\pi/2)^2}{4} = \frac{C}{4}$$

$$\frac{3C}{4} = \frac{(\pi/2)^2}{4}$$

$$C = \frac{\pi^2}{12}$$

You can use this process to show the previous sum, as well.

01-Oct-2008 MATH 6101 21

Euler's Trigonometric Sums

Note that from our earlier results, for $x \in (\pi, -\pi)$.

$$x = 2 \left(\sin x - \frac{1}{2} \sin 2x + \frac{1}{3} \sin 3x - \frac{1}{4} \sin 4x + \frac{1}{5} \sin 5x - \dots \right)$$

$$x^2 = 4 \left(\frac{\pi^2}{12} - \cos x - \frac{1}{4} \cos 2x + \frac{1}{9} \cos 3x - \frac{1}{16} \cos 4x + \frac{1}{25} \cos 5x - \dots \right)$$

This ability to represent functions as trigonometric series languished until the work of Fourier in 1807 when he showed that these trigonometric series could be used to model the propagation of heat through a material.

01-Oct-2008

MATH 6101

22

Euler's Trigonometric Sums

Note:

$$x = 2 \left(\sin x - \frac{1}{2} \sin 2x + \frac{1}{3} \sin 3x - \frac{1}{4} \sin 4x + \frac{1}{5} \sin 5x - \dots \right)$$

So

$$\frac{\pi}{3} = 2 \left(\sin \left(\frac{\pi}{3} \right) - \frac{1}{2} \sin \left(\frac{2\pi}{3} \right) + \frac{1}{3} \sin \left(\frac{3\pi}{3} \right) - \frac{1}{4} \sin \left(\frac{4\pi}{3} \right) + \frac{1}{5} \sin \left(\frac{5\pi}{3} \right) - \frac{1}{6} \sin \left(\frac{6\pi}{3} \right) + \frac{1}{7} \sin \left(\frac{7\pi}{3} \right) - \dots \right)$$

$$\frac{\pi}{3} = 2 \left(\frac{\sqrt{3}}{2} - \frac{1\sqrt{3}}{2 \cdot 2} + \frac{1}{3} \cdot 0 - \frac{1}{4} \left(-\frac{\sqrt{3}}{2} \right) + \frac{1}{5} \left(\frac{\sqrt{3}}{2} \right) - \frac{1}{6} \cdot 0 + \frac{1\sqrt{3}}{7 \cdot 2} - \dots \right)$$

$$\frac{\pi}{3} = \sqrt{3} - \sqrt{3} \frac{1}{2} + \sqrt{3} \frac{1}{4} - \sqrt{3} \frac{1}{5} + \sqrt{3} \frac{1}{7} - \dots$$

$$\frac{\pi}{3\sqrt{3}} = 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{5} + \frac{1}{7} - \dots$$

01-Oct-2008

MATH 6101

23

Your Turn

Show:

$$\frac{2\pi}{3\sqrt{3}} = 1 + \frac{1}{2} - \frac{1}{4} - \frac{1}{5} + \frac{1}{7} + \frac{1}{8} - \frac{1}{10} - \frac{1}{11} + \dots$$

01-Oct-2008

MATH 6101

24
