

## MATH 6101 Fall 2008

Functions, Sequences and Limits

---

---

---

---

---

---

---

---

## The Topology of the Reals

We will make some simple definitions. Let  $a$  and  $b$  be any two real numbers with  $a < b$ .

$$(a,b) = \{x \in \mathbb{R} \mid a < x < b\}$$

$$[a,b] = \{x \in \mathbb{R} \mid a \leq x \leq b\}$$

$$(a,b] = \{x \in \mathbb{R} \mid a < x \leq b\}$$

$$[a,b) = \{x \in \mathbb{R} \mid a \leq x < b\}$$

$$(a,\infty) = \{x \in \mathbb{R} \mid a < x\}$$

$$[a,\infty) = \{x \in \mathbb{R} \mid a \leq x\}$$

$$(-\infty,b) = \{x \in \mathbb{R} \mid x < b\}$$

$$(-\infty,b] = \{x \in \mathbb{R} \mid x \leq b\}$$

22-Oct-2008

MATH 6101

2

---

---

---

---

---

---

---

---

## Topology of the Reals

If  $r \in \mathbb{R}$  then a *neighborhood* of  $r$  is an open interval  $(a,b)$  so that  $r \in (a,b)$ .

The neighborhood is *centered* at  $r$  if

$$r = (a + b)/2$$

If  $\varepsilon$  and  $a$  are reals, then the  $\varepsilon$ -*neighborhood* of  $a$  is the interval  $(a - \varepsilon, a + \varepsilon)$

22-Oct-2008

MATH 6101

3

---

---

---

---

---


---

---

---

## Functions

Nicole Oresme – 1350 – described the laws of nature as laws giving a dependence of one quantity on another.



22-Oct-2008

MATH 6101

4

---

---

---

---

---


---

---

---

## History of Function

Galileo – 1638 – studies of motion contain the clear understanding of a relation between variables



22-Oct-2008

MATH 6101

5

---

---

---

---

---

---

---

---

## History of Function

Descartes - an equation in two variables, geometrically represented by a curve, indicates a dependence between variable quantities



22-Oct-2008

MATH 6101

6

---

---

---

---

---

---

---

---

## Euclid's Rational Numbers

Newton – showed how functions arise from infinite power series

Leibniz – 1673 – the first to use the term *function*. He took function to designate, in very general terms, the dependence of geometrical quantities on the shape of a curve.



22-Oct-2008

MATH 6101

7

---

---

---

---

---

---

---

---

## History of Function

- Jean Bernoulli - 1718 - function of a variable as a quantity that is composed in some way from that variable and constants
- Euler – 1748 - A function of a variable quantity is an analytic expression composed in any way whatsoever of the variable quantity and numbers or constant quantities.
- Euler – 1755 - If some quantities so depend on other quantities that if the latter are changed the former undergoes change, then the former quantities are called functions of the latter.

22-Oct-2008

MATH 6101

8

---

---

---

---

---

---

---

---

## History of Function

- Cauchy – 1821 – still thinking of a function in terms of a formula (either explicit or implicit)
- Fourier – 1822 – introduced general Fourier series but fell back on old definitions
- Dirichlet – 1837 – defined general function and continuity (in modern terms)
- Weierstrauss – 1885 – any continuous function is the limit of a uniformly convergent sequence of polynomials
- Goursat – 1923 – modern definition

22-Oct-2008

MATH 6101

9

---

---

---

---

---

---

---

---

### Definitions

Bernoulli – 1718 – *One calls here a function of a variable a quantity composed in any manner whatever of this variable and constants.*

Basically this meant +, -, ×, ÷, √, logs and sines.

They would say that  $f(x)$  depended *analytically* on the variable  $x$ .

22-Oct-2008

MATH 6101

10

---

---

---

---

---

---

---

---

### Definitions

Fourier – 1822 – *In general the function  $f(x)$  represents a succession or ordinates each of which is arbitrary. An infinity of values being given to the abscissa  $x$ , there are an equal number of ordinates  $f(x)$ . All have actual numerical values, either positive or negative or null. We do not suppose these ordinates to be subject to common law; they succeed each other in any manner whatever, and each of them is given as it were a single quantity.*

22-Oct-2008

MATH 6101

11

---

---

---

---

---

---

---

---

### Definitions

Fourier removed the requirement of “analytic” from the definition. It was not widely accepted for years.

22-Oct-2008

MATH 6101

12

---

---

---

---

---

---

---

---

## Definitions

Dirichlet – 1837 – *Let us suppose that  $a$  and  $b$  are two definite values and  $x$  is a variable quantity which is to assume, gradually, all values located between  $a$  and  $b$ . Now, if to each  $x$  there corresponds a unique, finite  $y$  ..., then  $y$  is called a ...function of  $x$  for this interval. It is, moreover, not at all necessary, that  $y$  depends on  $x$  in this whole interval according to the same law; indeed, it is not necessary to think of only relations that can be expressed by mathematical operations.*

22-Oct-2008

MATH 6101

13

---

---

---

---

---

---

---

---

---

---

## Definitions

Every “Bernoulli” function is a “Fourier” or a “Dirichlet” function.

Dirichlet:

$$f(x) = \begin{cases} 1 & \text{if } x \text{ is rational and } 0 \leq x \leq 1 \\ 0 & \text{if } x \text{ is irrational and } 0 \leq x \leq 1 \end{cases}$$

22-Oct-2008

MATH 6101

14

---

---

---

---

---

---

---

---

---

---

## Another “Bad Example”

d’Alembert was working on the problem of describing a vibrating string. The initial position for the string is not the graph of any analytical expression.



22-Oct-2008

MATH 6101

15

---

---

---

---

---

---

---

---

---

---

## A More Modern Definition

Let  $D$  be a set of real numbers. A function  
 $f: D \rightarrow \mathbb{R}$   
 is a rule that assigns a number  $f(x)$  to every  
 element  $x$  of  $D$ .

22-Oct-2008

MATH 6101

16

---

---

---

---

---

---

---

---

## Modern Set Theory Definition

A function  $f$  is an ordered triple of sets  $(F, X, Y)$  with  
 restrictions, where  $F$  (the **graph**) is a set of ordered pairs  $(x, y)$ ,  $X$   
 (the **source**) contains all the first elements of  $F$  and perhaps  
 more, and  $Y$  (the **target**) contains all the second elements of  $F$   
 and perhaps more.

The most common restrictions are that  $F$  pairs each  $x$  with  
 just one  $y$ , and that  $X$  is just the set of first elements of  $F$  and no  
 more.

When *no* restrictions are placed on  $F$ , we speak of a *relation*  
 between  $X$  and  $Y$  rather than a function. The relation is "single-  
 valued" when the first restriction holds:  $(x, y_1) \in F$  and  $(x, y_2) \in F$   
 together imply  $y_1 = y_2$ .

Relations that are not single valued are sometimes called  
*multivalued* functions. A relation is *total* when a second  
 restriction holds: if  $x \in X$  then  $(x, y) \in F$  for some  $y$ . Thus we can  
 also say that

A function from  $X$  to  $Y$  is a single-valued, total relation  
 between  $X$  and  $Y$ .

22-Oct-2008

MATH 6101

17

---

---

---

---

---

---

---

---

## Sequences

Let  $\mathbb{N}$  = the set of natural numbers (it will not  
 matter if it starts with 0 or with 1).

A sequence is a function  $a: \mathbb{N} \rightarrow \mathbb{R}$ .

We will normally denote a sequence by its set  
 of outputs  $\{a_n\}$ , where  $a_n = a(n)$ .

Occasionally you will see  $a_0, a_1, a_2, a_3, \dots$  or  
 $\{a_n\}_{n=0}^{\infty}$

22-Oct-2008

MATH 6101

18

---

---

---

---

---

---

---

---

### Examples

- 1)  $\{1,2,3,4,5,6,\dots\}$  – an arithmetic progression  
 $(f(n) = n)$
- 2)  $\{a + bn \mid n=0,1,2,3,\dots\}$  – a different type of arithmetic progression –  $(f(n) = a + bn)$
- 3)  $\{a^0, a^1, a^2, a^3, a^4, \dots\}$  – a geometric progression  
 $(f(n) = a^n)$
- 4)  $\{1, 1/2, 1/3, 1/4, 1/5, \dots\}$  –  $(f(n) = 1/n)$
- 5)  $f(n) = a_n = (-1)^n$ . Note that the range is  $\{-1, 1\}$

22-Oct-2008 MATH 6101 19

---

---

---

---

---

---

---

---

### Examples

- 1)  $f(n) = a_n = \cos(\pi n / 3)$   
 $a_1 = \cos(\pi/3) = \cos 60^\circ = 1/2$   
 $\{a_n\} = \{1/2, -1/2, -1, -1/2, 1/2, 1, 1/2, -1/2, -1, -1/2, 1/2, 1, \dots\}$ . The function takes on only a finite number of values, but the sequence has an infinite number of elements.
- 2)  $f(n) = a_n = n^{1/n}$   
 $\{1, 2^{1/2}, 3^{1/3}, 4^{1/4}, \dots\} = \{1, 1.41421, 1.44225, 1.41421, 1.37973, 1.34801, 1.32047, 1.29684, 1.27652, 1.25893, \dots\}$   
 Also  $a_{100} = 1.04713$ ,  $a_{10,000} = 1.00092$
- 3)  $b_n = (1+1/n)^n$   
 $\{2, (3/2)^2, (4/3)^3, (5/4)^4, \dots\} = \{2, 2.25, 2.37037, 2.44141, 2.48832, 2.52163, 2.54650, 2.56578, 2.58117, 2.59374, \dots\}$   
 Also  $a_{100} = 2.74081$  and  $a_{10,000} = 2.71815$

22-Oct-2008 MATH 6101 20

---

---

---

---

---

---

---

---

### Almost all ...

**Definition:** It is said that *almost all the terms* of the sequence  $\{a_n\}$  have a certain property provided that there is an index  $N$  such that  $\{a_n\}$  possesses this property whenever  $n \geq N$ .

22-Oct-2008 MATH 6101 21

---

---

---

---

---

---

---

---

## Convergence

**Definition 1:** A sequence of real numbers is said to *converge* to a real number  $L$  if for every  $\varepsilon > 0$  there is an integer  $N > 0$  such that if  $k > N$  then  $|a_k - L| < \varepsilon$ .

**Definition 2:** A sequence of real numbers is said to *converge* to a real number  $L$  if every neighborhood of  $L$  contains almost all of the terms of  $\{a_n\}$ .

The number  $L$  is called the *limit* of the sequence.

22-Oct-2008

MATH 6101

22

---

---

---

---

---

---

---

---

## Convergence

**Lemma 1:** The sequence  $\{1/n\}$  converges to 0.

**Proof:** Let  $(a, b)$  be any neighborhood of 0. This means that  $a < 0 < b$ . Let  $N > [1/b]$ , be an integer greater than  $1/b$ . Then  $1/N < b$  and for every integer  $n > N$ , we have that

$$a < 0 < 1/n < 1/N < b$$

and  $(a, b)$  contains almost all of the elements of the sequence. Thus, the sequence converges to 0.

22-Oct-2008

MATH 6101

23

---

---

---

---

---

---

---

---

## Convergence

**Lemma 1:** The sequence  $\{1/n\}$  converges to 0.

**Proof:** You prove this using Definition 1.

22-Oct-2008

MATH 6101

24

---

---

---

---

---

---

---

---



## Convergence

**Definition:** A sequence is *convergent* if it has a limit. If it is not convergent it is called *divergent*.

**Lemma 2:** The sequence  $\{a_n\}$  converges to  $L$  if and only if every neighborhood of  $L$  that is centered at  $L$  contains almost all of the terms of the sequence.

Note that this tells us that the two definitions are the same.

22-Oct-2008

MATH 6101

25

---

---

---

---

---

---

---

---

---

---

## Example

Let  $a_n = n/2^n$ .  $\{a_n\} = \{1/2, 2/2^2, 3/2^3, 4/2^4, \dots\}$

Educated guess:  $\{a_n\} \rightarrow 0$ .

Let  $\epsilon = 0.1, 0.01, 0.001, 0.0001, 0.00001$ .

We need to find an integer  $N$  so that  $|n/2^n - 0| < \epsilon$

$\epsilon$	$N$
1	$N > 0$
0.1	$N > 5$
0.01	$N > 9$
0.001	$N > 14$
0.0001	$N > 18$
0.00001	$N > 22$

Look in the table of values. Note that for  $N = 6$  the above is true if  $\epsilon = 0.1$

22-Oct-2008

MATH 6101

26

---

---

---

---

---

---

---

---

---

---

## **Theorem**(Convergent sequences are bounded)

Let  $\{a_n\}$  be a convergent sequence. Then the sequence is bounded, and the limit is unique.

### Proof:

(i) Uniqueness: Suppose the sequence has two limits,  $L$  and  $K$ . Let  $\epsilon > 0$ . There is an integer  $N_K$  such that  $|a_n - K| < \epsilon/2$  if  $n > N_K$ .

Also, there is an integer  $N_L$  such that  $|a_n - L| < \epsilon/2$  if  $n > N_L$ .

By Triangle Inequality:

$$|L - K| < |a_n - L| + |a_n - K| < \epsilon/2 + \epsilon/2 = \epsilon$$

if  $n > \max\{N_K, N_L\}$ .

Therefore  $|L - K| < \epsilon$  for any  $\epsilon > 0$ . But this means that  $L = K$ .

22-Oct-2008

MATH 6101

27

---

---

---

---

---

---

---

---

---

---

**Theorem** (Convergent sequences are bounded)

**Proof:**

(ii) Boundedness. Since the sequence converges, choose any  $\varepsilon > 0$ . Specifically take  $\varepsilon = 1$ . There is  $N$  so that

$$|a_n - L| < 1 \text{ if } n > N.$$

Fix  $N$ . Then

$$|a_n| \leq |a_n - L| + |L| < 1 + |L| = P \text{ for all } n > N.$$

Let  $M = \max\{|a_1|, |a_2|, \dots, |a_N|, P\}$ . Thus  $|a_n| < M$  for all  $n$ , which makes the sequence bounded.

22-Oct-2008

MATH 6101

28

---

---

---

---

---

---

---

---

---

---

**Theorem:** If  $\{a_n\} \rightarrow L$ ,  $\{b_n\} \rightarrow M$  and  $\alpha$  is a real number, then

1.  $\lim_{n \rightarrow \infty} \alpha = \alpha$ .
2.  $\lim_{n \rightarrow \infty} (a_n \pm b_n) = L \pm M$
3.  $\lim_{n \rightarrow \infty} (a_n \times b_n) = L \times M$
4.  $\lim_{n \rightarrow \infty} (\alpha a_n) = \alpha L$
5. If  $a_n \leq b_n$  for all  $n \geq m$ , then  $L \leq M$
6. If  $b_n \neq 0$  for all  $n$  and if  $M \neq 0$ , then  $\text{glb}\{|b_n|\} > 0$ .
7.  $\lim_{n \rightarrow \infty} (a_n/b_n) = L/M$ , provided  $M \neq 0$ .

22-Oct-2008

MATH 6101

29

---

---

---

---

---

---

---

---

---

---

**Proof:**

1.  $\lim_{n \rightarrow \infty} \alpha = \alpha$

Since  $\alpha - \alpha = 0$ , for any  $\varepsilon > 0$ ,  $|\alpha - \alpha| < \varepsilon$  and we are done.

2.  $\lim_{n \rightarrow \infty} (a_n \pm b_n) = L \pm M$

Do this for the sum. The difference is similar. Let  $\varepsilon > 0$ , there exist  $N_a$  and  $N_b$  so that

$$|a_n - L| < \varepsilon/2 \text{ if } n > N_a \text{ and}$$

$$|b_n - M| < \varepsilon/2 \text{ if } n > N_b.$$

Let  $K = \max\{N_a, N_b\}$ , then if  $n > K$

$$|(a_n + b_n) - (L + M)| \leq |a_n - L| + |b_n - M| < \varepsilon/2 + \varepsilon/2 = \varepsilon$$

22-Oct-2008

MATH 6101

30

---

---

---

---

---

---

---

---

---

---

$$3. \lim_{n \rightarrow \infty} (a_n \times b_n) = L \times M$$

Note:

$$\begin{aligned} |(a_n b_n) - (LM)| &\leq |(a_n - L) b_n + L(b_n - M)| \\ &\leq |(a_n - L) b_n| + |L(b_n - M)| \\ &= |(a_n - L)| |b_n| + |L| |b_n - M| \end{aligned}$$

Then use the fact that  $\{b_n\}$  is bounded.

$$4. \lim_{n \rightarrow \infty} (a a_n) = aL$$

Consider  $\varepsilon/a$  if  $a \neq 0$ . If  $a = 0$  this is easy.

22-Oct-2008

MATH 6101

31

---

---

---

---

---

---

---

---

---

---

5. If  $a_n \leq b_n$  for all  $n \geq m$ , then  $L \leq M$

6. If  $b_n \neq 0$  for all  $n$  and if  $M \neq 0$ , then  $\text{glb}\{b_n\} > 0$ .

Let  $\varepsilon = |M|/2 > 0$ .  $\{b_n\} \rightarrow M$  so there is  $N$  so that if  $n > N$  then  $|b_n - M| < |M|/2$ .

So if  $n > N$  we must have  $|b_n| \geq |M|/2$ .

If not by the Triangle Inequality

$$\begin{aligned} |M| &= |M - b_n + b_n| \leq |M - b_n| + |b_n| \\ &< |M|/2 + |M|/2 = |M| \end{aligned}$$

So set

$$m = \min \{|M|/2, |b_1|, |b_2|, \dots, |b_N|\}.$$

Then  $m > 0$  and  $|b_n| \geq m$  for all  $n$

22-Oct-2008

MATH 6101

32

---

---

---

---

---

---

---

---

---

---

7.  $\lim_{n \rightarrow \infty} (a_n/b_n) = L/M$ , provided  $M \neq 0$ .

Reduce to  $\lim_{n \rightarrow \infty} (1/b_n) = 1/M$  – How?

Let  $\varepsilon > 0$ . By (6) there is  $m > 0$  so that  $|b_n| \geq m$ . Since  $\{b_n\}$  is convergent there is  $N$  so that if  $n > N$

$$|M - b_n| < \varepsilon m |M|$$

Then for  $n > N$

$$\begin{aligned} |1/b_n - 1/M| &= |b_n - M|/|b_n M| \\ &\leq |b_n - M|/(m|M|) < \varepsilon \end{aligned}$$

22-Oct-2008

MATH 6101

33

---

---

---

---

---

---

---

---

---

---

### Example

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{3n^4 + 4n^3 - 7n^2 - 5280n + 3216547}{7n^4 + 5588741226n^2 - 7} \\ &= \lim_{n \rightarrow \infty} \frac{3 \frac{n^4}{n^4} + 4 \frac{n^3}{n^4} - 7 \frac{n^2}{n^4} - 5280 \frac{n}{n^4} + 3216547 \frac{1}{n^4}}{7 \frac{n^4}{n^4} + 5588741226 \frac{n^2}{n^4} - 7 \frac{1}{n^4}} \\ &= \lim_{n \rightarrow \infty} \frac{3 + \frac{4}{n} - \frac{7}{n^2} - \frac{5280}{n^3} + \frac{3216547}{n^4}}{7 + \frac{5588741226}{n^2} - \frac{7}{n^4}} = \frac{3}{7} \end{aligned}$$

22-Oct-2008

MATH 6101

34

---

---

---

---

---

---

---

---

---

---

### The Squeeze Theorem

**Theorem:** If  $\{a_n\} \rightarrow L$ ,  $\{b_n\} \rightarrow L$  and  $a_n \leq c_n \leq b_n$  for all  $n \geq m$ . Then  $\{c_n\} \rightarrow L$ .

22-Oct-2008

MATH 6101

35

---

---

---

---

---

---

---

---

---

---

### The Power Theorem

**Theorem:** Let  $a$  be fixed. Then

$$\lim_{n \rightarrow \infty} a^n = \begin{cases} 0 & \text{if } |a| < 1 \\ 1 & \text{if } a = 1 \\ dne & \text{if } |a| > 1 \\ dne & \text{if } a = -1 \end{cases}$$

22-Oct-2008

MATH 6101

36

---

---

---

---

---

---

---

---

---

---

Find

$$\lim_{n \rightarrow \infty} \frac{3n^2 + 5n + 1}{n + 1}$$

22-Oct-2008

MATH 6101

37

---



---



---



---



---



---



---



---

Find

$$\lim_{n \rightarrow \infty} \frac{3n^2 + 5n + 1}{n^2 + 1}$$

22-Oct-2008

MATH 6101

38

---



---



---



---



---



---



---



---

Find

$$\lim_{n \rightarrow \infty} \frac{3n^2 + 5n + 1}{n^3 + 1}$$

22-Oct-2008

MATH 6101

39

---



---



---



---



---



---



---



---

Find

$$\lim_{n \rightarrow \infty} \frac{0.5^n + 3\sin(n)}{\sqrt{n}}$$

22-Oct-2008

MATH 6101

40

---



---



---



---



---



---



---



---

Find

$$\lim_{n \rightarrow \infty} \frac{2^n - 1}{3^n + 1}$$

22-Oct-2008

MATH 6101

41

---



---



---



---



---



---



---



---

Find

$$\lim_{n \rightarrow \infty} \frac{2^n + 1}{3^n - 1}$$

22-Oct-2008

MATH 6101

42

---



---



---



---



---



---



---



---

Find

$$\lim_{n \rightarrow \infty} \frac{3^n + 2^n}{3^n - 2^n}$$

22-Oct-2008

MATH 6101

43

---



---



---



---



---



---



---

Find

$$\lim_{n \rightarrow \infty} \frac{3^n + 4^{n-3}}{5^{n+2} - 2^{n+4}}$$

22-Oct-2008

MATH 6101

44

---



---



---



---



---



---



---

Find

$$\lim_{n \rightarrow \infty} \frac{4^{2n-3} + 2^{5n+6}}{5^{3n-2} - 3^{n+10}}$$

22-Oct-2008

MATH 6101

45

---



---



---



---



---



---



---

Find

$$\lim_{n \rightarrow \infty} n^n$$

22-Oct-2008

MATH 6101

46

---

---

---

---

---

---

---

---

Find

$$\lim_{n \rightarrow \infty} \frac{1}{n^n}$$

22-Oct-2008

MATH 6101

47

---

---

---

---

---

---

---

---

Find

$$\lim_{n \rightarrow \infty} \left( 1 - \left| \frac{\sin(n)}{n} \right| \right)$$

22-Oct-2008

MATH 6101

48

---

---

---

---

---

---

---

---



Find

$$\lim_{n \rightarrow \infty} \frac{n}{2^n}$$

22-Oct-2008

MATH 6101

49

---

---

---

---

---

---

---

---

Find

$$\lim_{n \rightarrow \infty} \frac{n^2}{2^n}$$

22-Oct-2008

MATH 6101

50

---

---

---

---

---

---

---

---

Find

$$\lim_{n \rightarrow \infty} \frac{n!}{n^n}$$

22-Oct-2008

MATH 6101

51

---

---

---

---

---

---

---

---

Find

$$\lim_{n \rightarrow \infty} \frac{n!}{2^n}$$

22-Oct-2008

MATH 6101

52

---

---

---

---

---

---

---

---