

STA 291 Lecture 24

- *Two kinds of Error in Testing hypothesis*
- *Examples.*

About **bonus** project

Due in Lab April 20- 22

- Some survey show a majority people believe in "hot hand". A follow -up question then is: if there were "hot hand", how much better/worse a shooter can become by the previous shoots? (i.e. what is a reasonable difference to expect)
- i.e. whether he made or missed the two previous shots, how much difference do you think this has the effect on the present shoot? (in terms of hitting percentages) 5%? 10%? or even 20%?

- Since a small difference will need more data to detect.
- A larger difference can be discovered with less data.
- Some clue: margin of error calculations.

- Bonus is worth equivalent to one LAB
- Lab will start to give “practical quiz”

Decisions and Types of Errors in Tests of Hypotheses

- Terminology:
 - The alpha-level (significance level) is a *threshold number* such that one rejects the null hypothesis if the p -value is less than or equal to it. The most common alpha-levels are 0.05 and 0.01
 - The choice of the alpha-level reflects how cautious the researcher wants to be (when it come to reject null hypothesis)

Type I and Type II Errors

- Type I Error: The null hypothesis is rejected, even though it is true.
- Type II Error: The null hypothesis is not rejected, even though it is false.
- **Setting the alpha-level low protect us from type I Error. (the probability of making a type I error is less than alpha)**

Type I and Type II Errors

		Decision	
		Reject null	Do not reject null
the null hypothesis	True	Type I error	Correct
	False	Correct	Type II error

STA 291 - Lecture 24

7

Type I and Type II Errors

- Terminology:
 - **Alpha** = Probability of make a Type I error
 - **Beta** = Probability of make a Type II error
 - **Power** = 1 – Probability of a Type II error = 1 - Beta
- For a given data, the smaller the probability of Type I error, the larger the probability of Type II error and the smaller the power
- i.e. If you set alpha very small, it is more likely that you fail to detect a real difference (larger Beta).

STA 291 - Lecture 24

8

- When sample size(s) increases, both error probabilities could be made to decrease.
- Our Strategy:
 - keep type I error probability small by pick a small alpha.
 - Increase sample size to make Beta small.
- Depend on how expensive to obtain data, a Beta = 0.15 is not uncommon.

STA 291 - Lecture 24

9

Type I and Type II Errors

- In practice, alpha is specified, and the probability of Type II error could be calculated, but the calculations are usually difficult (sample size calculation)
- **How to choose alpha?**
- If the consequences of a Type I error are very serious, then choose a smaller alpha, like 0.01.
- For example, you want to find evidence that someone is guilty of a crime.
- In exploratory research, often a larger probability of Type I error is acceptable (like 0.05 or even 0.1)

STA 291 - Lecture 24

10

Example: New drug development

- The null hypothesis usually state that the new drug is “no difference” to the placebo.
- A type I error in this context is: falsely conclude a drug is useful when it is actually “NO effect”
- A type II error in this context is: falsely dismiss a useful drug.

STA 291 - Lecture 24

11

Alternative and p-value computation

$$H_0 : p = p_0$$

	One-Sided Tests		Two-Sided Test
alternative Hypothesis	$H_A : p < p_0$	$H_A : p > p_0$	$H_A : p \neq p_0$
p-value	$P(Z < z_{obs})$	$P(Z > z_{obs})$	$2 \cdot P(Z > z_{obs})$

$$z_{obs} = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0) / n}}$$

STA 291 - Lecture 24

12

Example

- Two consumer products (shampoo, laundry detergent etc) comparison. Call them A vs. B
- n consumers are given both products in the identical packaging. After one week of use of both products, state a preference.
- If there were no difference, then we should see 50%-50%

STA 291 - Lecture 24

13

- Suppose in n=236 consumers, 110 prefer product A. Let p = popu. proportion prefer A. Use alpha = 0.05
- Null: $H_0 : p = 0.5$
- Alternative: $H_A : p \neq 0.5$
- Compute z_{obs}

STA 291 - Lecture 24

14

- Sample proportion, $\hat{p} = 110/236 = 0.4661$

$$z_{obs} = \frac{0.4661 - 0.5}{\sqrt{0.5(1-0.5)/236}} = -1.04156$$

- Finally look the Z table for P-value:
- P-value = $2P(Z > 1.04) = 2(1 - 0.8508) = 0.2984$

STA 291 - Lecture 24

15

- Conclusion, we do not reject null hypothesis since P-value is not less than alpha.
- Since 0.2984 is not less than 0.05

STA 291 - Lecture 24 16

Two sample cases are similar, with two differences:

- Hypothesis involve 2 parameters from 2 populations
- Test statistic, z_{obs} , is different, involve 2 samples

STA 291 - Lecture 24 17

Alternative and p-value computation			
$H_0 : p_1 - p_2 = 0$			
	One-Sided Tests		Two-Sided Test
alternative Hypothesis	$H_A : p_1 - p_2 < 0$	$H_A : p_1 - p_2 > 0$	$H_A : p_1 - p_2 \neq 0$
p-value	$P(Z < z_{obs})$	$P(Z > z_{obs})$	$2 \cdot P(Z > z_{obs})$

STA 291 - Lecture 24 18

Two p's

$H_0 : p_1 = p_2$ which is equivalent to $H_0 : p_1 - p_2 = 0$,

$$z_{obs} = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\frac{\hat{p}(1-\hat{p})}{n_1} + \frac{\hat{p}(1-\hat{p})}{n_2}}}$$

STA 291 - Lecture 24

19

- Where the \hat{p} in the denominator is the combined (pooled) sample proportion.
= Total number of successes over total number of observations

So there are 3 different sample proportions: from sample one, from sample two and from both samples.

STA 291 - Lecture 24

20

- P for P-value in a test hypothesis setting
- p for population proportion
- \hat{p} for sample proportion
- p_0 for the hypothesized population proportion value

STA 291 - Lecture 24

21

Attendance Survey Question 24

- On a 4"x6" index card
 - Please write down your name and section number
 - Today's Question:
 - Probability of making a type II error is denoted by:
 - a. Alpha b. Beta c. Power

STA 291 - Lecture 24

22

Example: compare 2 proportions

- A nation wide study: an aspirin every other day can sharply reduce a man's risk of heart attack. (New York Times, reporting Jan. 27, 1987)
- Aspirin group: 104 Heart Att. in 11037
- Placebo group: 189 Heart Att. in 11034
- Randomized, double-blinded study

STA 291 - Lecture 24

23

Example – cont.

- Let aspirin = group 1; placebo = group 2
 - p_1 = popu. proportion of Heart att. for group 1
 - p_2 = popu. proportion of Heart att. for group 2

$H_0 : p_1 = p_2$ which is equivalent to $H_0 : p_1 - p_2 = 0$

$H_A : p_1 \neq p_2$ or $H_A : p_1 - p_2 \neq 0$

STA 291 - Lecture 24

24

Example – cont.

- We may use software to compute a p-value
 - p-value = $7.71 \times 10^{-7} = 0.000000771$
- Or we can calculate by hand:

$$z_{obs} = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\frac{\hat{p}(1-\hat{p})}{n_1} + \frac{\hat{p}(1-\hat{p})}{n_2}}}$$

STA 291 - Lecture 24

25

Example – cont.

- $n_1 = 11037, n_2 = 11034$
 $\hat{p}_1 = 104/11037 = 0.00942285$
 $\hat{p}_2 = 189/11034 = 0.01712887$
 $\hat{p} = (104 + 189)/(11037 + 11034) = 0.013275$
 $z = -0.00770602/0.001540777$
 $= -5.001386$

STA 291 - Lecture 24

26

$$z_{obs} = \frac{0.00942285 - 0.01712887}{\sqrt{\frac{\hat{p}(1-\hat{p})}{11037} + \frac{0.013275(1-0.013275)}{11034}}}$$

STA 291 - Lecture 24

27

Example – cont.

- P-value= $2 \times P(Z > |-5.00|)$
- It falls out of the range of our Z- table, so.....

P-value is approx. zero. (much smaller than 0.0000?)

What is alpha level? Say it was 0.01. Since P-value is smaller than alpha, we reject the null hypothesis.
