STA 291

Lecture 28

• Final exam: CB 106 May 6 Thursday 6:00pm – 8:00pm

STA 291 - Lecture 28

Exam II curve: conversion formula

- If your original score is 83 or above, then converted score = 90+(x-83)10/17
- If your original score is $71 \rightarrow 82$, then converted score = 80 + (x - 71)9/11
- If your original score is $59 \rightarrow 70$, then converted score = 70 + (x - 59)9/11
- If your original score is $48 \rightarrow 58$, then converted score = 60 + (x - 48)9/10
- If your original score is $1 \rightarrow 47$, then converted score = $x \frac{59}{47}$ 2

STA 291 - Lecture 27


Final Exam, Thursday, May 6

- When: 6:00PM 8:00PM
- Where: CB 106

Make-up exam:

- Only by prior arrangement:
- Friday May 7, 10:00am 12:00noon
- come to 8th floor POT for room assignment

STA 291 - Lecture 28



New materials after Exam II


- Chapter 11: 11.1 11.9
- Chapter 12: 12.5
- Chapter 13: 13.1 13.3
- Chapter 14: 14.1, 14.2, 14.3, 14.4

Final Exam

STA 291 - Lecture 28

5

- You bring a calculator.
- Will be given a formula sheet/table
- Any technology that can receive/transmit information wirelessly is *not* permitted during the exam
- Turn off cell phone etc.



Introduction to Linear Models -- a preview of What's next in statistics

- More Confidence intervals and
- More Testing hypothesis

plus

 Statistical models for more complex setting (beyond one/two samples)

More than two samples

STA 291 - Lecture 28

8


9

• Three groups (three populations): no drug, 10mg/day, 20mg/day

35 subjects in each group. Do blood pressure decrease as dosage increase?

We postulate a *Model*: the average decrease of blood pressure islinearly related to the dosage.


In this example, this assumes the effect must be doubled in the 20mg group compared to 10mg group



Example of a linear model $m = b_0 - 1.3 \times (dosage)$

- Here b_0 is the placebo effect, those taking a pill of corn starch, will decrease blood pressure by b_0
- Group with 10mg will further decrease
 13=1.3x10
- Group with 20mg will further decrease 26=1.3x20

STA 291 - Lecture 28



- The number -1.3 in the previous page is just for example. In reality we need to estimate that number (called slope). (estimator mostly produced by a computer, we fed data into a program...)
- Construct confidence interval for the slope
- Test hypothesis that slope =0; -- equivalent to testing no effect of drug

STA 291 - Lecture 28

13

14

15

• The parameters here: • Slope (-1.3) and the intercept (b_0) Confidence interval/testing for these two


parameters, etc


STA 291 - Lecture 28

Models can be more and more complicated....


- But we always test a hypothesis about the parameters in the model.
- Construct a confidence interval about (an all important slope of) the model.
- Once the model is established, it can be very useful: use the model to predict......

STA 291 - Lecture 28









- Two player may have two different curve, but both decreasing.
- Is there a favorable spot of shooting?

more examples:

• Kentucky Utility need to predict the electricity usage, as how it changes as temperature change.

STA 291 - Lecture 28

19

20

Financial modeling

- No matter how complicated the model is,
- No matter how many parameters it involves,
- the correspondence between confidence interval and p-value of testing hypothesis is always there

STA 291 - Lecture 28

Today's Q

- Your name/ID and 291 section number
- A: Sta291 is my last Statistics course in college.
- B: I will be taking another statistics course.
 (Which one?)

STA 291 - Lecture 28