	_
Logic and Proof	
Logic and Proof	
On my first day of school my	
parents dropped me off at the wrong nursery. There I	
wassurrounded by trees and	
bushes!	
26-Aug-2011 MA 341 001 2	
	1
Requirements for Proof	
Mutual understanding of the words and symbols used	
Acceptance of certain statements called axioms without justification	
3. Agreement on how and when one	
statement follows logically from another, that is, agreement on certain rules of	
reasoning.	
26-Aug-2011 <b>MA 341 001</b> 3	

	-
Sets	
A set is a collection of objects satisfying some condition.	
It is possible to have no objects in a set. This set is called the <b>empty set</b> and is	
denoted by { } or Ø	
26-Aug-2011 MA 341 001 4	
Set Notation	
Roster method: {a,b,c,z}	
Set builder notation: $A = \{ x \mid P(x) \text{ is satisfied } \}$	
Set elements:	
αεΑ α is in Α	
a is a member of A	
a is an element of A	
26-Aug-2011 MA 341 001 5	
Set Notation	
Set elements:	
a $\not\in$ A means a is not a member of A $A \subset B$ means that A is a subset of B which	
means every element of A is also an element of B	
if a $\epsilon$ A then a $\epsilon$ B	

Note:  $\varnothing \subset A$  for every set A

26-Aug-2011

Set	Equa	li+\
3e1		.1113

A = B means that every element of A is also and element of B and vice versa.

$$A = B \iff A \subset B \text{ and } B \subset A$$

26-Aug-2011

MA 341 001

### Set Intersection

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\}$$

Set Union

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\}$$

26-Aug-2011

MA 341 001

# Set Complement

$$A^{c} = A' = CA = \{x \mid x \notin A\}$$

Note that

$$\mathbf{A} \cap \mathbf{A}^{c} = \emptyset$$

$$A \cup A^{c} = U$$

26-Aug-2011

	1
Mathematical Statements	
A declarative sentence which is true or	
false, but not both, is called a statement.	
   Examples: 1+1=3	
2 - 1 = 1	
The grass is blue.	
26-Aug-2011 <b>MA 341 00</b> 1 10	
Set Variables	
A variable is a symbol or an icon that can	
be used to represent various elements of the universal set.	
Examples: He is a Wildcat.	
This is true if "he=Adolf Rupp". This is not	
true if "he=Rick Pitino".	
26-Aug-2011 <b>MA 341 00</b> 1 11	
15 / Ng 15 / N	
	1
Logical Connectives & Truth Tables	
If P and Q are statements, then the	
statement <u>P and Q</u> is called the	
conjunction of P and Q.	
Notation: PAQ	
Examples:	
Launipies	
26-Aug-2011 <b>MA 341 001</b> 12	

#### Truth Values

Р	Q	$P \wedge Q$
True	True	True
True	False	False
False	True	False
False	False	False

26-Aug-2011

MA 341 001

## Logical Connectives & Truth Tables

If P and Q are statements, then the statement <u>P or Q</u> is called the **disjunction** of P and Q.

Notation:  $P \vee Q$ 

Note: mathematicians use the inclusive "or" Examples:

26-Aug-2011

MA 341 001

14

## Truth Values

Р	Q	$P \vee Q$
True	True	True
True	False	True
False	True	True
False	False	False

26-Aug-2011

	1
Negation	
If P is a statement, then the statement <u>not</u> <u>P</u> is called the <b>negation</b> of P.	
Notation: ~P (sometimes -P or ¬P)  We assume that either P or ~P is true, but  not both and not neither	
26-Aug-2011 <b>MA 341 001</b> 16	
Negation	
~(P $\wedge$ Q) is equivalent to ~P $\vee$ ~Q.	
~(P ∨ Q) is equivalent to ~P ∧ ~Q ~(~P) is equivalent to P	
( ) jie oquivaleni je i	
26-Aug-2011 <b>MA 341 001</b> 17	
The Conditional	
If P and Q are statements, then the	
statement <u>if P then Q</u> is called the <b>conditional statement</b>	
Notation: $P \Rightarrow Q$	
Examples: If it is cold, it is snowing.	
If it is blue, it is UK.	

26-Aug-2011

#### Truth Values

Р	Q	$P \Rightarrow Q$
True	True	True
True	False	False
False	True	True
False	False	True

See additional notes in Blackboard for further discussion.

26-Aug-2011

MA 341 001

### Truth Values

P = "The animal is a tiger."

Q = "The animal is a mammal."

When must the statement be false?

P	Q	P⇒Q
is a tiger	is a mammal	
is a tiger	is not a mammal	
is not a tiger	is mammal	
is not a tiger	is not a mammal	

MA 341 001

#### Truth Values

Statements of the form "If P then Q" do not mean the same thing in math as in ordinary English.

#### In ordinary English,

- "If P then Q" may suggest order of occurrence.
- "If we go outside, the neighbors will see us" implies that the neighbors will see us **after** we go outside
- If P then Q" can suggest **causation**. This example has the connotation that the neighbors will see us **because** we went outside.

26-Aug-2011

26-Aug-2011

MA 341 001

20

The Conditional	
$P \Rightarrow Q$	
If P then $Q \leftarrow PREFERED$	
Q if P	
P only if Q	
Q provided P	
Q when P	

#### Related Statements

MA 341 001

 $P \Rightarrow Q$ 

Conditional: If P, then Q  $(P \Rightarrow Q)$ Converse: If Q, then  $P(Q \Rightarrow P)$ 

Inverse: If not P, then not Q ( $P \Rightarrow Q$ ) Contrapositive: If not Q, then not P

 $(\sim Q \Rightarrow \sim P)$ 

(Conditional logically equiv to Contrapositive) (Converse logically equiv to Inverse)

26-Aug-2011

26-Aug-2011

MA 341 001

#### Related Statements

If there is smoke, then there is fire. Converse:

If there is fire, then there is smoke. Inverse:

If there is no smoke, then there is no fire. Contrapositive:

If there is no fire, then there is no smoke.

MA 341 001

23

<u> </u>	_		•	
The	Ri	conc	lition	า

If P and Q are statements, then the statement <u>P if and only if Q</u> is called the biconditional statement

Notation: P⇔ Q

Meaning: P  $\Leftrightarrow$  Q means P  $\Rightarrow$  Q and Q  $\Rightarrow$  P.

26-Aug-2011

MA 341 001

### Truth Values

Р	Q	$P \Leftrightarrow Q$
True	True	True
True	False	False
False	True	False
False	False	True

26-Aug-2011

MA 341 001

26

## Compound Statements

If x is perpendicular to y and y is perpendicular to z, then x is parallel to z.

Let

P: x is perpendicular to y

 $\mathbf{Q}$ : y is perpendicular to z

 $\mathbf{R}$ : x is parallel to z.

 $P \wedge Q \Rightarrow R$ 

26-Aug-2011

	Р	Q	R	$P \land Q \Rightarrow R$	
	True	True	True	True	
	True	True	False	False	
	True	False	True	True	
	True	False	False	True	
	False	True	True	True	
	False	True	False	True	
	False	False	True	True	
	False	False	False	True	
26-Aug-2011	ug-2011 MA 341 001				

Universal Quantifier

If P is a statement that depends on a variable x, then the universal quantifier is For every x, P(x)

for every = for each = for all

$$\forall x, P(x)$$

22%-Alugg-22000

MA 341 001

# Existential Quantifier

If P is a statement that depends on a variable x, then the existential quantifier is

There exists an x such that P(x)

There is at least one = for at least one = some

 $\exists x, P(x)$ 

26-Aug-2011

•	-	•
7	•	١

## Translate the following

 $\forall x \forall y, x+y = 0$  For every x and for every y,

x+y = 0.

 $\forall x \; \exists y, \, x \!+\! y \!=\! 0 \quad \text{For every } x \; \text{there exists a } y$ 

so that x+y=0.

There exists an x so that for ∃x ∀y, x+y=0

every y, x+y = 0.

∃x ∃y, x+y=0 There exists an x and there

exists a y so that x+y = 0.

26-Aug-2011 MA 341 001

### Translate the following

For every x, if x is even, then there exists a ysuch that x = 2y.

 $\forall x (x \text{ is even} \Rightarrow \exists y, x = 2y)$ 

26-Aug-2011 MA 341 001

28

## Rules of Reasoning

A tautology is a sentence which is true no matter what the truth value of its constituent parts.

Example:

 $P \Rightarrow (P \vee Q)$ 

26-Aug-2011 MA 341 001

## Logic Axiom 1

Every tautology is a rule of reasoning.

- ♦  $(P\Rightarrow Q) \Leftrightarrow (\sim Q\Rightarrow \sim P)$  (The Contrapositive)
- **♦**  $[P \land (P \Rightarrow Q)] \Rightarrow Q$  (Modus ponens)
- ♦  $[(P\Rightarrow Q) \land (Q\Rightarrow R)]\Rightarrow (P\Rightarrow R)$  (Law of Syllogism)
- ~(P⇒Q) ⇔ (P∧~Q) (basis for Proof by Contradiction)
- $\bullet$  [(P $\Rightarrow$ R)  $\land$  (Q $\Rightarrow$ R)] $\Rightarrow$ [(P $\lor$ Q) $\Rightarrow$ R] Proof by Cases

26-Aug-2011

MA 341 001

34

### Logic Axiom 2

Let U denote a universal set. Each of the following is a rule of reasoning.

- $[\forall x, P(x) \Rightarrow Q(x)] \Rightarrow [\forall x, P(x) \Rightarrow \forall x, Q(x)]$
- $\forall x, P(x) \Leftrightarrow P(a)$  for any  $a \in U$
- $\exists x, P(x) \Leftrightarrow [P(a) \text{ for some } a \in U]$

26-Aug-2011

MA 341 001

35

## Logic Axiom 3

(Rule of Substitution) Suppose P  $\Leftrightarrow$  Q. Then P and Q may be substituted for one another in any sentence.

## Logic Axiom 4

Every sentence of the type  $\sim [\forall x, P(x)] \Leftrightarrow [\exists x \text{ so that } \sim P(x)]$  is true.

### Logic Axiom 5

Every sentence of the type  $\sim$ [ $\exists x$  so that P(x)]  $\Leftrightarrow$  [ $\forall x, \sim P(x)$ ] is true.

26-Aug-2011

MA 341 001

32