

## Logic, Proof, Axiom Systems

MA 341 - Topics in Geometry  
Lecture 03

---

---

---

---

---

---

---

---

$$\int_0^1 \frac{x^4(1-x)^4}{1+x^2} dx = \frac{22}{7} - \pi$$

$$111,111,111 \times 111,111,111 = 12,345,678,987,654,321$$

29-Aug-2011

MA 341 001

2

---

---

---

---

---

---

---

---

## Rules of Reasoning

A tautology is a sentence which is true no matter what the truth value of its constituent parts.

Example:

$$P \Rightarrow (P \vee Q)$$

29-Aug-2011

MA 341 001

3

---

---

---

---

---

---

---

---

### Logic Axiom 1

Every tautology is a rule of reasoning.

- ◆  $(P \Rightarrow Q) \Leftrightarrow (\sim Q \Rightarrow \sim P)$   
(The Contrapositive)
- ◆  $[P \wedge (P \Rightarrow Q)] \Rightarrow Q$   
(Modus ponens)
- ◆  $[(P \Rightarrow Q) \wedge (Q \Rightarrow R)] \Rightarrow (P \Rightarrow R)$   
(Law of Syllogism)
- ◆  $\sim(P \Rightarrow Q) \Leftrightarrow (P \wedge \sim Q)$  (basis for Proof by Contradiction)
- ◆  $[(P \Rightarrow R) \wedge (Q \Rightarrow R)] \Rightarrow [(P \vee Q) \Rightarrow R]$   
Proof by Cases

29-Aug-2011

MA 341 001

4

---

---

---

---

---

---

---

---

### Logic Axiom 2

Let  $U$  denote a universal set. Each of the following is a rule of reasoning.

- $[\forall x, P(x) \Rightarrow Q(x)] \Rightarrow [\forall x, P(x) \Rightarrow \forall x, Q(x)]$
- $\forall x, P(x) \Leftrightarrow P(a)$  for any  $a \in U$
- $\exists x, P(x) \Leftrightarrow [P(a)$  for some  $a \in U]$

29-Aug-2011

MA 341 001

5

---

---

---

---

---

---

---

---

### Logic Axiom 3

(Rule of Substitution) Suppose  $P \Leftrightarrow Q$ . Then  $P$  and  $Q$  may be substituted for one another in any sentence.

### Logic Axiom 4

Every sentence of the type  $\sim[\forall x, P(x)] \Leftrightarrow [\exists x$  so that  $\sim P(x)]$  is true.

### Logic Axiom 5

Every sentence of the type  $\sim[\exists x$  so that  $P(x)] \Leftrightarrow [\forall x, \sim P(x)]$  is true.

29-Aug-2011

MA 341 001

6

---

---

---

---

---

---

---

---

## Mathematical System

Consists of

1. Set of undefined concepts,
2. Universal set
3. Set of relations
4. Set of operations
5. Set of logical axioms
6. Set of axioms
7. Set of theorems
8. Set of definitions
9. An underlying set theory

29-Aug-2011

MA 341 001

7

---

---

---

---

---

---

---

---

---

---

## Proof

Suppose  $A_1, A_2, \dots, A_k$  are all the axioms and previously proved theorems of a mathematical system. A formal proof, or deduction, of a sentence  $P$  is a sequence of statements  $S_1, S_2, \dots, S_n$ , where

1.  $S_n$  is  $P$  and
2. One of the following holds
  - a)  $S_i$  is one of  $A_1, A_2, \dots, A_k$
  - b)  $S_i$  follows from the previous statements by valid argument

29-Aug-2011

MA 341 001

8

---

---

---

---

---

---

---

---

---

---

## Proving Conditionals

Suppose  $A_1, A_2, \dots, A_k$  are all the axioms and previously proved theorems of a mathematical system. A formal proof, or deduction, of a sentence  $P$  is a sequence of statements  $S_1, S_2, \dots, S_n$ , where

1.  $S_n$  is  $P$  and
2. One of the following holds
  - a)  $S_i$  is one of  $A_1, A_2, \dots, A_k$
  - b)  $S_i$  follows from the previous statements by valid argument

29-Aug-2011

MA 341 001

9

---

---

---

---

---

---

---

---

---

---

### Proof of $P \Rightarrow Q$

1. Assume  $P$  to be true.
2. Use  $P$  and all other theorems and axioms to deduce  $Q$ .
3. Once  $Q$  is deduced in this manner you have completed a proof of  $P \Rightarrow Q$ .

You have not shown that  $Q$  is true.

You have shown only that  $Q$  is true IF  $P$  is true

29-Aug-2011

MA 341 001

10

---

---

---

---

---

---

---

---

### Rule of Conditional Proof or Deduction Theorem

Suppose that  $A_1, A_2, \dots, A_k$  are the axioms and previously proved theorems. To prove  $P \Rightarrow Q$  is to show that

From  $A_1, A_2, \dots, A_k$  deduce  $P \Rightarrow Q$  is valid.  
To do this temporarily assume  $P$  to be an axiom and show that

From  $A_1, A_2, \dots, A_k, P$  deduce  $Q$  is valid.

29-Aug-2011

MA 341 001

11

---

---

---

---

---

---

---

---

### Proof by Contrapositive

We can prove  $P \Rightarrow Q$  by proving  $\sim Q \Rightarrow \sim P$ .

Rule of conditional proof used to prove the contrapositive

29-Aug-2011

MA 341 001

12

---

---

---

---

---

---

---

---

## Proving Biconditionals

Three techniques:

1. Prove  $P \Rightarrow Q$  and  $Q \Rightarrow P$ .
2. Prove  $P \Rightarrow Q$  and  $\sim P \Rightarrow \sim Q$ .
3. Prove an IFF string

29-Aug-2011

MA 341 001

13

---

---

---

---

---

---

---

---

## Proving $\forall x P(x)$

1. Let  $x$  represent arbitrary element of universal set
2. Prove that  $P(x)$  is true
3. Since  $x$  was arbitrary we know that  $\forall x P(x)$  is true.

The justification is Logical Axiom 2.

29-Aug-2011

MA 341 001

14

---

---

---

---

---

---

---

---

## Proof by Cases

Proving a statement of the form:  
 $(P \vee Q) \Rightarrow R$

Use the tautology:

$$[P \Rightarrow R] \wedge [Q \Rightarrow R] \Leftrightarrow [(P \vee Q) \Rightarrow R]$$

29-Aug-2011

MA 341 001

15

---

---

---

---

---

---

---

---

## Mathematical Induction

Let  $P(n)$  be a statement for any integer  $n$ .

Principle of Induction:

$$[P(1) \wedge (\forall k, P(k) \Rightarrow P(k+1))] \Rightarrow \forall n, P(n)$$

2 step process:

Basic Step: Prove  $P(1)$ .

Inductive step: Prove  $\forall k, P(k) \Rightarrow P(k+1)$

29-Aug-2011

MA 341 001

16

---

---

---

---

---

---

---

---

---

---

## Proof by Contradiction

Contradiction: statement which is false no matter truth value of its constituent parts

$$R \wedge \sim R$$

Tautology:  $[\sim P \Rightarrow (R \wedge \sim R)] \Rightarrow P$

Usually used to prove statement of type  $P \Rightarrow Q$ .

Deduction Theorem: Assume  $P$  and deduce  $P \Rightarrow Q$

Contrapositive: Assume  $\sim Q$  and deduce  $\sim Q \Rightarrow \sim P$

Contradiction: Assume  $P$  AND  $\sim Q$  and deduce

$$R \wedge \sim R$$

29-Aug-2011

MA 341 001

17

---

---

---

---

---

---

---

---

---

---

## Proving $\exists x P(x)$

Prove that there is at least one  $x$  so that  $P(x)$  is true. Here is one case where an example is good enough.

The other case is when you find a "counterexample" to show  $\sim(\forall x P(x))$ .

29-Aug-2011

MA 341 001

18

---

---

---

---

---

---

---

---

---

---

### How to Create a Proof

Suggestions - NOT RULES

1. Translate to symbolic logic
2. Analogy - look at other proofs
3. Reverse engineering - you know where you want to end, figure out each step it would require to get there (analogy - trig identities)
4. Use definitions
5. Use **previously** proved theorems

29-Aug-2011 MA 341 001 19

---

---

---

---

---

---

---

---

### Euclid's Five Axioms

Let the following be postulated

1. To draw a straight line from any point to any point.
2. To produce a finite straight line continuously in a straight line.
3. To describe a circle with any center and distance.
4. That all right angles are equal to one another.
5. That, if a straight line falling on two straight lines make the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which are the angles less than two right angles. (Euclid's Parallel Postulate)

29-Aug-2011 MA 341 001 20

---

---

---

---

---

---

---

---

### Hilbert's Axioms

**Incidence Axioms**

I-1: For every point P and for every point Q not equal to P there exists a unique line that passes through P and Q.  
 I-2: For every line there exist at least two distinct points incident with it.  
 I-3: There exist three distinct points with the property that no line is incident with all three of them.

29-Aug-2011 MA 341 001 21

---

---

---

---

---

---

---

---

### Hilbert's Axioms

**Betweenness Axioms**

B-1: If  $A*B*C$ , then  $A$ ,  $B$ , and  $C$  are 3 distinct points all lying on the same line and  $C*B*A$ .

B-2: Given any two distinct points  $B$  and  $D$ , there exists points  $A$ ,  $C$ , and  $E$  lying on  $BD$  such that  $A*B*D$ ,  $B*C*D$ , and  $B*D*E$ .

B-3: If  $A$ ,  $B$ , and  $C$  are three distinct point lying on the same line, then one and only one of the points is between the other two.

29-Aug-2011 MA 341 001 22

---

---

---

---

---

---

---

---

### Hilbert's Axioms

**Betweenness Axioms**

B-4: (Plane Separation Axiom) For every line  $l$  and for any 3 points  $A$ ,  $B$ , and  $C$  not lying on  $l$ :

- i. If  $A$  and  $B$  are on the same side of  $l$  and  $B$  and  $C$  are on the same side of  $l$ , then  $A$  and  $C$  are on the same side of  $l$ .
- ii. If  $A$  and  $B$  are on opposite sides of  $l$  and  $B$  and  $C$  are on opposite sides of  $l$ , then  $A$  and  $C$  are on the same side of  $l$ .

29-Aug-2011 MA 341 001 23

---

---

---

---

---

---

---

---

### Hilbert's Axioms

**Congruence Axioms**

C-1: If  $A$  and  $B$  are distinct points and if  $A'$  is any point, then for each ray  $r$  emanating from  $A'$  there is a unique point  $B'$  on  $r$  such that  $B' \neq A'$  and  $AB \cong A'B'$ .

C-2: If  $AB \cong CD$  and  $AB \cong EF$ , then  $CD \cong EF$ . Also, every segment is congruent to itself.

C-3: If  $A*B*C$  and  $A'*B'*C'$  and  $AB \cong A'B'$  and  $BC \cong B'C'$ , then  $AC \cong A'C'$ .

C-4: Given any  $\angle BAC$  and given a ray  $A'B'$  emanating from point  $A'$ ,  $\exists!$  ray  $A'C'$  on a given side of line  $A'B'$  so that  $\angle BAC \cong \angle B'A'C'$ .

29-Aug-2011 MA 341 001 24

---

---

---

---

---

---

---

---



### Hilbert's Axioms

**Congruence Axioms**

C-5: If  $\angle A \cong \angle B$  and  $\angle B \cong \angle C$  then  $\angle A \cong \angle C$ . Also, every angle is congruent to itself.

C-6: (Side-Angle-Side) If two sides and the included angle of one triangle are congruent respectively to two sides and the included angle of another triangle, then the two triangles are congruent.

29-Aug-2011 MA 341 001 25

---

---

---

---

---

---

---

---

### Hilbert's Axioms

**Continuity Axioms**

Archimedes Axiom: If  $AB$  and  $CD$  are any segments, then there is a number  $n$  such that if segment  $CD$  is laid off  $n$  times on the ray  $AB$  emanating from  $A$ , then a point  $E$  is reached where  $n CD \cong AE$  and  $B$  is between  $A$  and  $E$ .

Dedekind's Axiom: Suppose that the set of all points on a line  $l$  is the union  $\Sigma_1 \cup \Sigma_2$  of two nonempty subsets such that no point of  $\Sigma_1$  is between two points of  $\Sigma_2$  and vice versa. Then there is a unique point,  $O$ , lying on  $l$  such that  $P_1 * O * P_2$  if and only if  $P_1 \in \Sigma_1$  and  $P_2 \in \Sigma_2$  and  $O \neq P_1, P_2$ .

29-Aug-2011 MA 341 001 26

---

---

---

---

---

---

---

---

### Hilbert's Axioms

**Continuity Axioms**

Elementary Continuity Principle: If one endpoint of a segment is inside a circle and the other outside, then the segment intersects the circle.

Circular Continuity Principle: If a circle has one point inside and one point outside another circle, then the two circles intersect in two points.

29-Aug-2011 MA 341 001 27

---

---

---

---

---

---

---

---

### Birkhoff's Axioms

B1. There exist nonempty subsets of the plane called lines, with the property that each two points belong to exactly one line.

B2. Corresponding to any two points  $A$  and  $B$  in the plane there exists a unique real number  $d(AB) = d(BA)$ , the distance from  $A$  to  $B$ , which is 0 if and only if  $A = B$ .

29-Aug-2011

MA 341 001

28

---

---

---

---

---

---

---

---

---

---

### Birkhoff's Axioms

B3. (Birkhoff Ruler Axiom) If  $k$  is a line and  $\mathbf{R}$  denotes the set of real numbers, there exists a one-to-one correspondence  $(X \rightarrow x)$  between the points  $X$  in  $k$  and the numbers  $x$  in  $\mathbf{R}$  such that  $d(A, B) = |a - b|$  where  $A \rightarrow a$  and  $B \rightarrow b$ .

B4. For each line  $k$  there are exactly two nonempty convex sets  $R'$  and  $R''$  satisfying

- a)  $R' \cup k \cup R''$  is the entire plane,
- b)  $R' \cap R'' = \emptyset$ ,  $R' \cap k = \emptyset$ , and  $R'' \cap k = \emptyset$
- c) If  $X$  in  $R'$  and  $Y$  in  $R''$  then  $XY \cap k \neq \emptyset$

29-Aug-2011

MA 341 001

29

---

---

---

---

---

---

---

---

---

---

### Birkhoff's Axioms

B5. For each angle  $\angle ABC$  there exists a unique real number  $x$  with  $0 \leq x \leq 180$  which is the (degree) measure of the angle  $x = \angle ABC^\circ$ .

B6. If ray  $\overrightarrow{BD}$  lies in  $\angle ABC$ , then  $\angle ABD^\circ + \angle DBC^\circ = \angle ABC^\circ$ .

29-Aug-2011

MA 341 001

30

---

---

---

---

---

---

---

---

---

---

### Birkhoff's Axioms

B7. If  $\vec{AB}$  is a ray in the edge,  $k$ , of an open half plane  $H(k;P)$  then there exist a one-to-one correspondence between the open rays in  $H(k;P)$  emanating from  $A$  and the set of real numbers between 0 and 180 so that if  $\vec{AX} \rightarrow x$  then  $\angle BAX^\circ = x$ .

29-Aug-2011

MA 341 001

31

---

---

---

---

---

---

---

---

---

---

### Birkhoff's Axioms

B8. (SAS) If a correspondence of two triangles, or a triangle with itself, is such that two sides and the angle between them are respectively congruent to the corresponding two sides and the angle between them, the correspondence is a congruence of triangles.

29-Aug-2011

MA 341 001

32

---

---

---

---

---

---

---

---

---

---

### SMSG Axioms

3 undefined terms: point, line, plane

22 axioms - a mixture of Birkhoff's and Hilbert's meant to make it easier to teach geometry in high school and college.

29-Aug-2011

MA 341 001

33

---

---

---

---

---

---

---

---

---

---