

## 1. CONTINUITY AND LIMITS: DEFINITIONS

In the notes on the continuum we defined a continuous function as follows.

**Definition 1.** A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if for every open set  $U \subseteq \mathbb{R}$ ,  $f^{-1}(U)$  is open.

Theorem 5.5 from those notes provides an equivalent definition:

**Definition 2.** A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if for every  $p \in \mathbb{R}$  and every open interval  $I_1$  which contains  $f(p)$ , there exists an open interval  $I_2$  which contains  $p$  such that  $f(I_2) \subseteq I_1$ .

Replacing open intervals by open balls and rephrasing in terms of inequalities gives the following definition:

**Definition 3.** A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous if for every  $p \in \mathbb{R}$  and every  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$$\text{if } |x - p| < \delta \text{ then } |f(x) - f(p)| < \varepsilon.$$

**Theorem 1.1.** The above three definitions are all equivalent.

**Definition 4.** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$ , and  $p \in \mathbb{R}$ . We say that  $\lim_{x \rightarrow p} f(x) = L$  if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that

$$\text{if } 0 < |x - p| < \delta \text{ then } |f(x) - L| < \varepsilon.$$

The limit is well defined:

**Theorem 1.2.** If the limit exists then it is unique: i.e. if

$$\lim_{x \rightarrow p} f(x) = L \text{ and } \lim_{x \rightarrow p} f(x) = M$$

then  $L = M$ .

Now we have a fourth definition of continuity!

**Definition 5.** A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous at  $p$  if

$$\lim_{x \rightarrow p} f(x) = f(p).$$

We say  $f$  is continuous if this holds for every  $p \in \mathbb{R}$ .

Examination of the definition of limit shows that only the values of  $f(x)$  for  $x$  near (and not equal to)  $p$  determine the limit at  $p$ . This allows us to define the limit for functions whose domain is not all of  $\mathbb{R}$ .

**Definition 6.** Suppose  $A \subset \mathbb{R}$  is open, and  $p \in A$ , and suppose  $f : B \rightarrow \mathbb{R}$  where  $A \setminus \{p\} \subseteq B$ .

We say that  $\lim_{x \rightarrow p} f(x) = L$  if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that

$$\text{if } 0 < |x - p| < \delta \text{ and } x \in B \text{ then } |f(x) - L| < \varepsilon.$$

**Remark:** In the spirit of the  $\varepsilon - \delta$  definitions of  $\lim_{x \rightarrow a} f(x) = L$ , we can also give definitions for

$$\lim_{x \rightarrow \infty} f(x) = L, \quad \lim_{x \rightarrow a} f(x) = \infty,$$

and several other variations.

## 2. LIMIT THEOREMS

**Theorem 2.1.** *Suppose*

$$\lim_{x \rightarrow p} f(x) = L \quad \text{and} \quad \lim_{x \rightarrow p} g(x) = M.$$

*Then*

$$\begin{aligned} \lim_{x \rightarrow p} (f(x) + g(x)) &= L + M, \\ \lim_{x \rightarrow p} f(x)g(x) &= LM, \\ \text{and } \lim_{x \rightarrow p} \frac{1}{g(x)} &= \frac{1}{M} \quad \text{if } M \neq 0. \end{aligned}$$

**Remark:** This shows that the sum, product, and quotient of continuous functions is continuous (as long as the denominator is nonzero, in the case of the quotient).

**Theorem 2.2** (Squeeze Theorem \*). *Suppose  $f(x) \leq g(x) \leq h(x)$  for all  $x \in \mathbb{R}$ , and*

$$\lim_{x \rightarrow p} f(x) = \lim_{x \rightarrow p} h(x) = L.$$

*Then  $\lim_{x \rightarrow p} g(x) = L$ .*

**Theorem 2.3** (\*). *Suppose  $\lim_{x \rightarrow p} g(x) = L$ , and  $f$  is continuous at  $L$ . Then*

$$\lim_{x \rightarrow p} f(g(x)) = f(L).$$

**Proposition 2.4** (\*). *Suppose  $f(x) \geq 0$ . Then  $\lim_{x \rightarrow a} f(x) \geq 0$  if the limit exists.*

## 3. DERIVATIVES

**Definition 7.** *Suppose  $A \subset \mathbb{R}$  and  $f : A \rightarrow \mathbb{R}$ . We say  $f$  is differentiable at  $a$  if the limit*

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

*exists. Then we say that this limit is the derivative of  $f$  at  $a$ , and denote this by  $f'(a)$ .*

**Theorem 3.1.** *If  $f$  is differentiable at  $a$ , then  $f$  is continuous at  $a$ .*

Note that the converse is false: find a counterexample!

A straightforward application of the definition shows that the identity function  $f(x) = x$  and the constant function  $g(x) = c$  are differentiable everywhere, and their derivatives are 1 and 0, respectively, for all  $x$ .

Now the following rules for sums, products, and quotients let us differentiate any rational function.

**Theorem 3.2** (Sum, Product, and Quotient Rules). *If  $f$  and  $g$  are differentiable at  $x$ , then*

- $f + g$  is differentiable at  $x$  and  $(f + g)'(x) = f'(x) + g'(x)$ .
- $f \cdot g$  is differentiable at  $x$ , and  $(fg)'(x) = f'(x)g(x) + f(x)g'(x)$ .
- If  $g'(x) \neq 0$  then  $1/g$  is differentiable at  $x$ , and  $(1/g)'(x) = -g'(x)/g^2(x)$ .

Finally, we prove the chain rule. First a lemma.

**Lemma 3.3.** *Suppose  $g$  is differentiable at  $a$  and  $f$  is differentiable at  $g(a)$ . Define*

$$\phi(h) = \begin{cases} \frac{f(g(a+h)) - f(g(a))}{f'(g(a))} & \text{if } g(a+h) - g(a) \neq 0 \\ f'(g(a)) & \text{otherwise} \end{cases}$$

*Show that  $\phi$  is continuous at 0.*

**Theorem 3.4** (Chain Rule). *Suppose  $g$  is differentiable at  $a$  and  $f$  is differentiable at  $g(a)$ . Then  $f \circ g$  is differentiable at  $a$ , and*

$$(f \circ g)'(a) = f'(g(a))g'(a).$$

**Definition 8.** *Let  $A$  be a subset of the domain of  $f$ . We say that  $x \in A$  is a maximum point for  $f$  on  $A$  if  $f(x) \geq f(y)$  for all  $y \in A$ . A similar definition holds for minimum point.*

**Theorem 3.5** (\*). *Suppose  $f$  is defined on  $(a, b)$ , and  $x$  is a maximum or minimum point of  $f$  on  $(a, b)$ . If  $f$  is differentiable at  $x$  then  $f'(x) = 0$ .*

**Lemma 3.6** (Rolle's Theorem). *Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , and  $f(a) = f(b)$ . Then there exists  $x \in (a, b)$  such that  $f'(x) = 0$ .*

**Theorem 3.7** (Mean Value Theorem \*). *Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . Then there exists  $x \in (a, b)$  such that*

$$f'(x) = \frac{f(b) - f(a)}{b - a}.$$

**Corollary 3.8.** *If  $f'(x) = 0$  for all  $x \in (a, b)$ , then  $f$  is constant on  $(a, b)$ .*

**Corollary 3.9** (Cauchy Mean Value Theorem). *Suppose  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ . Then there exists  $x \in (a, b)$  such that*

$$f'(x) = \frac{f(b) - f(a)}{b - a}.$$

**Theorem 3.10** (L'Hôpital's Rule). *Suppose  $\lim_{x \rightarrow p} f(x) = \lim_{x \rightarrow p} g(x) = 0$  and  $\lim_{x \rightarrow p} \frac{f'(x)}{g'(x)}$  exists.*

*Then*

$$\lim_{x \rightarrow p} \frac{f(x)}{g(x)} = \lim_{x \rightarrow p} \frac{f'(x)}{g'(x)}.$$

*In particular the limit on the left exists.*

## 4. UNIFORM CONTINUITY

**Definition 9.** We say  $f$  is uniformly continuous on  $A$  if for all  $\varepsilon > 0$  there exists  $\delta > 0$  such that for all  $x, p \in A$ ,

$$\text{if } |x - p| < \delta \text{ then } |f(x) - f(p)| < \varepsilon.$$

**Exercise 4.1.** Give an example of a function that is continuous on  $(0, 1)$  but not uniformly continuous there.

**Theorem 4.2** (\*). If  $A$  is compact and  $f$  is continuous on  $A$  then  $f$  is uniformly continuous on  $A$ .

## 5. INTEGRALS

**Definition 10.** A partition  $P$  of  $[a, b]$  is a finite collection of points  $t_0, \dots, t_n$  such that

$$a = t_0 < \dots < t_n = b.$$

**Definition 11.** Suppose  $f$  is bounded on  $[a, b]$ , and  $P$  is a partition of  $[a, b]$ . The upper and lower sums of  $f$  on  $[a, b]$  with respect to  $P$  are the quantities

$$U(f, P) = \sum_{i=1}^n M_i(t_i - t_{i-1}) \quad \text{and} \quad L(f, P) = \sum_{i=1}^n m_i(t_i - t_{i-1})$$

where  $M_i$  and  $m_i$  are the supremum and infimum, respectively, of the sets

$$\{f(x) | x \in [t_{i-1}, t_i]\}.$$

**Lemma 5.1.** Let  $P$  and  $Q$  be partitions of  $[a, b]$  such that all the points of  $P$  are also in  $Q$ . Then

$$L(f, P) \leq L(f, Q) \quad \text{and} \quad U(f, P) \geq U(f, Q).$$

**Theorem 5.2.** Let  $P_1, P_2$  be partitions of  $[a, b]$ . Then

$$L(f, P_1) \leq U(f, P_2).$$

Note that this implies that  $\inf_P U(f, P) \geq \sup_P L(f, P)$ .

**Definition 12.** Suppose  $f$  is bounded on  $[a, b]$ . We say  $f$  is (Riemann) integrable on  $[a, b]$  if

$$\inf_P U(f, P) = \sup_P L(f, P),$$

and define this quantity to be the integral

$$\int_a^b f(x) dx.$$

**Lemma 5.3** (\*). Suppose  $f$  is bounded on  $[a, b]$ . Then  $f$  is Riemann integrable on  $[a, b]$  if and only if for all  $\varepsilon > 0$ , there exists a partition  $P$  of  $[a, b]$  such that

$$U(f, P) - L(f, P) < \varepsilon.$$

**Theorem 5.4.** Suppose  $f$  is continuous on  $[a, b]$ . Then  $f$  is integrable on  $[a, b]$ .

**Theorem 5.5.** *Suppose  $f$  is integrable on  $[a, b]$  and on  $[b, c]$ . Then  $f$  is integrable on  $[a, c]$ , and*

$$\int_a^c f(x)dx = \int_a^b f(x)dx + \int_b^c f(x)dx.$$

**Remark:** This formula justifies the definition

$$\int_b^a f(x)dx = - \int_a^b f(x)dx.$$

**Theorem 5.6.** *Suppose  $f$  is integrable on  $[a, b]$ , and  $c$  is constant. Then  $cf$  is integrable on  $[a, b]$ , and*

$$\int_a^b cf(x)dx = c \int_a^b f(x)dx$$

**Theorem 5.7.** *Suppose  $f, g$  are integrable on  $[a, b]$ . Then  $f + g$  is integrable on  $[a, b]$ , and*

$$\int_a^b (f + g)(x)dx = \int_a^b f(x)dx + \int_a^b g(x)dx.$$

**Lemma 5.8.** *Suppose  $m \leq f(x) \leq M$  for all  $x \in [a, b]$ . Then*

$$m(b - a) \leq \int_a^b f(x)dx \leq M(b - a).$$

**Proposition 5.9.** *Suppose  $f$  and  $g$  are integrable on  $[a, b]$  and  $f \leq g$  on  $[a, b]$ . Then*

$$\int_a^b f(x)dx \leq \int_a^b g(x)dx.$$

**Proposition 5.10 (\*)**. *Suppose  $f$  is integrable on  $[a, b]$  and*

$$F(x) = \int_a^x f(t)dt.$$

*Then  $F$  is continuous on  $[a, b]$ .*

**Theorem 5.11** (1st Fundamental Theorem\*). *Suppose  $f$  is integrable on  $[a, b]$  and define*

$$F(x) = \int_a^x f(t)dt.$$

*If  $f$  is continuous at  $c$  then  $F$  is differentiable at  $c$  and*

$$F'(c) = f(c).$$

**Theorem 5.12** (2nd Fundamental Theorem). *Suppose  $f$  is integrable on  $[a, b]$  and  $f = g'$  for some  $g$ . Then*

$$\int_a^b f(x)dx = g(b) - g(a).$$