

## 5 Angles

### 5.1 The Angle Axioms and Basic Theorems

The following is a summary of Section 2.7 of Kay.

**Axiom A-1 — Existence of Angle Measure:** To every angle  $\angle A$  there corresponds a unique, real number  $\theta = m\angle A$ ,  $0 < \theta < 180$ , called its *measure*.

**Axiom A-2 — Angle Addition Postulate:** If  $D$  lies in the interior of  $\angle ABC$ , then  $m\angle ABC = m\angle ABD + m\angle DBC$ , and conversely.

**Axiom A-3 — Protractor Postulate:** The set of rays having a common origin  $O$  and lying on one side of line  $\ell = \overleftrightarrow{OA}$ , including ray  $\overrightarrow{OA}$ , may be assigned to the real numbers  $\theta$  for which  $0 \leq \theta < 180$ , called *coordinates*, in such a manner that

1. Each ray is assigned a unique coordinate  $\theta$ .
2. Each coordinate  $\theta$  is assigned to a unique ray.
3. The coordinate of  $\overrightarrow{OA}$  is 0.
4. If rays  $\overrightarrow{OP}$  and  $\overrightarrow{OQ}$  have coordinates  $\theta$  and  $\phi$ , respectively, then  $m\angle POQ = |\theta - \phi|$ .

**Definition:** Suppose that  $\overrightarrow{OA}$ ,  $\overrightarrow{OB}$ , and  $\overrightarrow{OC}$  are concurrent rays, all having the same endpoint  $O$ . Then if these rays are distinct (no two are the same ray), and if  $m\angle AOB + m\angle BOC = m\angle AOC$ , then  $\overrightarrow{OB}$  is said to lie *between*  $\overrightarrow{OA}$  and  $\overrightarrow{OC}$ , and we write  $\overrightarrow{OA}-\overrightarrow{OB}-\overrightarrow{OC}$ .

**Notation:** Write  $A[a]$  if  $a$  is the coordinate of a point  $A$  under the Ruler Postulate. Write  $\overrightarrow{OA}[a]$  if  $a$  is the coordinate of a ray  $\overrightarrow{OA}$  under the Protractor Postulate.

**Theorem 2.7.1:** If  $A[a]$ ,  $B[b]$ , and  $C[c]$  are three collinear points (and  $\overrightarrow{OA}[a]$ ,  $\overrightarrow{OB}[b]$ ,  $\overrightarrow{OC}[c]$  three concurrent rays) with their coordinates, then  $A-B-C$  ( $\overrightarrow{OA}-\overrightarrow{OB}-\overrightarrow{OC}$ ) if and only if  $a < b < c$  or  $c < b < a$ . (This is Theorem 1 of Kay, Section 2.7.)

**Corollary:** Suppose that four distinct collinear points are given with their coordinates:  $A[a]$ ,  $B[b]$ ,  $C[c]$ ,  $D[d]$ . If  $A-B-C$  and  $A-C-D$ , then  $A-B-C-D$ , and similarly for rays  $\overrightarrow{OA}[a]$ ,  $\overrightarrow{OB}[b]$ ,  $\overrightarrow{OC}[c]$ ,  $\overrightarrow{OD}[d]$ .

**Lemma:** A segment, ray, or line is a convex set.

**Lemma:** If  $A$  and  $B$  are two distinct points, and  $C \in \overrightarrow{AB}$ , with  $A \neq C$ , then  $\overrightarrow{AB} \subseteq \overrightarrow{AC}$ .

**Theorem 2.7.2:** If  $C \in \overrightarrow{AB}$  and  $A \neq C$ , then  $\overrightarrow{AB} = \overrightarrow{AC}$ . (This is Theorem 2 of Kay, Section 2.7.)

**Theorem 2.7.3 (Segment Construction Theorem):** If  $\overline{AB}$  and  $\overline{XY}$  are any two segments and  $AB \neq XY$ , then there is a unique point  $C$  on ray  $\overrightarrow{AB}$  such that  $AC = XY$ , with  $A-C-B$  if  $XY < AB$ , or  $A-B-C$  if  $XY > AB$ . (This is Theorem 3 of Kay, Section 2.7.)

**Definition:** A point  $M$  on a segment  $\overline{AB}$  is called a *midpoint* if it has the property that  $AM = MB$ . Such a midpoint is also said to *bisect* the segment, and any line, segment, or ray passing through that midpoint is also said to *bisect* the segment.

**Theorem 2.7.4 (Midpoint Construction Theorem):** The midpoint of any segment exists and is unique. (This is Theorem 4 of Kay, Section 2.7.)

**Theorem 2.7.5 (Segment Doubling Theorem):** There exists a unique point  $C$  on ray  $\overrightarrow{AB}$  such that  $B$  is the midpoint of  $\overline{AC}$ . (This is Theorem 5 of Kay, Section 2.7.)

**Definition:** A ray  $\overrightarrow{OM}$  such that  $\overrightarrow{OA}-\overrightarrow{OM}-\overrightarrow{OB}$  is said to be an *angle bisector* of  $\angle AOB$  if  $m\angle AOM = m\angle OMB$ . Any line or ray containing an angle bisector is said to *bisect* the angle.

**Theorem 2.7.3' (Angle Construction Theorem):** If  $\angle ABC$  and  $\angle XYZ$  are any two nondegenerate angles and  $m\angle ABC \neq m\angle XYZ$ , then there exists a unique ray  $\overrightarrow{BD}$  on the  $C$ -side of  $\overleftrightarrow{AB}$  such that  $m\angle XYZ = m\angle ABD$ , and either  $\overrightarrow{BA}-\overrightarrow{BD}-\overrightarrow{BC}$  if  $m\angle XYZ < m\angle ABC$ , or  $\overrightarrow{BA}-\overrightarrow{BC}-\overrightarrow{BD}$  if  $m\angle XYZ > m\angle ABC$ . (This is Theorem 3' of Kay, Section 2.7.)

**Theorem 2.7.4' (Angle Bisection Theorem):** Every angle has a unique bisector. (This is Theorem 4' of Kay, Section 2.7.)

**Theorem 2.7.5' (Angle Doubling Theorem):** Given any angle  $\angle ABC$  having measure  $< 90$ , there exists a ray  $\overrightarrow{BD}$  such that  $\overrightarrow{BC}$  is the bisector of  $\angle ABD$ . (This is Theorem 5' of Kay, Section 2.7.)

## 5.2 More Theorems on Angles

This is a summary of Section 2.8 of Kay.

**Theorem 2.8.1 (Crossbar Theorem):** If  $D$  is in the interior of  $\angle BAC$ , then ray  $\overrightarrow{AD}$  meets segment  $\overline{BC}$  at some interior point  $E$ . (This is Theorem 1 of Kay, Section 2.8.)

**Definition:** If  $A-B-C$  then  $\overrightarrow{BA}$  and  $\overrightarrow{BC}$  are called *opposite rays*.

**Lemma:** For every ray  $\overrightarrow{PQ}$  there exists a unique opposite ray.

**Definition:** If the sides of one angle are opposite rays to the respective sides of another angle, the angles are said to form a *vertical pair*.

**Definition:** Two angles are said to form a *linear pair* iff they have one side in common and the other two sides are opposite rays. We call any two angles whose angle measures sum to 180 a *supplementary pair*, or more simply, *supplementary*, and two angles whose angle measures sum to 90 a *complementary pair*, or *complementary*.

**Theorem 2.8.2:** Two angles which are supplementary (or complementary) to the same angle have equal angle measures. (This is Theorem 2 of Kay, Section 2.8.)

**Axiom A-4:** A linear pair of angles is a supplementary pair.

**Theorem 2.8.3 (Vertical Pair Theorem):** Vertical angles have equal measures. (This is Theorem 3 of Kay, Section 2.8.)

**Definition:** If line  $\ell$  intersects another line  $m$  at some point  $A$  and forms a supplementary pair of angles at  $A$  having equal measures, then  $\ell$  is said to be *perpendicular* to  $m$ , and we write  $\ell \perp m$ .

**Definition:** An angle having measure 90 is called a *right angle*. Angles having measure less than 90 are *acute angles*, and those with measure greater than 90, *obtuse angles*.

**Theorem 2.8.4:** One line is perpendicular to another line iff the two lines form four right angles at their point of intersection. (This is Theorem 4 of Kay, Section 2.8.)

**Corollary:** Line  $\ell$  is perpendicular to line  $m$  iff  $\ell$  and  $m$  contains the sides of a right angle.

**Theorem 2.8.5 (Existence and Uniqueness of Perpendiculars):** Suppose that in some plane line  $m$  is given and an arbitrary point  $A$  on  $m$  is located. Then there exists a unique line  $\ell$  that is perpendicular to  $m$  at  $A$ . (This is Theorem 5 of Kay, Section 2.8.)