
MA123 — Elem. Calculus	Fall 2017	Name:	500.
Exam 3	2017-11-16		Sec.:

Do not remove this answer page — you will turn in the entire exam. No books or notes may be used. You may use an ACT-approved calculator during the exam, but NO calculator with a Computer Algebra System (CAS), networking, or camera is permitted. Absolutely no cell phone use during the exam is allowed.

The exam consists of two short answer questions and eighteen multiple choice questions. Answer the short answer questions on the back of this page, and record your answers to the multiple choice questions on this page. For each multiple choice question, you will need to fill in the circle corresponding to the correct answer. It is your responsibility to make it CLEAR which response has been chosen. For example, if (a) is correct, you must write



You have two hours to do this exam. Please write your name and section number on this page.




GOOD LUCK!

For grading use:

Multiple Choi	ce Short Ans	swer
(number right) (5 points	s each) (out of 10 po	ints)

Total		
	(out of 100 points)	

1. Sketch the graph of a **continuous** function y = f(x) which satisfies f(1) = 4, f'(x) > 0 on $(-\infty,3)$, f'(x) < 0 on $(3,\infty)$, and f(x) is concave up on both $(-\infty,3)$ and $(3,\infty)$.



A farmer builds a rectangular grid of pens with 1 row and 3 columns using 950 feet of fencing. Find the dimensions (overall length and width) that will maximize the total area of the pen. You must clearly use calculus to find <u>and justify</u> your answer.



Name:

Multiple Choice Questions

Show all your work on the page where the question appears. Clearly mark your answer both on the cover page on this exam and in the corresponding questions that follow.

3. Where is the function $f(t) = (t - 69)^{-2}$ decreasing?

Possibilities:

- (a) f(t) is always decreasing except at t = 69
- (b) f(t) is never decreasing
- (c) t < 69
- (d) -1 < t < 69
- (e) t > 69

4. Where is the function $f(t) = t^4 - 16t^3 - 3$ concave up?

- (a) t < 0 and t > 8
- (b) 0 < t < 8
- (c) t < 12
- (d) t > 12
- (e) f(t) is always concave up

5. Suppose the derivative of g(t) is $g'(t) = 15(t-5)(t-9)^2$. Find the interval(s) of values of t in which g is decreasing.

Possibilities:

- (a) $(9, \infty)$
- (b) (5,9)
- (c) $(5,9) \cup (15,\infty)$
- (d) $(-\infty, 5)$
- (e) $(-\infty, 5) \cup (9, \infty)$


6. Suppose the derivative of g(t) is g'(t) = 15(t-5)(t-9). For t in which interval(s) is g concave up?

- (a) $(-\infty,7)$
- (b) $(7, \infty)$
- (c) $(5,7) \cup (9,15)$
- (d) (5,9)
- (e) $(-\infty,5) \cup (9,\infty)$

7. The following is the graph of the derivative, f'(x), of the function f(x). Where is the regular function f(x) decreasing?


Possibilities:

- (a) $(-\infty, -1)$
- (b) $(-\infty, 4)$
- (c) $(4,\infty)$
- (d) (-2,1)
- (e) $(-\infty, -2)$ and $(1, \infty)$



8. The following is the graph of the derivative, f'(x), of the function f(x). Where is the regular function f(x) concave up?

- (a) $(-\infty, -1)$
- (b) $(-\infty, -2)$ and $(1, \infty)$
- (c) $(-\infty, 4)$
- (d) $(4,\infty)$
- (e) (-2, 1)




9. Find the critical numbers of the function $f(x) = 7xe^{19x}$.

Possibilities:

(a)
$$-\frac{7}{19}$$
, 0
(b) $-\frac{1}{19}$
(c) $-\frac{7}{19}$
(d) $-\frac{1}{19}$, 0, e^{19}
(e) 0

10. Consider the graph of the original function, f(x). For this function, what are the signs of f'(-3) and f''(-3)?

- (a) f'(-3) = 0 and f''(-3) < 0
- (b) f'(-3) > 0 and f''(-3) < 0
- (c) f'(-3) < 0 and f''(-3) < 0
- (d) f'(-3) < 0 and f''(-3) > 0
- (e) f'(-3) > 0 and f''(-3) > 0



11. The manager of a large apartment complex knows from experience that 110 units will be occupied if the rent is 460 dollars per month. A market survey suggests that, on the average, one additional unit will remain vacant for each 7 dollar increase in rent. Similarly, one additional unit will be occupied for each 7 dollar decrease in rent. What rent should the manager charge to maximize revenue?

Possibilities:

- (a) \$614.20 per month
- (b) \$614.60 per month
- (c) \$615.80 per month
- (d) \$615.60 per month
- (e) \$615.00 per month

12. A landscape architect wishes to enclose a rectangular garden on one side by a brick wall costing \$80 per foot, and on the other three sides by a metal fence costing \$40 per foot. If the area of the garden is 300 square feet, find the lowest possible cost to enclose the garden.


- (a) \$3393.61
- (b) \$3395.11
- (c) \$3394.11
- (d) \$3394.61
- (e) \$3395.61

13. Given the function $f(x) = \begin{cases} x & \text{if } x < 60 \\ 60 & \text{if } x \ge 60 \end{cases}$ evaluate the definite integral


$$\int_0^{70} f(x) \, \mathrm{d}x$$

Possibilities:

- (a) 2400
- (b) 2401
- (c) 2402
- (d) 2403
- (e) 2404



- (a) 2.5
- (b) 7.5
- (c) 21.5
- (d) 6
- (e) 1.5



15. Suppose that $\int_{4}^{13} f(x) dx = 20$, $\int_{28}^{45} f(x) dx = 48$, and $\int_{4}^{45} f(x) dx = 12$. Find the value of $\int_{13}^{28} f(x) dx$.

Possibilities:

- (a) 80
- (b) -56
- (c) -80
- (d) 16
- (e) **-**504

16. Suppose that $\int_{5}^{20} f(x) dx = 14$. Find the value of $\int_{5}^{20} (3f(x) + 60) dx$.

- (a) 102
- (b) 57
- (c) 942
- (d) 222
- (e) 1242

17. Find the **average value** of f(x) on the interval [7, 15] given that $f(x) = \begin{cases} 40 & \text{if } x < 10 \\ -80 & \text{if } x \ge 10. \end{cases}$

Possibilities:

- (a) -15
- (b) -140
- (c) -20
- (d) -35
- (e) 6

18. Estimate the area under the graph of $y = -x^2 + 50x$ for x between 5 and 17, by using a partition that consists of 4 equal subintervals of [5, 17] and using the **right** endpoint of each subinterval as a sample point.

- (a) 5490
- (b) 1830
- (c) 4482
- (d) 6165
- (e) 5004

19. Suppose you estimate the area under the graph of $f(x) = x^3$ from x = 7 to x = 25 by adding the areas of the rectangles as follows: partition the interval into 6 equal subintervals and use the right endpoint of each interval to determine the height of the rectangle. What is the area of the 5th rectangle?

Possibilities:

- (a) 10648
- (b) 121275
- (c) 31944
- (d) 20577
- (e) $\frac{103935}{4}$

20. Suppose you are given the following data points for a function f(x).

x	1	2	3	4
f(x)	6	12	17	18

If f is a **linear** function on each interval between the given points, find $\int_1^4 f(x) \, \mathrm{d}x$.

- (a) 47
- (b) 53
- (c) 35
- (d) 153
- (e) 41

1. Areas:

(a) Triangle
$$A = \frac{bh}{a}$$

- (a) Triangle $A = \frac{1}{2}$ (b) Circle $A = \pi r^2$
- (c) Rectangle A = lw

(d) Trapezoid
$$A = \frac{h_1 + h_2}{2} b$$

2. Volumes:

- (a) Rectangular Solid V = lwh
- (b) Sphere $V = \frac{4}{3}\pi r^3$
- (c) Cylinder $V = \pi r^2 h$

(d) Cone
$$V = \frac{1}{3}\pi r^2 h$$


MA123 — Elem. Calculus	Fall 2017	Name:	500.
Exam 3	2017-11-16		Sec.:

Do not remove this answer page — you will turn in the entire exam. No books or notes may be used. You may use an ACT-approved calculator during the exam, but NO calculator with a Computer Algebra System (CAS), networking, or camera is permitted. Absolutely no cell phone use during the exam is allowed.

The exam consists of two short answer questions and eighteen multiple choice questions. Answer the short answer questions on the back of this page, and record your answers to the multiple choice questions on this page. For each multiple choice question, you will need to fill in the circle corresponding to the correct answer. It is your responsibility to make it CLEAR which response has been chosen. For example, if (a) is correct, you must write



You have two hours to do this exam. Please write your name and section number on this page.



For grading use:

Multiple Choice	Short Answer
(number right) (5 points each)	(out of 10 points)

Total	
	(out of 100 points)

GOOD LUCK!