· · · · · · · · · · · · · · · · · · ·						
MA123 — Elem. Calculus Exam 3	Spring 2017 2017-04-13	Name: _	SOLUTIONS Sec.:			
You may use an ACT-approved	calculator during t	che exam, bu	e exam. No books or notes may be used. t NO calculator with a Computer Algebra ely no cell phone use during the exam is			
answer questions on the back of this page. For each multiple cho	f this page, and re ice question, you w	cord your an	nultiple choice questions. Answer the short swers to the multiple choice questions on l in the circle corresponding to the correct onse has been chosen. For example, if (a)			
	(a) (b)	(c) (d) (e)				
You have two hours to do this e	xam. Please write	your name or	n this page, and at the top of page three.			
GOOD LUCK!						
3. (a) (b)) c d e	12. (a b c d e			
4. (a) (b		13. (a b c d e			
5. (a) (b) c d e	14. (a b c d e			

7.	(a) (b) (c) (d) (e)	16. (a) (b) (c) (d) (e)
8.	(a) (b) (c) (d) (e)	17. (a) (b) (c) (d) (e)
9.	(a) (b) (c) (d) (e)	18. (a) (b) (c) (d) (e)
10.	(a) (b) (c) (d) (e)	19. (a) (b) (c) (d) (e)

15. (a) (b) (c) (d) (e)

(a) (b) (c) (d) (e)

11. (a) (b) (c) (d) (e) 20. (a) (b) (c) (d)

For grading use:


Multiple Choice		Short Answer		
	·			
(number right)	(5 points each)	(out of 10 points)		


Total (total 100 points)

Spring 2017 Exam 3 Short Answer Questions

Write answers on this page. Your work must be clear and legible to be sure you will get full credit.

1. Sketch the graph of a **continuous** function y = f(x) for which f is increasing on $(-\infty,4)$, f''(x) < 0 on $(-\infty,2)$ and $(7,\infty)$; f''(x) < 0 on (2,7).





2. Suppose we know two nonnegative numbers x and y satisfying 2x + y = 13. Find the maximum possible value of their product xy. You must clearly use calculus to find and justify your answer. Your final answer does not need to be simplified.

Maximum possible product:

Let P=xy

Get P in terms of one variable by using 2x+y=13 to substitute

y=13-2x

 $P = \times (13-2x) = 13x - 2x^2$

Take derivative and find crit. points

$$p' = 13-4x$$

$$0 = 13-4x$$

$$4x = 13$$

$$x = \frac{13}{4}$$

$$p' \text{ is never undefined}$$

Method 1: Use closed interval $[0, \frac{13}{2}]$ P(0) = 0 $P(\frac{13}{2}) = 0$ $P(\frac{13}{4}) = \frac{13}{4}(13 - 3(\frac{13}{4}))$ $= \frac{13}{4}(\frac{13}{2}) = \frac{169}{8} \leftarrow \text{maximum ratue}$

Method 2: Sign chart for P)

P(1) = 13-4(1)=9

P(10)-13-4(10)=-27

Sign of P) + 4

P (incr.) Max. occurs at $x = \frac{13}{4}$ since P changes from incr. to decr.

plug this into P to get the maximum value of product

Multiple Choice Questions

Show all your work on the page where the question appears. Clearly mark your answer both on the cover page on this exam and in the corresponding questions that follow.

3. Where is the function $f(t) = t^3 + 6t^2 - 36t + 8$ decreasing?

Possibilities:

(a)
$$t < -2$$

=9


(b)
$$f(t)$$
 is always decreasing f' is never undefined
$$(c) -6 < t < 2$$


(d)
$$t > -2$$

$$0 = 3t^2 + 12t - 36$$
 (Divide by 3) $f^3(3) = (3+6)(3-2)$

(e)
$$t < -6$$
 and $t > 2$

$$0 = t^2 + 4t - 12$$





4. Where is the function
$$f(t) = \frac{1}{t-51}$$
 concave up?
$$f'(t) = -(t-51)^{-2}$$
Possibilities:
$$f(t) = (t-51)^{-1}$$

(a) f(t) is never concave up

(b) -1 < t < 51

(c) t < 51

$$(d) \ t > 51$$

$$0 = \frac{2}{(4-61)^3}$$

 $G = \frac{2}{(t-61)^3}$ (Multiply both sides by denom.)

(e) f(t) is always concave up except at t = 51

$$f''(50) = \frac{2}{-1} = -2$$

5. Suppose the derivative of g(t) is $g'(t) = 11(t-3)^2(t-7)$. For t in which interval(s) is g increasing?

Possibilities:

when 9(4)>0

(a)
$$(-\infty,3) \cup (7,\infty)$$

(a)
$$(-\infty, 3) \cup (7, \infty)$$
 $0 = 11 (t-3)^2 (t-7)$


(b)
$$(-\infty, 7)$$

(c)
$$(3,7)$$

(d)
$$(3,7) \cup (11,\infty)$$

t=3,7

(d)
$$(3,7) \cup (11,\infty)$$
 g) is never undefined



$$g^{1}(2) = 11(-1)^{2}(-5)$$

= $(+)(+)(-)$

$$g'(5) = 11(2)^{2}(-2)$$

= $(+)(+)(-)$

$$g'(8) = 11(5)^{2}(1)$$

= (4)(+)(+)

6. Suppose the derivative of g(t) is $g'(t) = 81 - t^2$. Where is the function g(t) concave up?


(a)
$$f(t)$$
 is always concave up

(a)
$$f(t)$$
 is always concave

(b)
$$t < -9$$
 and $t > 9$

$$(d) -9 < t < 9$$

(e)
$$t > 0$$




$$g''(1)=-2$$
 \Longrightarrow g is conc. up on $(-\infty,0)$

7. The following is the graph of the derivative, f'(x), of the function f(x). Where is the original function f(x) decreasing?

f is decreasing when f) is negative




- This happens on (-00,-3) (a) nowhere
- (b) $(-\infty, -3)$
- (c) everywhere
- (d) $(2,\infty)$
- (e) $(-3, \infty)$




8. The following is the graph of the derivative, f'(x), of the function f(x). Where is the original function f(x) concave up?


- f is concave up when
- (a) nowhere
- f' is increasing which happens everywhere
- (b) $(2,\infty)$
- (c) $(-\infty, -3)$
- (d) $(-3,\infty)$
- ((e) everywhere





9. A farmer builds a rectangular pen with 8 vertical partitions (9 vertical sides) using 400 feet of fencing. What is the maximum possible total area of the pen?

X









$$A^{1}(1) = 191$$
 $A^{1}(30) = -70$
 $A^{2}(1) = 191$ $A^{3}(30) = -70$
 $A^{3}(1) = 191$ $A^{3}(30) = -70$

10. A car rental agency rents 190 cars per day at a rate of \$27 dollars per day. For each 1 dollar increase in the daily rate, 3 fewer cars are rented. At what rate should the cars be rented to produce maximum income? R= revenue R=NP

(e) \$45.97 per day Slope =
$$-\frac{3}{1} = -3$$

equation:
$$N-190=-3(P-27)=-3P+81$$

 $N=-3P+271$

$$R = (-3P + 271) P = -3P^{2} + 271 P$$

$$R' = -6P + 271 + \text{this is never ordefined}$$

$$O = -6P + 271 + 6P = 271 + P = \frac{271}{6} \approx 45.45$$

$$P = rate per car$$
 $27, 190)$
 $P = 1$
 271
 $P = 50$
 $R'(1) = 265$
 $R'(50) = -29$

$$\Rightarrow$$
 R has a max. at $P = \frac{271}{6}$

11. Find the critical numbers of the function $f(x) = xe^{13x+3}$. Product Rule and Chain Rule.

Possibilities:

(a)
$$-\frac{3}{13}$$
, 0

(c)
$$-\frac{1}{13}$$
, 0, e^{13}

$$(d) -\frac{1}{13}$$

 $f'(x) = \chi \left(e^{13x+3}\right)^{1} + e^{13x+3}(\chi)^{1}$

 $f'(x) = x e^{i3x+3}(i3) + e^{i3x+3}(i)$

Set
$$f'(x) = 0$$

 $0 = 13xe^{13x+3} + e^{13x+3}$

$$Q = e^{13x+3} (13x+1)$$


$$\Rightarrow 13x+1=0$$

 $13x=-1 \Rightarrow x=-\frac{1}{13}$

12. Given the function $f(x) = \begin{cases} -x & \text{if } x < 0 \\ x & \text{if } x \ge 0 \end{cases}$ evaluate the definite integral




$$(c)$$
 850




13. The graph of y = f(x) shown below consists of straight lines. Evaluate the definite integral $\int_{-3}^3 f(x) \, \mathrm{d}x.$

- (a) 7.5
- $= \frac{1}{2}(4)(3) (1)(3) \frac{1}{2}(1)(3)$
- (b) 2.5
- = 6-3-3 (c) 6
- (d) 21.5
- = 1.5 (e) 1.5



14. The graph of y = f(x) shown below includes semicircles and a straight line. Evaluate the definite integral $\int_{-4}^{2} f(x) dx$. Use $\pi = 3.14$.

- $= A_a A_1$ = 17(1)2 = 7(2)2 (a) 7.85
- (b) -9.42
- (c) -4.71 = T = 4T = 2 = 2
- =-317 2-4.71 (e) 4.71



15. Suppose that $\int_{15}^{17} f(x) dx = 22$ and $\int_{6}^{17} f(x) dx = 7$. Find the value of $\int_{6}^{15} f(x) dx$.

$$\int_{0}^{15} \int_{0}^{17} \int_{0}^{17$$

$$\begin{array}{ccc}
\hline
\text{(c)} & -\frac{5}{3} & = & 7 - \\
\text{(d)} & \text{20} & = & -15
\end{array}$$

- (d) -29
- (e) 15

- (c) 544
- (d) 94
- (e) 96

16. Suppose that
$$\int_{7}^{23} f(x) dx = 9$$
 and $\int_{7}^{23} g(x) dx = 19$. Find the value of $\int_{7}^{23} (2f(x) + 4g(x)) dx$.

Possibilities:

(a) 1504

(b) 8

$$= 2(9) + 4(19)$$

$$(a) -\frac{25}{2}$$

(b)
$$\frac{55}{2}$$

17. Find the average value of
$$f(x)$$
 on the interval [3, 11] given that $f(x) = \begin{cases} 90 & \text{if } x < 6 \\ -10 & \text{if } x \ge 6. \end{cases}$
Possibilities:

(a) $-\frac{25}{2}$

(b) Possibilities:

(a) $-\frac{25}{2}$

(b) Possibilities:


$$=\int f(x)dx$$

$$\frac{\int f(x) dx}{8} = \frac{\int f(x) dx + \int f(x) dx}{8} = \frac{90(3) - 10(5)}{8}$$

18. Estimate the area under the graph of
$$y = -x^2 + 30x$$
 for x between 2 and 14, by using a partition that consists of 4 equal subintervals of $[2,14]$ and using the **right** endpoint of each subinterval as a sample point.

Possibilities:

- (a) 2370
- (b) 2202
- (c) 734
- (d) 1698
- (e) 1968





176


10

19. Suppose you estimate the area under the graph of $f(x) = \frac{1}{x}$ from x = 5 to x = 45 by adding the

areas of the rectangles as follows: partition the interval into 20 equal subintervals and use the right endpoint of each interval to determine the height of the rectangle. What is the area of the 10th rectangle?







Area of 10th rectangle
= (base) (height)
=
$$2 + (25)$$

= $2 + (25)$
= $2 + (25)$
= $2 + (25)$

20. Suppose you are given the following data points for a function f(x).

5

40

If f is a linear function on each interval between the given points, find $\int_1^4 f(x) dx$.

Possibilities:


(a) 41

(b) 45

(c) 139

(d) 33

(e) 25



Area of 3 Trapezoids =
$$T_1 + T_2 + T_3$$

= $(1)(\frac{4+8}{2}) + (1)(\frac{8+13}{2}) + (1)(\frac{13+20}{2})$
= $\frac{12}{2} + \frac{21}{2} + \frac{33}{2}$
= $\frac{66}{2} = 33$

Some Formulas

1. Areas:

- (a) Triangle $A = \frac{bh}{2}$
- (b) Circle $A = \pi r^2$
- (c) Rectangle A = lw
- (d) Trapezoid $A = \frac{h_1 + h_2}{2} b$

2. Volumes:

- (a) Rectangular Solid V = lwh
- (b) Sphere $V = \frac{4}{3}\pi r^3$
- (c) Cylinder $V = \pi r^2 h$
- (d) Cone $V = \frac{1}{3}\pi r^2 h$

MA123 — Elem. 6 Exam 3	Calculus	Spring 2017 2017-04-13	Name:		Sec.:
You may use an ACT	-approved	calculator during	the exam, but N	O calcula	books or notes may be used. ator with a Computer Algebra bhone use during the exam is
answer questions on this page. For each m	the back of ultiple cho ponsibility	this page, and redice question, you v	ecord your answ will need to fill in	ers to the	ce questions. Answer the short e multiple choice questions on le corresponding to the correct en chosen. For example, if (a)
		(a) (b)	(c) (d) (e)		
You have two hours to	o do this e	xam. Please write	your name on t	his page,	and at the top of page three.
		GOO	D LUCK!		
3.	(a) (b)	(c) (d) (e)	12. a	(b) (c) d e
4.	(a) (b)	(c) (d) (e)	13. (a	(b) (c) d e
5.	(a) (b)	(c) (d) (e)	14. (a	b c) d e
6.	(a) (b)	(c) (d) (e)	15. (a	b c) d e
7.	(a) (b)	(c) (d) (e)	16. (a)	(b) (c) d (e)
8.	(a) (b)	(c) (d) (e)	17. (a)	b c) (d) (e)
9.	(a) (b)	(c) (d) (e)	18. (a)	(b) (c)) (d) (e)
10). (a) (b)	$\begin{array}{c} (c) & (d) & (e) \end{array}$	19. (a)	(b)	(\mathbf{d})
11.	(a) (b)	(c) (d) (e)	20. (a	(b) (c) d e
		For gr	rading use:		
Multiple	Choice	Short Answer		Total	
(number right)	(5 points each)	(out of 10 points)			(total 100 points)