

MA137 – Calculus 1 with Life Science Applications

Exponential and Logarithmic Functions

(Sections 1.1 and 1.2)

Alberto Corso

`<alberto.corso@uky.edu>`

Department of Mathematics
University of Kentucky


August 30, 2017

Exponential Functions


The **exponential function**

$$f(x) = a^x \quad (a > 0, a \neq 1)$$

has domain \mathbb{R} and range $(0, \infty)$. The graph of $f(x)$ has one of these shapes:



$$f(x) = a^x \quad \text{for } a > 1$$



$$f(x) = a^x \\ \text{for } 0 < a < 1$$

Laws of Exponents

Let a and b be real numbers so that $a, b > 0$ and $a, b \neq 1$.

- $a^0 = 1$

- $a^u a^v = a^{u+v}$

- $\frac{a^u}{a^v} = a^{u-v}$ In particular, $\frac{1}{a^v} = \frac{a^0}{a^v} = a^{0-v} = a^{-v}$

- $(a^u)^v = a^{uv}$ In particular, $a^{1/n} = \sqrt[n]{a}$

- $(ab)^u = a^u b^u$ $\left(\frac{a}{b}\right)^u = \frac{a^u}{b^u}$

Example 1:

Use the graph of $f(x) = 3^x$ to sketch the graph of each function:

$$g(x) = -3^x$$

$$h(x) = 1 - 3^{-x}$$

The Number 'e' (Euler's constant)

The most important base is the number denoted by the letter e .

The number e is defined as the value that $\left(1 + \frac{1}{n}\right)^n$ approaches as n becomes very large.

Correct to five decimal places (note that e is an irrational number), $e \approx 2.71828$.

n	$\left(1 + \frac{1}{n}\right)^n$
1	2.00000
5	2.48832
10	2.59374
100	2.70481
1,000	2.71692
10,000	2.71815
100,000	2.71827
1,000,000	2.71828

The Natural Exponential Function

The Natural Exponential Function

The **natural exponential function** is the exponential function

$$f(x) = e^x$$

with base e . It is often referred to as the exponential function.


Note:

Sometimes we write

$$f(x) = \exp(x)$$

to denote the exponential function.

Since $2 < e < 3$, the graph of $y = e^x$ lies between the graphs of $y = 2^x$ and $y = 3^x$.



Example 2:

When a certain drug is administered to a patient, the number of milligrams remaining in the patient's bloodstream after t hours is modeled by

$$D(t) = 50 e^{-0.2t}.$$

How many milligrams of the drug remain in the patient's bloodstream after 3 hours?

Logarithmic Functions

Every exponential function $f(x) = a^x$, with $0 < a \neq 1$, is a one-to-one function (Horizontal Line Test). Thus, it has an inverse function, called the *logarithmic function with base a* and denoted by $\log_a x$.

Definition

Let a be a positive number with $a \neq 1$. The **logarithmic function** with base a , denoted by \log_a , is defined by

$$y = \log_a x \iff a^y = x.$$


That is, $\log_a x$ is the exponent to which a must be raised to give x .

Properties of Logarithms

1. $\log_a 1 = 0$
2. $\log_a a = 1$
3. $\log_a a^x = x$
4. $a^{\log_a x} = x$

Graphs of Logarithmic Functions

The graph of $f^{-1}(x) = \log_a x$ is obtained by reflecting the graph of $f(x) = a^x$ in the line $y = x$. Thus, the function $y = \log_a x$ is defined for $x > 0$ and has range equal to \mathbb{R} .



The point $(1, 0)$ is on the graph of $y = \log_a x$ (as $\log_a 1 = 0$) and the y -axis is a vertical asymptote.

Natural Logarithms

Of all possible bases a for logarithms, it turns out that the most convenient choice for the purposes of Calculus is the number e .

Definition

The logarithm with base e is called the **natural logarithm** and denoted:

$$\ln x := \log_e x.$$

We recall again that, by the definition of inverse functions, we have

$$y = \ln x \quad \iff \quad e^y = x.$$

Properties of Natural Logarithms

1. $\ln 1 = 0$

2. $\ln e = 1$

3. $\ln e^x = x$

4. $e^{\ln x} = x$

Common Logarithms

Another convenient choice of base for the purposes of the Life Sciences is the number 10.

Definition

The logarithm with base 10 is called the **common logarithm** and denoted:

$$\log x := \log_{10} x.$$

We recall again that, by the definition of inverse functions, we have

$$y = \log x \quad \iff \quad 10^y = x.$$

Properties of Natural Logarithms

1. $\log 1 = 0$
2. $\log 10 = 1$
3. $\log 10^x = x$
4. $10^{\log x} = x$

Laws of Logarithms

Since logarithms are 'exponents', the Laws of Exponents give rise to the Laws of Logarithms:

Laws of Logarithms

Let a be a positive number, with $a \neq 1$. Let A , B and C be any real numbers with $A > 0$ and $B > 0$.

1. $\log_a(AB) = \log_a A + \log_a B$;
2. $\log_a\left(\frac{A}{B}\right) = \log_a A - \log_a B$;
3. $\log_a(A^C) = C \log_a A$.

Proof of Law 1.: $\log_a(AB) = \log_a A + \log_a B$

Let us set

$$\log_a A = u \quad \text{and} \quad \log_a B = v.$$

When written in exponential form, they become

$$a^u = A \quad \text{and} \quad a^v = B.$$

$$\begin{aligned} \text{Thus:} \quad \underline{\log_a(AB)} &= \log_a(a^u a^v) \\ &= \log_a(a^{u+v}) \\ &\stackrel{\text{why?}}{=} u + v \\ &= \underline{\log_a A + \log_a B}. \end{aligned}$$

In a similar fashion, one can prove **2.** and **3.**

Expanding and Combining Logarithmic Expressions

Example 3:

Use the Laws of Logarithms to combine the expression

$$\log_a b + c \log_a d - r \log_a s - \log_a t$$

into a single logarithm.

Change of Base

For some purposes, we find it useful to change from logarithms in one base to logarithms in another base. One can prove that:

$$\log_b x = \frac{\log_a x}{\log_a b}.$$

Proof: Set $y = \log_b x$. By definition, this means that $b^y = x$. Apply now $\log_a(\cdot)$ to $b^y = x$. We obtain

$$\log_a(b^y) = \log_a x \quad \rightsquigarrow \quad y \log_a b = \log_a x.$$

Thus

$$\log_b x = y = \frac{\log_a x}{\log_a b}.$$

Example:

$$\log_5 2 = \frac{\log 2}{\log 5} = \frac{\ln 2}{\ln 5} \approx 0.43068.$$

Exponential Equations

An exponential equation is one in which the variable occurs in the exponent. For example,

$$3^{x+2} = 7.$$

We take the (either common or natural) logarithm of each side and then use the Laws of Logarithms to 'bring down the variable' from the exponent:

$$\log(3^{x+2}) = \log 7$$

$$\rightsquigarrow (x + 2) \log 3 = \log 7$$

$$\rightsquigarrow x + 2 = \frac{\log 7}{\log 3}$$

$$\rightsquigarrow x = \frac{\log 7}{\log 3} - 2 \approx -0.228756$$

Example 4: (Online Homework HW03, # 6)

Solve the given equation for x:

$$2^{5x-4} = 3^{10x-10}$$

Logarithmic Equations

A logarithmic equation is one in which a logarithm of the variable occurs. For example,

$$\log_2(25 - x) = 3.$$

To solve for x , we write the equation in exponential form, and then solve for the variable:

$$25 - x = 2^3 \quad \rightsquigarrow \quad 25 - x = 8 \quad \rightsquigarrow \quad x = 17.$$

Alternatively, we raise the base, 2, to each side of the equation; we then use the Laws of Logarithms:

$$2^{\log_2(25-x)} = 2^3 \quad \rightsquigarrow \quad 25 - x = 2^3 \quad \rightsquigarrow \quad x = 17.$$

Example 5: (Online Homework HW03, # 5)

Solve the given equation for x :

$$\log_{10} x + \log_{10}(x + 21) = 2$$