

# MA 137 — Calculus 1 with Life Science Applications

## The Definite Integral

(Section 6.1)

**Alberto Corso**  
(alberto.corso@uky.edu)

Department of Mathematics  
University of Kentucky

December 1, 2017

## Sigma ( $\Sigma$ ) Notation

In approximating areas we have encountered sums with many terms. A convenient way of writing such sums uses the Greek letter  $\Sigma$  (which corresponds to our capital S) and is called *sigma notation*. More precisely, if  $a_1, a_2, \dots, a_n$  are real numbers we denote the sum

$$a_1 + a_2 + \dots + a_n$$

by using the notation

$$\sum_{k=1}^n a_k.$$

The integer  $k$  is called an *index* or *counter* and takes on the values  $1, 2, \dots, n$ . For example,

$$\sum_{k=1}^6 k^2 = 1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2 = 1 + 4 + 9 + 16 + 25 + 36 = 91$$

whereas

$$\sum_{k=3}^6 k^2 = 3^2 + 4^2 + 5^2 + 6^2 = 9 + 16 + 25 + 36 = 86.$$

## Summation Rules

The rules and formulas given next allow us to compute fairly easily Riemann sums where the number  $n$  of subintervals is rather large. We can also get compact and manageable expressions for the sum so that we can readily investigate what happens as  $n$  approaches infinity.

$$[\text{sr}_1] \quad \sum_{k=1}^n c = nc \qquad [\text{sr}_2] \quad \sum_{k=1}^n (c a_k) = c \sum_{k=1}^n a_k$$

$$[\text{sr}_3] \quad \sum_{k=1}^n (a_k \pm b_k) = \sum_{k=1}^n a_k \pm \sum_{k=1}^n b_k$$

**Note:** The summations rules are nothing but the usual rules of arithmetic rewritten in the  $\Sigma$  notation.

For example,  $[\text{sr}_2]$  is nothing but the distributive law of arithmetic  $c a_1 + c a_2 + \dots + c a_n = c (a_1 + a_2 + \dots + a_n)$ ;

$[\text{sr}_3]$  is nothing but the commutative law of addition  $(a_1 \pm b_1) + \dots + (a_n \pm b_n) = (a_1 + \dots + a_n) \pm (b_1 + \dots + b_n)$ .

## Formulas [Neuhauser, Example #3 (p. 279); Problem # 31 (p. 291)]

$$[\text{sf}_1] \quad \sum_{k=1}^n k = \frac{n(n+1)}{2} \qquad [\text{sf}_2] \quad \sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$$

**Proof:** In the case of  $[\text{sf}_1]$ , let  $S$  denote the sum of the integers  $1, 2, 3, \dots, n$ . Let us write this sum  $S$  twice: we first list the terms in the sum in increasing order whereas we list them in decreasing order the second time:

$$\begin{array}{r} S = 1 + 2 + \dots + n \\ S = n + n-1 + \dots + 1 \end{array}$$

If we now add the terms along the vertical columns, we obtain

$$2S = \underbrace{(n+1) + (n+1) + \dots + (n+1)}_{n \text{ times}} = n(n+1).$$

This gives our desired formula, once we divide both sides of the above equality by 2.

In the case of  $[\text{sf}_2]$ , let  $S$  denote the sum of the integers  $1^2, 2^2, 3^2, \dots, n^2$ . The *trick* is to consider the sum

$$\sum_{k=1}^n [(k+1)^3 - k^3]. \quad \text{On the one hand, this new sum collapses to}$$

$$(2^3 - 1^3) + (3^3 - 2^3) + (4^3 - 3^3) + \dots + (n^3 - (n-1)^3) + ((n+1)^3 - n^3) = (n+1)^3 - 1^3 = n^3 + 3n^2 + 3n$$

On the other hand, using our summation rules together with  $[\text{sf}_1]$  gives us

$$\sum_{k=1}^n [(k+1)^3 - k^3] = \sum_{k=1}^n [3k^2 + 3k + 1] = 3 \sum_{k=1}^n k^2 + 3 \sum_{k=1}^n k + \sum_{k=1}^n 1 = 3S + 3 \frac{n(n+1)}{2} + n$$

Equating the right hand sides of the above identities gives us:  $3S + 3 \frac{n(n+1)}{2} + n = n^3 + 3n^2 + 3n$ .

If we solve for  $S$  and properly factor the terms, we obtain our desired expression.

## More Formulas

The next formulas can be verified in a sequential order using the same type of trick used in the proof for [sf<sub>2</sub>]. The proofs get increasingly more tedious.

$$[\text{sf}_3] \quad \sum_{k=1}^n k^3 = \frac{n^2(n+1)^2}{4}$$

$$[\text{sf}_4] \quad \sum_{k=1}^n k^4 = \frac{n(n+1)(2n+1)(3n^2+3n-1)}{30}$$

5/17

## Example 1: (Online Homework, HW 23, # 15)

Find the numerical value of the sums below:

$$\bullet \sum_{j=3}^7 (4j - 1)$$

$$\bullet \sum_{i=3}^5 (i^2 - i)$$

6/17

$$\begin{aligned} \sum_{j=3}^7 (4j-1) &= [4 \cdot \underline{3} - 1] + [4 \cdot \underline{4} - 1] + [4 \cdot \underline{5} - 1] + [4 \cdot \underline{6} - 1] + [4 \cdot \underline{7} - 1] \\ &= 11 + 15 + 19 + 23 + 27 = 95 \end{aligned}$$

$$\begin{aligned} \sum_{i=3}^5 (i^2 - i) &= [3^2 - 3] + [4^2 - 4] + [5^2 - 5] \\ &= 6 + 12 + 20 = 38 \end{aligned}$$

## Example 2:

Find the numerical value of the sums below:

$$\bullet \sum_{j=3}^n (4j - 1)$$

$$\bullet \sum_{i=3}^n (i^2 - i)$$

7/17

$$\begin{aligned}
\sum_{j=3}^n (4j-1) &= \sum_{j=1}^n (4j-1) - \sum_{j=1}^2 (4j-1) \\
&= \sum_{j=1}^n (4j-1) - \underbrace{[(4 \cdot 1 - 1) + (4 \cdot 2 - 1)]}_{=10} \\
&= 4 \sum_{j=1}^n j - \sum_{j=1}^n 1 - 10 \\
&= 4 \frac{n(n+1)}{2} - n - 10 \\
&= 2n(n+1) - n - 10 \\
&= 2n^2 + 2n - n - 10 \\
&= \boxed{2n^2 + n - 10} \quad \text{||| m}
\end{aligned}$$

$$\begin{aligned}
\sum_{i=3}^n (i^2 - i) &= \sum_{i=1}^n (i^2 - i) - \sum_{i=1}^2 (i^2 - i) \\
&= \sum_{i=1}^n (i^2 - i) - \underbrace{[(1^2 - 1) + (2^2 - 2)]}_{=0+2=2} \\
&= \left( \sum_{i=1}^n i^2 \right) - \left( \sum_{i=1}^n i \right) - 2 \\
&\stackrel{\text{use the rules}}{=} \frac{n(n+1)(2n+1)}{6} - \frac{n(n+1)}{2} - 2 \\
&\stackrel{\text{simplify}}{=} \frac{2n^3 + 3n^2 + n - 3n(n+1) - 12}{6} \\
&= \frac{2n^3 + \cancel{3n^2} + n - \cancel{3n^2} - 3n - 12}{6} = \frac{2n^3 - 2n - 12}{6} \\
&= \frac{n^3 - n - 6}{3} \quad \text{||| l.}
\end{aligned}$$

## Back to the Area Problem: Partitions

The idea we have used so far is to “to partition” or subdivide the given interval  $[a, b]$  into smaller subintervals on each of which the variable  $x$ , and thus the function  $f(x)$ , does not change much.

### Definition of a Partition

A *partition* of an interval  $[a, b]$  is a set of points  $\{x_0, x_1, x_2, \dots, x_{n-1}, x_n\}$ , listed increasingly, on the  $x$ -axis with  $a = x_0$  and  $x_n = b$ . That is:

$$a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b.$$

These points subdivide the interval  $[a, b]$  into  $n$  subintervals

$$[a, x_1], [x_1, x_2], [x_2, x_3], \dots, [x_{n-1}, b].$$

The  $k$ -th subinterval is thus of the form  $[x_{k-1}, x_k]$  and it has length

$$\Delta x_k = x_k - x_{k-1}.$$

### Assumption

Set  $\|P\| = \max_{1 \leq i \leq n} \{\Delta x_i\}$ . We assume that our partition  $P$  is such that

$\|P\| \rightarrow 0$  as  $n \rightarrow \infty$ . In other words, we assume that the length of the longest (and, hence, of all) subinterval(s) tend(s) to zero whenever the number of subintervals in  $P$  becomes very large.

## The Definite Integral

Let  $f(x)$  be a function defined on an interval  $[a, b]$ .

- Partition the interval  $[a, b]$  in  $n$  subintervals of lengths  $\Delta x_1, \dots, \Delta x_n$ , respectively.
- For  $k = 1, \dots, n$  pick a representative point  $c_k$  in the corresponding  $k$ -th subinterval.

The **definite integral** of  $f$  from  $a$  to  $b$  is defined as

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n f(c_k) \cdot \Delta x_k = \lim_{\|P\| \rightarrow 0} \sum_{k=1}^n f(c_k) \cdot \Delta x_k$$

and it is denoted by  $\int_a^b f(x) dx$ .

The sum  $\sum_{k=1}^n f(c_k) \cdot \Delta x_k$  is called a *Riemann sum* in honor of the German mathematician Bernhard Riemann

(1826-1866), who developed the above ideas in full generality. The symbol  $\int$  is called the *integral sign*. It is an elongated capital S, of the kind used in the 1600s and 1700s. The letter S stands for the summation performed in computing a Riemann sum. The numbers  $a$  and  $b$  are called the *lower and upper limits of integration*, respectively. The function  $f(x)$  is called the *integrand* and the symbol  $dx$  is called the *differential* of  $x$ . You can think of the  $dx$  as representing what happens to the term  $\Delta x$  in the limit, as the size  $\Delta x$  of the subintervals gets closer and closer to 0.

- The role of  $x$  in a definite integral is the one of a *dummy variable*. In fact  $\int_a^b x^2 dx$  and  $\int_a^b t^2 dt$  have the same meaning. They represent the same number.
- We recall that a limit does not necessarily exist. However:


**Theorem**

If  $f$  is continuous on  $[a, b]$  then  $\int_a^b f(x) dx$  exists.

- As we observed earlier, it is computationally easier to partition the interval  $[a, b]$  into  $n$  subintervals of equal length. Therefore each subinterval has length  $\Delta x = \frac{b-a}{n}$  (we drop the index  $k$  as it is no longer necessary). In this case, there is a simple formula for the points of the partition, namely:  
 $x_0 = a + 0 \cdot \Delta x = a, x_1 = a + \Delta x, \dots, x_k = a + k \cdot \Delta x, \dots, x_n = a + n \cdot \Delta x = b$   
 or, more concisely,  $x_k = a + k \cdot \frac{b-a}{n}$  for  $k = 0, 1, 2, \dots, n$ .

## Definite Integrals and Areas

We stress the fact that if the function  $f$  takes on only positive values then the definite integral is nothing but the area of the region below the graph of  $f$ , lying above the  $x$ -axis, and bounded by the vertical lines  $x = a$  and  $x = b$ .




**Distance traveled by an object:**

If the positive valued function under consideration is the velocity  $v(t)$  of an object at time  $t$ , then the area underneath the graph of the velocity function and lying above the  $t$ -axis represents the total distance traveled by the object from  $t = a$  to  $t = b$ .

## What if the Function Takes on Negative Values?

If  $f$  happens to take on both positive and negative values, then the Riemann sum is the sum of the areas of rectangles that lie above the  $x$ -axis and the negatives of the areas of rectangles that lie below the  $x$ -axis. Passing to the limit, we obtain that, in general, a definite integral can be interpreted as a difference of areas:




$$\int_a^b f(x) dx = [\text{area of the region(s) lying above the } x\text{-axis}] - [\text{area of the region(s) lying below the } x\text{-axis}]$$

## Right Versus Left Endpoint Estimates

Observe that  $x_k$ , the right endpoint of the  $k$ -th subinterval, is also the left endpoint of the  $(k + 1)$ -th subinterval. Thus the Riemann sum estimate for the definite integral of a function  $f$  defined over an interval  $[a, b]$  can be written in either of the following two forms

$$\sum_{k=0}^{n-1} f(x_k) \cdot \Delta x_{k+1} \qquad \sum_{k=1}^n f(x_k) \cdot \Delta x_k$$

depending on whether we use left or right endpoints, respectively.




Left endpoint Riemann sum estimate

If we are dealing with a regular partition, the above sums become

$$\sum_{k=0}^{n-1} f(a + k \cdot \Delta x) \cdot \Delta x \qquad \sum_{k=1}^n f(a + k \cdot \Delta x) \cdot \Delta x$$

with  $\Delta x = \frac{b-a}{n}$  and  $x_k = a + k \cdot \Delta x$  for  $k = 0, 1, \dots, n$ .



Right endpoint Riemann sum estimate 13/17

**Example 3:** (Online Homework, HW 23, # 11)

Express the limit as a definite integral

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{2}{n} \left( 5 + \frac{2i}{n} \right)^{10}$$

14/17

From page 13, the formula for a Riemann sum using the right endpoints is:

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n f\left(a + k \cdot \underbrace{\frac{b-a}{n}}_{\Delta x}\right) \cdot \underbrace{\frac{b-a}{n}}_{\Delta x}$$

Hence in our case:

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \left( 5 + \frac{2i}{n} \right)^{10} \cdot \frac{2}{n}$$

says that this is  $\int_5^7 \frac{x^{10}}{f(x)} \cdot dx$

**Example 4:** (Online Homework, HW 23, # 12)

Express the limit as a definite integral

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{4}{n} \sqrt{1 + \frac{4i}{n}}$$

15/17

We can interpret  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{4}{n} \sqrt{1 + \frac{4i}{n}}$

as  $\int_0^4 \underbrace{\sqrt{1+x}}_{f(x)} dx$

(this is the type of area that WeBWork seeks)

(or  $\int_1^5 \sqrt{x} dx$  ... which are actually equivalent)


there is just an horizontal shift!

**Example 5:** (Online Homework, HW 23, # 7)

Evaluate the following integral by interpreting it in terms of areas:

$$\int_0^3 \left( \frac{1}{2}x - 1 \right) dx$$

Let's graph the function  $y = \frac{1}{2}x - 1$  on  $[0, 3]$




$$\int_0^3 \left( \frac{1}{2}x - 1 \right) dx = \boxed{-1} + \frac{1}{4} = \boxed{\underline{\underline{-\frac{3}{4}}}}$$

"signed area" as it is below the x-axis

**Example 6:** (Online Homework, HW 23, # 10)

Let  $g(x) = \int_0^x f(t) dt$ , where  $f$  is the function whose graph is shown below.




- Evaluate  $g(x)$  for  $x = 0, 1, 2, 3, 4, 5$ , and  $6$ .
- Estimate  $g(7)$ .
- At what value of  $x$  does  $g$  attain its maximum?
- At what value of  $x$  does  $g$  attain its minimum?


$$g(x) = \int_0^x f(t) dt$$

$$g(0) = \int_0^0 f(t) dt = 0$$

$$g(1) = \int_0^1 f(t) dt = \frac{1}{2}$$




$$g(2) = \int_0^2 f(t) dt = 0$$




$$g(3) = \int_0^3 f(t) dt$$

$$= \frac{1}{2} - 1 = \boxed{-\frac{1}{2}}$$



$$g(4) = \int_0^4 f(t) dt = \boxed{0}$$

$$= \frac{1}{2} - 1 + \frac{1}{2} = 0$$



$$g(5) = \boxed{\frac{3}{2}} \quad \text{and} \quad g(6) = \boxed{4} \quad \text{check}$$

$$g(7) \cong 6.2 \quad (\text{estimate})$$

Max of  $g$  occurs at  $\boxed{x=7}$ ; minimum at  $\boxed{x=3}$