

MA 138 – Calculus 2 with Life Science Applications
Integration by Parts
(Section 7.2)

Alberto Corso

⟨alberto.corso@uky.edu⟩

Department of Mathematics
University of Kentucky

Monday, January 23, 2017

Section 7.2: Integration by Parts

We saw that integration by parts is the product rule in integral form.

We also recall the following general formula:

Rule for Integration by Parts

If $f(x)$ and $g(x)$ are differentiable functions, then

$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx;$$

$$\int_a^b f(x)g'(x) dx = f(x)g(x) \Big|_a^b - \int_a^b f'(x)g(x) dx.$$

Example 1

A particle that moves along a straight line has velocity

$$v(t) = t^2 e^{-2t}$$

meters per second after t seconds.

How many meters will it travel during the first t seconds?

Using the notion of cumulative change

$$\underbrace{s(t) - s(0)}_{\text{distance traveled}} = \int_0^t \frac{ds}{du} \cdot du = \int_0^t v(u) \cdot du$$

$$= \int_0^t u^2 \cdot e^{-2u} \cdot du = \text{set } w = -2u \quad \frac{dw}{du} = -2$$
$$\boxed{u = -\frac{1}{2}w} \quad \boxed{du = -\frac{1}{2}dw}$$

Substitute and change the integration limits:

$$= \int_0^{-2t} \left(-\frac{1}{2}w\right)^2 \cdot e^w \cdot \left(-\frac{1}{2}dw\right) = -\int_0^{-2t} \frac{1}{8} w^2 e^w dw =$$

$$= \int_{-2t}^0 \frac{1}{8} w^2 e^w dw = \text{if you prefer relabel}$$

$w \longleftrightarrow x$

$$\therefore s(t) - s(0) = \int_{-2t}^0 \underbrace{\frac{1}{8} x^2}_f \underbrace{e^x}_{g'} dx = \text{integration by parts}$$

$$= \left[\frac{1}{8} x^2 \cdot e^x \right]_{-2t}^0 - \int_{-2t}^0 \frac{1}{8} 2x \cdot e^x dx$$

$$= \left(0 - \frac{1}{8} (-2t)^2 e^{-2t} \right) - \int_{-2t}^0 \frac{1}{4} x e^x dx$$

$$= -\frac{1}{2} t^2 e^{-2t} - \left\{ \left[\frac{1}{4} x e^x \right]_{-2t}^0 - \int_{-2t}^0 \frac{1}{4} e^x dx \right\}$$

integrate by parts again

$$= -\frac{1}{2} t^2 e^{-2t} - \left\{ \left(0 - \frac{1}{4} (-2t) e^{-2t} \right) - \left[\frac{1}{4} e^x \right]_{-2t}^0 \right\}$$

$$= -\frac{1}{2} t^2 e^{-2t} - \frac{1}{2} t e^{-2t} + \left[\frac{1}{4} e^0 - \frac{1}{4} e^{-2t} \right]$$

$$= \boxed{\frac{1}{4} - \left(\frac{1}{2} t^2 + \frac{1}{2} t + \frac{1}{4} \right) e^{-2t}}$$

Example 2 (Online Homework # 10)

Suppose that $f(1) = 4$, $f(4) = 6$, $f'(1) = -5$, $f'(4) = -5$,
and f'' is continuous. Find the value of

$$\int_1^4 x f''(x) dx.$$

$$\int_1^4 x f''(x) dx = ?$$

$$f(1) = 4$$

$$f(4) = 6$$

$$f'(1) = -5$$

$$f'(4) = -5$$

f'' continuous

integrate by parts:

$$\int_1^4 x \underbrace{(f')'}_{f''} dx = \left[x f' \right]_1^4 - \int_1^4 \underbrace{1 \cdot f'}_{\text{F.T.C.}} dx =$$

$$= \left[x f'(x) \right]_1^4 - \left[f(x) \right]_1^4 =$$

$$= \left[4 \cdot f'(4) - 1 \cdot f'(1) \right] - \left[f(4) - f(1) \right]$$

$$= 4(-5) - (-5) - (6 - 4) = \boxed{-17}$$

Example 3 (Problem # 8, Section 7.2, page 342)

Evaluate the indefinite integral: $\int 3xe^{-x/2} dx$.

$$\int 3x e^{-x/2} dx = \text{set } \boxed{u = -\frac{x}{2}} \quad \frac{du}{dx} = -\frac{1}{2}$$

$$\therefore \boxed{x = -2u}, \quad \boxed{-2 du = dx} \quad ; \quad \text{substitute}$$

$$= \int -6u e^u (-2 du) = \int \underbrace{12u}_f \underbrace{e^u}_{g'} du = \text{by parts}$$

$$= \underbrace{12u}_f \cdot \underbrace{e^u}_g - \int \underbrace{12}_{f'} \underbrace{e^u}_g du = 12u e^u - 12e^u + C$$

$$= \left(12\left(-\frac{x}{2}\right) - 12\right) e^{-x/2} + C = \boxed{-6(x+2)e^{-x/2} + C}$$

Example 4 (Online Homework # 7)

Find the integral: $\int_0^1 x^2 \sqrt[4]{e^x} dx.$

$$\int_0^1 x^2 \sqrt[4]{e^x} dx = \int_0^1 x^2 (e^x)^{1/4} dx = \int_0^1 x^2 e^{x/4} dx$$

Use the substitution

$$\boxed{u = \frac{x}{4}}$$

so

$$\boxed{x = 4u}$$

and $\frac{du}{dx} = \frac{1}{4}$ so

$$\boxed{4 du = dx}$$

Change also the limits of integration

$$\therefore \int_0^1 x^2 \sqrt[4]{e^x} dx = \int_0^{1/4} (4u)^2 \cdot e^u \cdot (4 du) = \int_0^{1/4} 64 u^2 \cdot e^u du$$

$$\int \underbrace{64 u^2}_{f} \cdot \underbrace{e^u}_{g'} du = \text{by parts } \underbrace{64 u^2}_{f} \cdot \underbrace{e^u}_{g} - \int \underbrace{128 u}_{f'} \cdot \underbrace{e^u}_{g} du =$$

$$= \text{by parts again } 64 u^2 e^u - \left[128 u e^u - \int 128 \cdot e^u du \right]$$

In conclusion :

$$\int_0^1 x^2 \sqrt[4]{e^x} dx = \int_0^{1/4} 64 u^2 \cdot e^u du = \left[64 u^2 e^u - 128 u e^u + 128 e^u \right]_0^{1/4}$$

$$= \left[64 (u^2 - 2u + 2) e^u \right]_0^{1/4}$$

$$= 64 \left(\frac{1}{16} - \frac{1}{2} + 2 \right) e^{1/4} - 64(2) \cdot e^0 =$$

$$= 64 \left(\frac{1 - 8 + 32}{16} \right) e^{1/4} - 128 = \underline{\underline{100 e^{1/4} - 128}}$$

$$\approx \underline{\underline{0.4025}}$$

Example 5 (Problem # 35, Section 7.2, page 343)

Evaluate the indefinite integral: $\int \frac{1}{x} \ln x \, dx.$

(1) Our text book suggests to compute $\int \frac{1}{x} \ln x \, dx$ using integration by parts.

$$\int \underbrace{\frac{1}{x}}_{g'} \cdot \underbrace{\ln x}_{f} \, dx = \underbrace{(\ln x)}_g \cdot \underbrace{\ln(x)}_f - \int \underbrace{(\ln x)}_g \cdot \underbrace{\frac{1}{x}}_{f'} \, dx$$

move this to the right-hand side

$$\therefore 2 \int \frac{1}{x} \cdot \ln x \, dx = (\ln x)^2 + C$$

$$\therefore \boxed{\int \frac{1}{x} \ln x \, dx = \frac{1}{2} (\ln x)^2 + \tilde{C}}$$

(2) Using the substitution $\boxed{u = \ln x}$ so $\boxed{du = \frac{1}{x} dx}$

$$\text{we also get: } \int \frac{1}{x} \cdot \ln x \, dx = \int u \cdot du = \frac{1}{2} u^2 + \tilde{C} = \boxed{\frac{1}{2} (\ln x)^2 + \tilde{C}}$$

Example 6 (Problem # 48, Section 7.2, page 343)

Evaluate the definite integral: $\int_0^1 x^3 \ln(x^2 + 1) dx.$

$$\int_0^1 x^3 \cdot \ln(x^2+1) dx = \text{use first the substitution } \boxed{u = x^2 + 1}$$

$$\text{so that } \frac{du}{dx} = 2x \quad \text{or} \quad \boxed{\frac{1}{2} du = x dx}$$

$$\text{and observe } \boxed{x^2 = u - 1}$$

Substitute and change the limits of integration

$$\int_0^1 x^2 \cdot \ln(x^2+1) \cdot x dx = \int_1^2 (u-1) \cdot \ln(u) \cdot \frac{1}{2} du =$$

$$= \int_1^2 \underbrace{\left(\frac{1}{2}u - \frac{1}{2}\right)}_{g'} \cdot \underbrace{\ln u}_{f} du \quad \text{by parts} \quad \left[\left(\frac{1}{4}u^2 - \frac{1}{2}u\right) \ln u \right]_1^2 - \int_1^2 \left(\frac{1}{4}u^2 - \frac{1}{2}u\right) \frac{1}{u} du$$

$$= \left[\left(\frac{1}{4}u^2 - \frac{1}{2}u\right) \ln u \right]_1^2 - \int_1^2 \left(\frac{1}{4}u - \frac{1}{2}\right) du =$$

$$= \left[\left(\frac{1}{4} \cdot \overset{0}{\cancel{2^2}} - \frac{1}{2} \cdot 2\right) \cdot \ln(2) - \left(\frac{1}{4} \cdot \overset{0}{\cancel{(1)^2}} - \frac{1}{2} \cdot (1)\right) \cdot \ln(1) \right] - \left[\frac{1}{8}u^2 - \frac{1}{2}u \right]_1^2$$

$$= - \left[\left(\frac{1}{8} \cdot 4 - 1\right) - \left(\frac{1}{8} - \frac{1}{2}\right) \right] = \boxed{\frac{1}{8}} \approx \underline{0.125}$$

Useful aside: Trigonometric addition formulas

- We also used the double angle formulas

$$\begin{aligned}\cos(2\alpha) &= \cos^2 \alpha - \sin^2 \alpha & \sin(2\alpha) &= 2 \sin \alpha \cos \alpha \\ &= 2 \cos^2 \alpha - 1 & \text{and} & \\ &= 1 - 2 \sin^2 \alpha\end{aligned}$$

to compute $\int \cos^2 x \, dx$ and $\int \sin x \cos x \, dx$.

- Is there a 'simple' way to remember formulas of this kind?
- **Euler's formula** establishes the fundamental relationship between the trigonometric functions and the complex exponential function. It states that, for any real number x ,

$$e^{ix} = \cos x + i \sin x,$$

where i is the imaginary unit ($i^2 = -1$).

- For any α and β , using Euler's formula, we have

$$\begin{aligned}\cos(\alpha + \beta) + i \sin(\alpha + \beta) &= e^{i(\alpha + \beta)} \\ &= e^{i\alpha} \cdot e^{i\beta} \\ &= (\cos \alpha + i \sin \alpha) \cdot (\cos \beta + i \sin \beta) \\ &= (\cos \alpha \cos \beta + i^2 \sin \alpha \sin \beta) \\ &\quad + i(\sin \alpha \cos \beta + \cos \alpha \sin \beta).\end{aligned}$$

- Thus, by comparing the terms, we obtain

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta.$$

- Thus, by setting $\alpha = \beta$, we obtain

$$\cos(2\alpha) = \cos^2 \alpha - \sin^2 \alpha \quad \text{and} \quad \sin(2\alpha) = 2 \sin \alpha \cos \beta.$$