

MA 138 – Calculus 2 with Life Science Applications

Integration by Parts

(Section 7.2)

Alberto Corso

(alberto.corso@uky.edu)

Department of Mathematics
University of Kentucky

Friday, January 20, 2017

<http://www.ms.uky.edu/~ma138>

Lecture 4

About Example 4 from the previous lecture

Last time we integrated $\int \frac{3}{3+e^x} dx$ by using the substitution $u = 3 + e^x$.

This lead to $du = e^x dx = (u - 3) dx$. Thus

$$\int \frac{3}{3+e^x} dx \iff \int \frac{3}{u} \cdot \frac{du}{u-3} = \int \frac{3}{u(u-3)} du.$$

A natural question to ask is:

“Why should I care about integrals of this form?”

Next, I will give you a good reason.

We will study more systematically integrals of this form in Section 7.3.

<http://www.ms.uky.edu/~ma138>

Lecture 4

The Logistic Growth Model

In Sections 3.3 and 4.1 you should have introduced the logistic growth model. In this growth model it is assumed that the population size $N(t)$ at time t satisfies the initial value problem

$$\frac{dN}{dt} = rN \left(1 - \frac{N}{K}\right) \quad N(0) = N_0,$$

where r (=growth rate) and K (=carrying capacity) are positive constants.


Rewriting this differential equation as

$$\frac{1}{N} \frac{dN}{dt} = r \left(1 - \frac{N}{K}\right)$$

says that the per capita growth rate in the logistic equation is a linearly decreasing function of population size.

<http://www.ms.uky.edu/~ma138>

Lecture 4



In Chapter 8 we will see that in order to solve the logistic differential equation we first separate the variables to obtain


$$\frac{1}{N(1 - N/K)} dN = r dt.$$

Then we integrate both sides with respect to N and t

$$\int \frac{K}{N(N - K)} dN = \int -r dt.$$

After several calculations we obtain that the solution of the IVP is

$$N(t) = \frac{K}{1 + (K/N_0 - 1)e^{-rt}}.$$



<http://www.ms.uky.edu/~ma138>

Lecture 4

Section 7.2: Integration by Parts

Integration by parts is the product rule in integral form.

Let $f = f(x)$ and $g = g(x)$ be differentiable functions. Then, differentiating the product fg with respect to x yields

$$(fg)' = f'g + fg'$$

or, after rearranging,

$$fg' = (fg)' - f'g.$$

Integrating both sides with respect to x , we find that

$$\int fg' dx = \int (fg)' dx - \int f'g dx.$$

Since fg is an antiderivative of $(fg)'$, it follows that

$$\int (fg)' dx = fg + C.$$

Therefore

$$\int fg' dx = fg - \int f'g dx.$$

(Note that the constant C can be absorbed into the indefinite integral on the right-hand side.) Because $f' = df/dx$ and $g' = dg/dx$, we can also write the preceding equation in the short form

$$\int f dg = fg - \int g df.$$

We summarize this discussion, by stating the following **general rule**:

Rule for Integration by Parts

If $f(x)$ and $g(x)$ are differentiable functions, then

$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx;$$

$$\int_a^b f(x)g'(x) dx = f(x)g(x) \Big|_a^b - \int_a^b f'(x)g(x) dx.$$

Example 1 (Problem #61, Section 7.2, page 343)

Evaluate the indefinite integral: $\int \ln x dx$.

Integration by parts says that

$$\int fg' dx = fg - \int f'g dx$$

In our case:

$$\int \ln x dx = \int \underbrace{(\ln x)}_f \cdot \underbrace{1}_{g'} dx$$

Hence:

$$\int \ln x dx = \underbrace{x}_g \cdot \underbrace{\ln x}_f - \int \underbrace{\frac{1}{x}}_{f'} \cdot x dx$$

$$= \int \ln x dx = x \cdot \ln x - \int 1 \cdot dx = \underline{x \cdot \ln x - x + C}$$

Example 2 (Online Homework # 9)

If $g(1) = -5$, $g(5) = 2$ and $\int_1^5 g(x) dx = -10$, evaluate

$$\int_1^5 x g'(x) dx.$$

$$\int_1^5 x g'(x) dx = ?$$

$$g(1) = -5$$

$$g(5) = 2$$

$$\int_1^5 g(x) dx = -10$$

integration by parts:

$$\int_1^5 x g'(x) dx = \left[x \cdot g(x) \right]_1^5 - \int_1^5 \underbrace{1}_{\text{derivative of } f(x)=x} \cdot g(x) dx =$$

$$= 5 \cdot g(5) - 1 \cdot g(1) - \underbrace{\int_1^5 g(x) dx}_{-10} =$$

$$= 5 \cdot 2 - 1 \cdot (-5) - (-10) = 10 + 5 + 10 = \underline{25}$$

Example 3 (Online Homework # 3)

Evaluate the indefinite integral: $\int e^{4x} \sin(6x) dx.$

$$\int \underbrace{e^{4x}}_{g'} \cdot \underbrace{\sin(6x)}_f dx = \left(\frac{1}{4} e^{4x} \right) \cdot \sin(6x) - \int \frac{1}{4} e^{4x} \cos(6x) \cdot 6 dx$$

$$\therefore \int e^{4x} \sin(6x) dx = \frac{1}{4} e^{4x} \cdot \sin(6x) - \frac{3}{2} \int e^{4x} \cos(6x) dx$$

now what? let's try integration by parts again

$$= \frac{1}{4} e^{4x} \sin(6x) - \frac{3}{2} \left[\frac{1}{4} e^{4x} \cos(6x) - \int \left(\frac{1}{4} e^{4x} \right) \cdot (-\sin(6x) \cdot 6) dx \right]$$

$$= \frac{1}{4} e^{4x} \sin(6x) - \frac{3}{8} e^{4x} \cos(6x) - \frac{9}{4} \int e^{4x} \sin(6x) dx$$

Now move the integral on the right-hand side of the inequality to the left-hand side.

We can add those 2 integrals

$$\underbrace{\left(1 + \frac{9}{4}\right)}_{\frac{13}{4}} \int e^{4x} \cdot \sin(6x) dx = \frac{1}{4} e^{4x} \sin(6x) - \frac{3}{8} e^{4x} \cos(6x) + C$$

\therefore

$$\begin{aligned} \int e^{4x} \cdot \sin(6x) dx &= \frac{4}{13} \left[\frac{1}{4} e^{4x} \sin(6x) - \frac{3}{8} e^{4x} \cos(6x) \right] + \tilde{C} \\ &= \frac{1}{13} e^{4x} \sin(6x) - \frac{3}{26} e^{4x} \cos(6x) + \tilde{C} \end{aligned}$$

$$\begin{aligned} \int x^9 \cdot \cos(x^5) dx &= \text{note that we can rewrite it as follows} \\ &= \int \underbrace{x^4 \cdot x^5}_{=x^9} \cdot \cos(x^5) dx \end{aligned}$$

So, set $u = x^5$ and observe that $\frac{du}{dx} = 5x^4$
so $\frac{1}{5} du = x^4 dx$. Thus

$$\int x^9 \cos(x^5) dx = \int \underbrace{\frac{1}{5}}_f \cdot \underbrace{u \cos(u)}_{g'} du =$$

use now integration by parts!

$$= \underbrace{\frac{1}{5}}_f \cdot \underbrace{\sin(u)}_g - \int \underbrace{\frac{1}{5}}_{f'} \underbrace{\sin(u)}_g du = \frac{1}{5} u \sin(u) + \frac{1}{5} \cos(u) + C$$

Example 4 (Online Homework # 4)

Evaluate the indefinite integral: $\int x^9 \cos(x^5) dx$.

(Hint: First make a substitution and then use integration by parts to evaluate the integral.)

\therefore

$$\int x^9 \cos(x^5) dx \stackrel{u=x^5}{=} \int \frac{1}{5} u \cos(u) du$$

$$\stackrel{\text{by parts}}{=} \frac{1}{5} u \sin(u) + \frac{1}{5} \cos(u) + C$$

$$= \frac{1}{5} x^5 \sin(x^5) + \frac{1}{5} \cos(x^5) + C$$

Example 5 (Problem #3, Section 7.2, page 342)

Evaluate the indefinite integrals:

$$\int \cos^2 x \, dx \qquad \int \cos^3 x \, dx.$$

<http://www.ms.uky.edu/~ma138>

Lecture 4

Thus:

$$\int \cos^2 x \, dx = \cos x \cdot \sin x + \int \sin^2 x \, dx$$

But $\cos^2 x + \sin^2 x = 1$ so $\sin^2 x = 1 - \cos^2 x$

$$= \cos x \sin x + \int (1 - \cos^2 x) \, dx$$

$$\therefore \int \cos^2 x \, dx = \cos x \sin x + \int dx - \int \cos^2 x \, dx$$

move it to the
left-hand side

$$\therefore 2 \int \cos^2 x \, dx = \cos x \cdot \sin x + x + C$$

$$\therefore \int \cos^2 x \, dx = \frac{1}{2} \cos x \cdot \sin x + \frac{1}{2} x + \tilde{C}$$

$\int \cos^3(x) \, dx$ is easy! In fact $\cos^3(x) = \cos(x)(\cos^2 x)$
 $= \cos(x) \cdot [1 - \sin^2(x)]$

$$= \int \cos(x) \cdot [1 - \sin^2(x)] \, dx = \int (1 - u^2) \, du$$

$u = \sin x$
 $\frac{du}{dx} = \cos x \quad \therefore du = (\cos x) \, dx$

$$= u - \frac{1}{3} u^3 + C = \sin x - \frac{1}{3} \sin^3 x + C$$

$\int \cos^2(x) \, dx$ is not as easy! We need integration by parts

$$\int \cos^2(x) \, dx = \int \underbrace{\cos x}_f \cdot \underbrace{\cos x}_{g'} \, dx = \underbrace{\cos x}_f \cdot \underbrace{\sin x}_g - \int \underbrace{(-\sin x)}_{f'} \cdot \underbrace{\sin x}_g \, dx$$

Alternative method: use the double angle

formula: $\cos(2x) = 2 \cos^2 x - 1$

or $\cos^2 x = \frac{\cos(2x) + 1}{2}$

Thus:

$$\int \cos^2 x \, dx = \int \left(\frac{1}{2} \cos(2x) + \frac{1}{2} \right) dx =$$

$$= \frac{1}{2} \cdot \frac{1}{2} \sin(2x) + \frac{1}{2} x + C$$

$$= \frac{1}{4} \sin(2x) + \frac{1}{2} x + C$$

$$= \frac{2 \sin x \cos x}{4} + \frac{1}{2} x + C = \frac{1}{2} \sin x \cos x + \frac{1}{2} x + C$$