

MA 138 – Calculus 2 with Life Science Applications
Rational Functions and Partial Fractions
(Section 7.3)

Alberto Corso
`(alberto.corso@uky.edu)`

Department of Mathematics
University of Kentucky

Wednesday, January 27, 2017

<http://www.ms.uky.edu/~ma138>

Lecture 7

We need to consider 2 cases:

$a = b$ the integral becomes $\int \frac{3}{(x+a)^2} dx$

so by letting $u = x+a$; $\frac{du}{dx} = 1$ so $du = dx$

$$\begin{aligned} \text{Thus } & \int \frac{3}{u^2} du = \int 3u^{-2} du = -3u^{-1} + C \\ & = -\frac{3}{u} + C = \boxed{-\frac{3}{x+a} + C} \end{aligned}$$

$a \neq b$ We need to write $\frac{3}{(x+a)(x+b)} \approx \frac{A}{x+a} + \frac{B}{x+b}$

for some A, B constants.

$$\frac{3}{(x+a)(x+b)} = \frac{A}{x+a} + \frac{B}{x+b} = \frac{A(x+b) + B(x+a)}{(x+a)(x+b)}$$

Example 1 (Online Homework # 7)

Evaluate the indefinite integral: $\int \frac{3}{(x+a)(x+b)} dx$.

<http://www.ms.uky.edu/~ma138>

Lecture 7

Thus the numerators must be the same:

$$\begin{aligned} 3 &= A(x+b) + B(x+a) \\ &= (A+B)x + (Ab+Ba) \end{aligned}$$

$$\therefore A+B=0 \quad \text{and} \quad Ab+Ba=3$$

$$\hookrightarrow A=-B$$

$$\hookrightarrow -Bb+Ba=3$$

$$\therefore \boxed{B = \frac{3}{a-b}} \quad \text{and} \quad \boxed{A = -B = \frac{-3}{a-b}}$$

$$\text{Thus } \frac{3}{(x-a)(x+b)} = \frac{-3/a-b}{x+a} + \frac{3/a-b}{x+b}$$

$$= \frac{3}{a-b} \left[\frac{1}{x+b} - \frac{1}{x+a} \right]$$

$$\therefore \int \frac{3}{(x-a)(x+b)} dx = \frac{3}{a-b} \left[\int \frac{1}{x+b} dx - \int \frac{1}{x+a} dx \right]$$

$$\int \frac{3}{(x+a)(x+b)} dx = \frac{3}{a-b} \left[\ln|x+b| - \ln|x+a| \right] + C$$

$$= \boxed{\frac{3}{a-b} \cdot \ln \left| \frac{x+b}{x+a} \right| + C}$$

$$= \boxed{\frac{3}{b-a} \cdot \ln \left| \frac{x+a}{x+b} \right| + C}$$

Let's try to find A and B such that

$$\begin{aligned} \frac{4x^2 - x - 1}{(x+1)^2(x-3)} &= \frac{A}{(x+1)^2} + \frac{B}{x-3} \\ &= \frac{A(x-3) + B(x+1)^2}{(x+1)^2(x-3)} \\ &= \frac{Ax - 3A + Bx^2 + 2Bx + B}{(x+1)^2(x-3)} \\ &= \frac{Bx^2 + (A+2B)x + (B-3A)}{(x+1)^2(x-3)} \end{aligned}$$

Thus: $\boxed{B=4}$, $\boxed{A+2B=-1}$, $\boxed{B-3A=-1}$

\downarrow $\boxed{A+8=-1} \Rightarrow \boxed{A=-9}$

But: $\boxed{4-3(-9)=-1}$ not

IMPOSSIBLE!!!

Example 2

Consider the rational function

$$f(x) = \frac{4x^2 - x - 1}{(x+1)^2(x-3)}$$

which has a repeated factor at the denominator.

Try to find constants A and B such that

$$\frac{4x^2 - x - 1}{(x+1)^2(x-3)} = \frac{A}{(x+1)^2} + \frac{B}{x-3}.$$

<http://www.ms.uky.edu/~ma138>

Lecture 7

Example 2 (again)

The previous calculation didn't work.

Try now to find constants A , B and C such that

$$\frac{4x^2 - x - 1}{(x+1)^2(x-3)} = \frac{A}{(x+1)} + \frac{B}{(x+1)^2} + \frac{C}{x-3}.$$

Then evaluate the definite integral

$$\int \frac{4x^2 - x - 1}{(x+1)^2(x-3)} dx.$$

<http://www.ms.uky.edu/~ma138>

Lecture 7

$$\frac{4x^2 - x - 1}{(x+1)^2(x-3)} = \frac{A}{(x+1)} + \frac{B}{(x+1)^2} + \frac{C}{x-3}$$

$$= \frac{A(x+1)(x-3) + B(x-3) + C(x+1)^2}{(x+1)^2(x-3)}$$

The denominators are the same:

$$4x^2 - x - 1 = A(x+1)(x-3) + B(x-3) + C(x+1)^2 \quad (*)$$

If we evaluate (*) for $x = -1$ we obtain

$$4(-1)^2 - (-1) - 1 = A \cdot 0 + B(-4) + C \cdot 0$$

$$\therefore -4B = 4 \quad \therefore \boxed{B = -1}$$

If we evaluate (*) at $x = 3$ we obtain

$$4(3)^2 - 3 - 1 = A \cdot 0 + B \cdot 0 + 16C$$

$$\therefore 16C = 32 \quad \therefore \boxed{C = 2}$$

Partial Fraction Decomposition (repeated linear factors)

Case of Repeated Linear Factors

$Q(x)$ is a product of m distinct linear factors to various powers. $Q(x)$ is thus of the form

$$Q(x) = a(x - \alpha_1)^{n_1}(x - \alpha_2)^{n_2} \cdots (x - \alpha_m)^{n_m}$$

where $\alpha_1, \alpha_2, \dots, \alpha_m$ are the m distinct roots of $Q(x)$ and n_1, n_2, \dots, n_m are positive integers such that $n_1 + n_2 + \cdots + n_m = \deg Q(x)$.

The rational function can then be written as

$$\frac{P(x)}{Q(x)} = \frac{1}{a} \left[\sum_{i=1}^m \frac{A_{i,1}}{x - \alpha_i} + \frac{A_{i,2}}{(x - \alpha_i)^2} + \cdots + \frac{A_{i,n_i}}{(x - \alpha_i)^{n_i}} \right].$$

Thus the (Heaviside) cover-up method doesn't produce directly A ! Well substitute in (*) for example $x = 0$ and get:

$$4 \cdot 0^2 - 0 - 1 = A \cdot (1)(-3) + (-1)(0-3) + (2)(0+1)^2$$

$$\text{So: } -1 = -3A + 3 + 2 \quad \leftrightarrow \quad 3A = 6 \quad \therefore \boxed{A = 2}$$

Bottom line:

$$\begin{aligned} \int \frac{4x^2 - x - 1}{(x+1)^2(x-3)} dx &= \int \frac{2}{x+1} dx + \int \frac{-1}{(x+1)^2} dx + \int \frac{2}{x-3} dx \\ &= 2 \ln|x+1| + \frac{1}{x+1} + 2 \ln|x-3| + C \\ &= \ln[(x+1)^2(x-3)^2] + \frac{1}{x+1} + C \end{aligned}$$

using the properties of logarithms

Example 3 (Online Homework # 5)

Evaluate the integral

$$\int \frac{-10x^2}{(x+1)^3} dx.$$

We need to find A, B, and C such that

$$\begin{aligned}\frac{-10x^2}{(x+1)^3} &= \frac{A}{x+1} + \frac{B}{(x+1)^2} + \frac{C}{(x+1)^3} \\ &= \frac{A(x+1)^2 + B(x+1) + C}{(x+1)^3}\end{aligned}$$

Thus: $-10x^2 = A(x+1)^2 + B(x+1) + C$ (*)

Evaluate (*) at $x=-1$. we get "C" !

$$-10(-1)^2 = A \cdot 0 + B \cdot 0 + C \quad \therefore C = -10$$

If we plug-in in (*) $x=0$ and $x=1$ for example we get :

$$x=0 : 0 = A(1)^2 + B(1) - 10$$

$$x=1 : -10 = A(2)^2 + B(2) - 10$$

$$\text{So: } \begin{cases} A+B=10 \\ 4A+2B=0 \end{cases} \Rightarrow \begin{cases} B=-2A \\ A+(2A)=10 \end{cases}$$

$$\therefore A = -10 \quad \text{and} \quad B = 20$$

$$\text{Thus: } \frac{-10x^2}{(x+1)^3} = \frac{-10}{x+1} + \frac{20}{(x+1)^2} + \frac{-10}{(x+1)^3}$$

$$\begin{aligned}\text{and } \int \frac{-10x^2}{(x+1)^3} dx &= -10 \int \frac{1}{x+1} dx + 20 \int \frac{1}{(x+1)^2} dx - 10 \int \frac{1}{(x+1)^3} dx \\ &= -10 \ln|x+1| + 20 \left[-\frac{1}{(x+1)} \right] - 10 \left[-\frac{1}{2} \frac{1}{(x+1)^2} \right] + C\end{aligned}$$

$$\boxed{\begin{aligned}&= -10 \ln|x+1| + \frac{-20}{x+1} + \frac{5}{(x+1)^2} + C\end{aligned}}$$

$$\boxed{\begin{aligned}&= -10 \ln|x+1| + \frac{-20x-15}{(x+1)^2} + C\end{aligned}}$$

$$\int \frac{-10x^2}{(x+1)^3} dx \quad \text{you can also } u=x+1 \quad du=dx$$

$$\Rightarrow x=u-1$$

$$= \int \frac{-10(u-1)^2}{u^3} du = \int \frac{-10(u^2-2u+1)}{u^3} du$$

$$= \int \left[10 \frac{u^2}{u^3} + 20 \frac{u}{u^3} - 10 \frac{1}{u^3} \right] du =$$

$$= \int \left[-10 \frac{1}{u} + 20 \frac{1}{u^2} - 10 \frac{1}{u^3} \right] du =$$

$$= -10 \ln|u| - 20 \frac{1}{u} + 5 \frac{1}{u^2} + C$$

$$\boxed{\begin{aligned}&= -10 \ln|x+1| - 20 \frac{1}{x+1} + 5 \frac{1}{(x+1)^2} + C\end{aligned}}$$

Example 4

Evaluate the integral

$$\int \frac{1}{x^2(x-1)^2} dx.$$

We want to write $\frac{1}{x^2(x-1)^2}$ as the following sum of fractions

$$\frac{1}{x^2(x-1)^2} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x-1} + \frac{D}{(x-1)^2}$$

for some constants A, B, C , and D . Give common denominator -

$$\begin{aligned} &= \frac{Ax(x-1)^2 + B(x-1)^2 + Cx^2(x-1) + Dx^2}{x^2(x-1)^2} \\ &= \frac{A(x^3 - 2x^2 + x) + B(x^2 - 2x + 1) + C(x^3 - x^2) + Dx^2}{x^2(x-1)^2} \\ &= \frac{x^3[A+C] + x^2[-2A+B-C+D] + x[A-2B]}{x^2(x-1)^2} \end{aligned}$$

Equate the coefficients of the denominators :

$$A+C=0 ; -2A+B-C+D=0 ; A-2B=0 ; B=1$$

Example 5 (Online Homework #11)

If $f(x)$ is a quadratic function such that $f(0)=1$ and $\int \frac{f(x)}{x^2(x+1)^3} dx$ is a rational function, find the value of $f'(0)$.

Thus $B=1$; $A=2B=2$; $C=-A=-2$.

Finally from $-2A+B-C+D=0$ we get

$$-4+1-(-2)+D=0 \quad \text{so} \quad D=1$$

Hence :

$$\frac{1}{x^2(x-1)^2} = \frac{2}{x} + \frac{1}{x^2} + \frac{-2}{x-1} + \frac{1}{(x-1)^2} \quad \text{using this, we get}$$

$$\begin{aligned} \int \frac{1}{x^2(x-1)^2} dx &= \left[2\ln|x| - \frac{1}{x} - 2\ln|x-1| - \frac{1}{(x-1)} + C \right] \\ &= \ln \left[\frac{x^2}{(x-1)^2} \right] + \frac{1-2x}{x(x-1)} + C \end{aligned}$$

after simplifying and using the properties of logarithms

$f(x)$ is a quadratic function; so $f(x)=ax^2+bx+c$

We know that $f(0)=1$ so $f(0)=a \cdot 0^2 + b \cdot 0 + c = \boxed{c=1}$

Thus $f(x)=ax^2+bx+1$. We are asked to compute

$f'(0)$. Now $f'(x)=2ax+b$. Hence

we are asked to compute $f'(0)=2a \cdot 0 + b = \boxed{b}$

We are asking that

$$\int \frac{f(x)}{x^2(x+1)^3} dx = \int \frac{ax^2+bx+1}{x^2(x+1)^3} dx \text{ is a rational function.}$$

Think about the partial fraction decomposition

$$\frac{ax^2+bx+1}{x^2(x+1)^3} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{(x+1)} + \frac{D}{(x+1)^2} + \frac{E}{(x+1)^3}$$

If A and C were to be $\neq 0$ those 2 fractions would give rise to logarithmic terms. Thus we need a partial fraction decomposition of the form:

$$\begin{aligned} \frac{ax^2+bx+c}{x^2(x+1)^3} &= \frac{B}{x^2} + \frac{D}{(x+1)^2} + \frac{E}{(x+1)^3} \\ &= \frac{B(x+1)^3 + D x^2(x+1) + E x^2}{x^2(x+1)^3} \\ &= \frac{B(x^3+3x^2+3x+1) + D(x^3+x^2) + E x^2}{x^2(x+1)^3} \\ &= \frac{x^3[B+D] + x^2[3B+D+E] + 3Bx + B}{x^2(x+1)^3} \end{aligned}$$

Thus the coefficients of the numerators must be the same:

$$B+D=0 \quad 3B+D+E=a \quad 3B=b \quad B=1$$

$$\text{Thus } B=1 \quad \text{so } b=3B=3 \quad \boxed{b=3B=3} \quad \leftarrow \quad \leftarrow$$

Note that $D=-B=-1$ so that

$$E=a-3B-D=a-3-(-1)=a-2$$

Hence

$$\frac{ax^2+bx+c}{x^2(x+1)^3} = \frac{1}{x^2} - \frac{1}{(x+1)^2} + \frac{a-2}{(x+1)^3}$$

Partial Fraction Decomposition (irreducible quadratic factors)

Irreducible quadratic factors in the denominator of a proper rational functions are dealt with in the partial-fraction decomposition as follows:

Case of Irreducible Quadratic Factors

If the irreducible quadratic factor $ax^2 + bx + c$ is contained n times in the factorization of the denominator of a proper rational function, then the partial-fraction decomposition contains terms of the form

$$\frac{B_1x+C_1}{ax^2+bx+c} + \frac{B_2x+C_2}{(ax^2+bx+c)^2} + \cdots + \frac{B_nx+C_n}{(ax^2+bx+c)^n}.$$

Example 6 (Example 6, Section 7.3, page 348)

Write the partial fraction decomposition of

$$f(x) = \frac{2x^3 - x^2 + 2x - 2}{(x^2 + 1)(x^2 + 2)}.$$

$$\frac{2x^3 - x^2 + 2x - 2}{(x^2+1)(x^2+2)} = \frac{Ax+B}{x^2+1} + \frac{Cx+D}{x^2+2}$$

for some A, B, C , and D .

$$= \frac{(Ax+B)(x^2+2) + (Cx+D)(x^2+1)}{(x^2+1)(x^2+2)}$$

$$= \frac{Ax^3 + 2Ax + Bx^2 + 2B + Cx^3 + Cx + Dx^2 + D}{(x^2+1)(x^2+2)}$$

$$= \frac{x^3[A+C] + x^2[B+D] + x[2A+C] + 2B+D}{(x^2+1)(x^2+2)}$$

so that, if we equate the coefficients of the numerators

$$\boxed{A+C=2}$$

$$\boxed{B+D=-1}$$

$$\boxed{2A+C=2}$$

$$\boxed{2B+D=-2}$$

So we need to solve

$$\begin{cases} A+C=2 \\ 2A+C=2 \end{cases}$$

$$\begin{cases} B+D=-1 \\ 2B+D=-2 \end{cases}$$

These 2 systems give

$$A=0 \quad C=2$$

$$B=-1 \quad D=0$$

Hence

$$\left| \frac{2x^3 - x^2 + 2x - 2}{(x^2+1)(x^2+2)} = \frac{-1}{x^2+1} + \frac{2x}{x^2+2} \right|$$

$$\left[\int \frac{2x^3 - x^2 + 2x - 2}{(x^2+1)(x^2+2)} = \ln(x^2+2) - \tan^{-1}x + C \right]$$