

MA 138 – Calculus 2 with Life Science Applications

Improper Integrals

(Section 7.4)

Alberto Corso

alberto.corso@uky.edu

Department of Mathematics
University of Kentucky

Wednesday, February 1, 2017

<http://www.ms.uky.edu/~ma138>

Lecture 9

Type 2: Unbounded Integrand

What if the integrand becomes infinite at one or both endpoints of the interval of integration?

- If f is continuous on $(a, b]$ and $\lim_{x \rightarrow a^+} f(x) = \pm\infty$, we define

$$\int_a^b f(x) dx := \lim_{c \rightarrow a^+} \int_c^b f(x) dx$$

provided that this limit exists.

- If f is continuous on $[a, b)$ and $\lim_{x \rightarrow b^-} f(x) = \pm\infty$, we define

$$\int_a^b f(x) dx := \lim_{c \rightarrow b^-} \int_a^c f(x) dx$$

provided that this limit exists.

If the limit does not exist, we say that the integral diverges.

<http://www.ms.uky.edu/~ma138>

Lecture 9

Improper Integrals

We discuss definite integrals of two types with the following characteristics:

- one or both limits of integration are infinite; that is, the integration interval is unbounded. For example

$$\int_1^\infty e^{-x} dx \quad \text{or} \quad \int_{-\infty}^\infty \frac{1}{1+x^2} dx;$$

(These integrals are very important in Probability and Statistics!)

- the integrand becomes infinite at one or more points of the interval of integration. For example

$$\int_{-1}^1 \frac{1}{x^2} dx \quad \text{or} \quad \int_0^1 \frac{1}{2\sqrt{x}} dx.$$

We call such integrals **improper integrals**.

<http://www.ms.uky.edu/~ma138>

Lecture 9

Example 1 (Problem #12, Section 7.4, page 362)

Determine whether the improper integral

$$\int_1^e \frac{1}{x\sqrt{\ln x}} dx$$

is convergent. If the integral is convergent, compute its value.

<http://www.ms.uky.edu/~ma138>

Lecture 9

$$\begin{aligned}
 &= \int_1^e \frac{1}{x \sqrt{\ln x}} dx \quad \text{use the substitution} \\
 &\qquad u = \ln x \\
 &\qquad du = \frac{1}{x} dx \\
 &= \int_0^1 \frac{1}{\sqrt{u}} du \quad \text{notice that the integrand is} \\
 &\qquad \text{not defined when } \underline{u=0} !! \\
 &= \lim_{a \rightarrow 0^+} \int_a^1 \frac{1}{\sqrt{u}} du = \lim_{a \rightarrow 0^+} \left[2\sqrt{u} \right]_a^1 = \\
 &= \lim_{a \rightarrow 0^+} \left[2\sqrt{1} - 2\sqrt{a} \right] = 2 - 2 \cdot \underbrace{\lim_{a \rightarrow 0^+} \sqrt{a}}_{=0} \\
 &= \boxed{2}
 \end{aligned}$$

$$\begin{aligned}
 &= \int_1^e \frac{1}{x \cdot \ln x} dx \quad \text{use the substitution } u = \ln x \\
 &\qquad du = \frac{1}{x} dx \quad . \text{ Thus} \\
 &= \int_0^1 \frac{1}{u} du \quad \text{notice that the integrand } \frac{1}{u} \\
 &\qquad \text{is not defined when } u=0. \\
 &= \lim_{a \rightarrow 0^+} \int_a^1 \frac{1}{u} du = \lim_{a \rightarrow 0^+} \left[\ln|u| \right]_a^1 = \\
 &= \lim_{a \rightarrow 0^+} \left[\underbrace{\ln(1)}_0 - \ln(a) \right] = \\
 &= - \lim_{a \rightarrow 0^+} \ln(a) = -(-\infty) = \boxed{+\infty} \\
 &\qquad \text{diverges}
 \end{aligned}$$

Example 2 (Problem #26, Section 7.4, page 362)

Determine whether the improper integral

$$\int_1^e \frac{1}{x \ln x} dx$$

is convergent. If the integral is convergent, compute its value.

<http://www.ms.uky.edu/~ma138>

Lecture 9

Example 3 (Online Homework #7)

Determine whether the improper integral

$$\int_0^9 \frac{4}{(x-6)^2} dx$$


is convergent. If the integral is convergent, compute its value.

<http://www.ms.uky.edu/~ma138>

Lecture 9

$$= \int_0^9 \frac{4}{(x-6)^2} dx$$

use the substitution $u = x-6$
so that $du = dx$



$$= \int_{-6}^3 \frac{4}{u^2} du$$

$$= \int_{-6}^0 \frac{4}{u^2} du + \int_0^3 \frac{4}{u^2} du \quad \text{the integrand } \frac{1}{u^2} \text{ is not defined at } u=0$$

Note that both integrals are " $+\infty$ ". So the whole integral does not exist (or diverges).

e.g. $\int_0^3 \frac{4}{u^2} du = \lim_{a \rightarrow 0^+} \int_a^3 \frac{4}{u^2} du = \lim_{a \rightarrow 0^+} \left[-\frac{4}{u} \right]_a^3 =$

$$= \lim_{a \rightarrow 0^+} \left(-\frac{4}{3} - \left(-\frac{4}{a} \right) \right)$$

$$= \lim_{a \rightarrow 0^+} \left[\frac{4}{a} - \frac{4}{3} \right] = +\infty - \frac{4}{3} = \underline{\underline{+\infty}}$$

Similarly:

$$\int_{-6}^0 \frac{4}{u^2} du = \lim_{b \rightarrow 0^-} \int_{-6}^b \frac{4}{u^2} du = \lim_{b \rightarrow 0^-} \left[-\frac{4}{u} \right]_{-6}^b$$

$$= \lim_{b \rightarrow 0^-} \left[-\frac{4}{b} - \left(-\frac{4}{(-6)} \right) \right] = \lim_{b \rightarrow 0^-} \left[-\frac{4}{b} - \frac{2}{3} \right]$$

$$= - \left[\lim_{b \rightarrow 0^-} \frac{4}{b} \right] - \frac{2}{3} = -(-\infty) - \frac{2}{3} = \underline{\underline{+\infty}}$$

Example 4 (Problem #34, Section 7.4, page 363)

Let p be a positive real number. Show that

$$\int_0^1 \frac{1}{x^p} dx = \begin{cases} \frac{1}{1-p} & \text{for } 0 < p < 1 \\ \infty & \text{for } p \geq 1. \end{cases}$$

E.g.: $\int_0^1 \frac{1}{x} dx$ and $\int_0^1 \frac{1}{x^2} dx$ both diverge (as $p = 1, 2$, respectively).

E.g.: $\int_0^1 \frac{1}{\sqrt{x}} dx = 2$ and $\int_0^1 \frac{1}{\sqrt[3]{x}} dx = \frac{3}{2}$ (as $p = 1/2, 1/3$, respectively).

$$\int_0^1 \frac{1}{x^p} dx = \lim_{a \rightarrow 0^+} \int_a^1 \frac{1}{x^p} dx = \lim_{a \rightarrow 0^+} \int_a^1 x^{-p} dx$$

for $p \neq 1$

$$= \lim_{a \rightarrow 0^+} \left[\frac{1}{1-p} x^{1-p} \right]_a^1 = \lim_{a \rightarrow 0^+} \left[\frac{1}{1-p} - \frac{1}{1-p} a^{1-p} \right]$$

$$= \frac{1}{1-p} - \frac{1}{1-p} \underbrace{\lim_{a \rightarrow 0^+} a^{1-p}}_{\text{converges}}$$

diverges

0 for
 $0 < p < 1$

for $p = 1$

$$\lim_{a \rightarrow 0^+} \int_a^1 \frac{1}{x} dx = \lim_{a \rightarrow 0^+} \left[\ln|x| \right]_a^1 = \lim_{a \rightarrow 0^+} [0 - \ln(a)]$$

$$= 0 - (-\infty) = \underline{\underline{+\infty}} \quad \text{diverges}$$

Example 5 (Problem #15, Section 7.4, page 362)


Determine whether the improper integral

$$\int_{-1}^1 \ln|x| dx.$$

is convergent. If the integral is convergent, compute its value.

<http://www.ms.uky.edu/~ma138>

Lecture 9



Notice that

$$\int_{-1}^1 \ln|x| dx = \int_{-1}^0 \ln|x| dx + \int_0^1 \ln|x| dx$$

If both integrals exist, by symmetry then

$$\int_{-1}^1 \ln|x| dx = 2 \int_0^1 \ln x dx = \boxed{2 \lim_{a \rightarrow 0^+} \int_a^1 \ln x dx}$$

Recall that (integration by parts):

$$\begin{aligned} \int \ln x dx &= x \cdot \ln x - \int x \cdot \frac{1}{x} dx = x \ln x - \int dx \\ &= x \ln x - x + C = x(\ln x - 1) + C \end{aligned}$$

Thus:

$$\begin{aligned} \int_{-1}^1 \ln|x| dx &= 2 \lim_{a \rightarrow 0^+} \int_a^1 \ln x dx \\ &= 2 \lim_{a \rightarrow 0^+} \left[x(\ln x - 1) \right]_a^1 \\ &= 2 \lim_{a \rightarrow 0^+} \left\{ [1 \cdot (\ln(1) - 1)] - [a(\ln a - 1)] \right\} \\ &= 2 \lim_{a \rightarrow 0^+} \left\{ -1 - \underbrace{a \cdot \ln(a)}_{\substack{\xrightarrow{0} \\ 0}} + \underbrace{a}_{\substack{\xrightarrow{0} \\ 0}} \right\} \\ &= \boxed{-2} \end{aligned}$$

Note: $\lim_{a \rightarrow 0^+} a \cdot \ln(a) = 0 \cdot (-\infty) = \lim_{a \rightarrow 0^+} \frac{\ln(a)}{\frac{1}{a}} \stackrel{(l'Hopital)}{=} \lim_{a \rightarrow 0^+} \frac{\frac{1}{a}}{-\frac{1}{a^2}} = \lim_{a \rightarrow 0^+} (-a) = 0$