

# MA 138 – Calculus 2 with Life Science Applications

## Integration by Parts

(Section 7.2)

**Alberto Corso**

⟨alberto.corso@uky.edu⟩

Department of Mathematics  
University of Kentucky

Monday, January 23, 2017

## Section 7.2: Integration by Parts

We saw that **integration by parts is the product rule in integral form.**

We also recall the following **general formula:**

### Rule for Integration by Parts

If  $f(x)$  and  $g(x)$  are differentiable functions, then

$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx;$$

$$\int_a^b f(x)g'(x) dx = f(x)g(x) \Big|_a^b - \int_a^b f'(x)g(x) dx.$$

**Example 1**

A particle that moves along a straight line has velocity

$$v(t) = t^2 e^{-2t}$$

meters per second after  $t$  seconds.

How many meters will it travel during the first  $t$  seconds?

**Example 2** (Online Homework # 10)

Suppose that  $f(1) = 4$ ,  $f(4) = 6$ ,  $f'(1) = -5$ ,  $f'(4) = -5$ ,  
and  $f''$  is continuous. Find the value of

$$\int_1^4 x f''(x) dx.$$

**Example 3** (Problem # 8, Section 7.2, page 342)

Evaluate the indefinite integral:  $\int 3xe^{-x/2} dx$ .

**Example 4** (Online Homework # 7)

Find the integral:  $\int_0^1 x^2 \sqrt[4]{e^x} dx.$

**Example 5** (Problem # 35, Section 7.2, page 343)

Evaluate the indefinite integral:  $\int \frac{1}{x} \ln x \, dx.$

**Example 6** (Problem # 48, Section 7.2, page 343)

Evaluate the definite integral:  $\int_0^1 x^3 \ln(x^2 + 1) dx.$



## Useful aside: Trigonometric addition formulas

- We also used the double angle formulas

$$\begin{aligned} \cos(2\alpha) &= \cos^2 \alpha - \sin^2 \alpha & \sin(2\alpha) &= 2 \sin \alpha \cos \alpha \\ &= 2 \cos^2 \alpha - 1 & \text{and} & \\ &= 1 - 2 \sin^2 \alpha \end{aligned}$$

to compute  $\int \cos^2 x \, dx$  and  $\int \sin x \cos x \, dx$ .

- Is there a 'simple' way to remember formulas of this kind?
- **Euler's formula** establishes the fundamental relationship between the trigonometric functions and the complex exponential function. It states that, for any real number  $x$ ,

$$e^{ix} = \cos x + i \sin x,$$

where  $i$  is the imaginary unit ( $i^2 = -1$ ).

- For any  $\alpha$  and  $\beta$ , using Euler's formula, we have

$$\begin{aligned}
 \cos(\alpha + \beta) + i \sin(\alpha + \beta) &= e^{i(\alpha+\beta)} \\
 &= e^{i\alpha} \cdot e^{i\beta} \\
 &= (\cos \alpha + i \sin \alpha) \cdot (\cos \beta + i \sin \beta) \\
 &= (\cos \alpha \cos \beta + i^2 \sin \alpha \sin \beta) \\
 &\quad + i(\sin \alpha \cos \beta + \cos \alpha \sin \beta).
 \end{aligned}$$

- Thus, by comparing the terms, we obtain

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta.$$

- Thus, by setting  $\alpha = \beta$ , we obtain

$$\cos(2\alpha) = \cos^2 \alpha - \sin^2 \alpha \quad \text{and} \quad \sin(2\alpha) = 2 \sin \alpha \cos \beta.$$