

MA 138 – Calculus 2 with Life Science Applications  
**Vector Valued Functions**  
(Section 10.4)

**Alberto Corso**  
(alberto.corso@uky.edu)

Department of Mathematics  
University of Kentucky

April 12, 2017

## Vector-valued functions

- So far, we have considered only real-valued functions  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ .
- We now extend our discussion to functions whose the range is a subset of  $\mathbb{R}^m$  — that is,  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- Such functions are vector-valued functions, since they take on values that are represented by vectors:

$$\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad (x_1, x_2, \dots, x_n) \mapsto \begin{bmatrix} f_1(x_1, x_2, \dots, x_n) \\ f_2(x_1, x_2, \dots, x_n) \\ \vdots \\ f_m(x_1, x_2, \dots, x_n) \end{bmatrix}.$$

- Here, each function  $f_i(x_1, \dots, x_n)$  is a real-valued function:

$$f_i : \mathbb{R}^n \rightarrow \mathbb{R} \quad (x_1, x_2, \dots, x_n) \mapsto f_i(x_1, x_2, \dots, x_n).$$

We will encounter vector-valued functions where  $n = m = 2$  in Chapter 11.

### Example

As an example, consider a community consisting of two species.

Let  $u$  and  $v$  denote the respective densities of the species and  $f(u, v)$  and  $g(u, v)$  the per capita growth rates of the species as functions of the densities  $u$  and  $v$ .

We can then write this relationship as a map

$$\mathbf{h} : \mathbb{R}^2 \longrightarrow \mathbb{R}^2 \quad (u, v) \mapsto \begin{bmatrix} f(u, v) \\ g(u, v) \end{bmatrix}.$$

**E.g.**, in the Lotka-Volterra predator-prey model:  $(u, v) \mapsto \begin{bmatrix} \alpha - \beta v \\ \gamma u - \delta \end{bmatrix}$ ,

where  $\alpha, \beta, \gamma$ , and  $\delta$  are constants.

## Review

- We have defined earlier the linearization at a point  $(x_0, y_0)$  of a real-valued function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ ; namely,

$$L_f(x, y) = f(x_0, y_0) + \frac{\partial f(x_0, y_0)}{\partial x}(x - x_0) + \frac{\partial f(x_0, y_0)}{\partial y}(y - y_0).$$

- We can write the above equation in matrix notation as

$$L_f(x, y) = f(x_0, y_0) + \underbrace{\begin{bmatrix} \frac{\partial f(x_0, y_0)}{\partial x} & \frac{\partial f(x_0, y_0)}{\partial y} \end{bmatrix}}_{1 \times 2 \text{ matrix}} \cdot \underbrace{\begin{bmatrix} x - x_0 \\ y - y_0 \end{bmatrix}}_{2 \times 1 \text{ matrix}}.$$

## Our Goal

- Our task is to define the linearization at a point  $(x_0, y_0)$  of vector-valued functions whose domain and range are  $\mathbb{R}^2$ ; that is,

$$\mathbf{h} : \mathbb{R}^2 \longrightarrow \mathbb{R}^2 \quad (x, y) \mapsto \begin{bmatrix} f(x, y) \\ g(x, y) \end{bmatrix}.$$

- To do so, we linearize at the point  $(x_0, y_0)$  each component of  $\mathbf{h}(x, y)$

$$L_f(x, y) = f(x_0, y_0) + \frac{\partial f(x_0, y_0)}{\partial x}(x - x_0) + \frac{\partial f(x_0, y_0)}{\partial y}(y - y_0)$$

$$L_g(x, y) = g(x_0, y_0) + \frac{\partial g(x_0, y_0)}{\partial x}(x - x_0) + \frac{\partial g(x_0, y_0)}{\partial y}(y - y_0).$$

- We define the linearization of  $\mathbf{h}(x, y)$  at the point  $(x_0, y_0)$  to be the vector-valued function  $\mathbf{L}(x, y)$

$$\mathbf{L}(x, y) = \begin{bmatrix} L_f(x, y) \\ L_g(x, y) \end{bmatrix}.$$

## The Jacobi (or Derivative) Matrix

We can rewrite the linearization  $\mathbf{L}(x, y)$  at a point  $(x_0, y_0)$  of the vector-valued functions  $\mathbf{h}(x, y)$  in the following matrix form

$$\begin{aligned}\mathbf{h}(x, y) \approx \mathbf{L}(x, y) &= \begin{bmatrix} L_f(x, y) \\ L_g(x, y) \end{bmatrix} \\ &= \begin{bmatrix} f(x_0, y_0) + \frac{\partial f(x_0, y_0)}{\partial x}(x - x_0) + \frac{\partial f(x_0, y_0)}{\partial y}(y - y_0) \\ g(x_0, y_0) + \frac{\partial g(x_0, y_0)}{\partial x}(x - x_0) + \frac{\partial g(x_0, y_0)}{\partial y}(y - y_0) \end{bmatrix} \\ &= \underbrace{\begin{bmatrix} f(x_0, y_0) \\ g(x_0, y_0) \end{bmatrix}}_{\mathbf{h}(x_0, y_0)} + \underbrace{\begin{bmatrix} \frac{\partial f(x_0, y_0)}{\partial x} & \frac{\partial f(x_0, y_0)}{\partial y} \\ \frac{\partial g(x_0, y_0)}{\partial x} & \frac{\partial g(x_0, y_0)}{\partial y} \end{bmatrix}}_{(D\mathbf{h})(x_0, y_0)} \cdot \begin{bmatrix} (x - x_0) \\ (y - y_0) \end{bmatrix}\end{aligned}$$

$(D\mathbf{h})(x_0, y_0)$  is a  $2 \times 2$  matrix called the **Jacobi matrix** of  $\mathbf{h}$  at  $(x_0, y_0)$ .

## Example 1 (Problem #10, Exam 3, Spring 2012)

Consider the vector valued function  $\mathbf{h} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  given by

$$\mathbf{h}(x, y) = \begin{bmatrix} x^2y - y^3 \\ 2x^3y^2 + y \end{bmatrix}.$$

- (a) Compute the **Jacobi matrix**  $(D\mathbf{h})(x, y)$  and evaluate it at the point  $(1, 2)$ .
- (b) Find the linear approximation of  $\mathbf{h}(x, y)$  at the point  $(1, 2)$ .

**Example 2** (Problem #46, Section 10.4, p. 536)

Find a linear approximation to

$$\mathbf{f}(x, y) = \begin{bmatrix} \sqrt{2x + y} \\ x - y^2 \end{bmatrix}$$

at  $(1, 2)$ . Use your result to find an approximation for  $\mathbf{f}(1.05, 2.05)$ .



### Example 3 (Example # 9, Section 10.4, p. 534)

Consider the function  $\mathbf{f} : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \quad (x, y) \mapsto \begin{bmatrix} u(x, y) \\ v(x, y) \end{bmatrix}$ , with

$$u(x, y) = y e^{-x} \quad \text{and} \quad v(x, y) = \sin x + \cos y.$$

Find the linear approximation to  $\mathbf{f}(x, y)$  at  $(0, 0)$ .

Compare  $\mathbf{f}(0.1, -0.1)$  with its linear approximation.

## The General Case

- Consider the function  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , say  $\mathbf{f}(x_1, \dots, x_n) = \begin{bmatrix} f_1(x_1, \dots, x_n) \\ \vdots \\ f_m(x_1, \dots, x_n) \end{bmatrix}$ ,

where  $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ , are  $m$  real-valued functions of  $n$  variables.

- The Jacobi matrix of  $\mathbf{f}$  is an  $m \times n$  matrix of the form

$$(D\mathbf{f})(x_1, \dots, x_n) = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \vdots & & \vdots \\ \frac{\partial f_m}{\partial x_1} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}$$

- The linearization of  $\mathbf{f}$  about the point  $(x_1^*, \dots, x_n^*)$  is then

$$\mathbf{L}(x_1, \dots, x_n) = \begin{bmatrix} f_1(x_1^*, \dots, x_n^*) \\ \vdots \\ f_m(x_1^*, \dots, x_n^*) \end{bmatrix} + (D\mathbf{f})(x_1^*, \dots, x_n^*) \cdot \begin{bmatrix} x_1 - x_1^* \\ \vdots \\ x_n - x_n^* \end{bmatrix}$$