

MA 138 – Calculus 2 with Life Science Applications
Linear Systems: Theory
(Section 11.1)

Alberto Corso

`<alberto.corso@uky.edu>`

Department of Mathematics
University of Kentucky

Friday, April 14, 2017

Systems of Differential Equations

- Suppose that we are given a set of variables x_1, x_2, \dots, x_n , each depending on an independent variable, say, t , so that

$$x_1 = x_1(t), \quad x_2 = x_2(t), \quad \dots, \quad x_n = x_n(t).$$

- Suppose also that the dynamics of the variables are linked by n differential equations (\equiv DEs) of the first-order; that is,

$$\begin{cases} \frac{dx_1}{dt} = g_1(t, x_1, x_2, \dots, x_n) \\ \vdots \\ \frac{dx_n}{dt} = g_n(t, x_1, x_2, \dots, x_n) \end{cases}$$

- This set of equations is called a **system of differential equations**.
- On the LHS are the derivatives of $x_i(t)$ with respect to t . On the RHS is a function g_i that depends on the variables x_1, x_2, \dots, x_n and on t .

Examples

■ Kermack & McKendrick epidemic disease model (SIR, 1927)

$$\begin{cases} \frac{dS}{dt} = -bSI \\ \frac{dI}{dt} = bSI - aI \\ \frac{dR}{dt} = aI \end{cases}$$

$S = S(t)$ = # of susceptible individuals

$I = I(t)$ = # of infected individuals

$R = R(t)$ = # of removed individuals (\equiv no longer susceptible)

a, b = constant rates

■ Lotka-Volterra predator-prey model (1910/1920):

$$\begin{cases} \frac{dN}{dt} = rN - aPN \\ \frac{dP}{dt} = abPN - dP \end{cases}$$

$N = N(t)$ = prey density

$P = P(t)$ = predator density

r = intrinsic rate of increase of the prey

a = attack rate

b = efficiency rate of predators in turning preys into new offsprings

d = rate of decline of the predators

Direction Field of a System of 2 Autonomous DEs

- Review the notion of the direction field of a DE of the first order $dy/dx = f(x, y)$. We encountered this notion just before Section 8.2 (Handout; February 15, 2017).
- Consider, now a system of two autonomous differential equations

$$\begin{cases} \frac{dx}{dt} = g_1(x, y) \\ \frac{dy}{dt} = g_2(x, y) \end{cases}$$

- Assuming that y is also a function of x and using the chain rule, we can eliminate t and obtain the DE


$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{g_2(x, y)}{g_1(x, y)}$$

of which we can plot the direction field.

Example (Lotka-Volterra)

Consider the system of DEs $\frac{dx}{dt} = x - 4xy$ and $\frac{dy}{dt} = 2xy - 3y$.

The direction field of the differential equation $\frac{dy}{dx} = \frac{(2x - 3)y}{x(1 - 4y)}$ has been produced with the SAGE commands in Chapter 8.



Notice that the trajectories are closed curves. Furthermore, they all seem to revolve around the point $P(3/2, 1/4)$. This is the point where the factors $2x - 3$ and $1 - 4y$ of dy/dt and dx/dt , respectively, are both zero.

Linear Systems of Differential Equations (11.1)

- We first look at the case when the g_i 's are linear functions in the variables x_1, x_2, \dots, x_n — that is,

$$\begin{cases} \frac{dx_1}{dt} = a_{11}(t)x_1 + \dots + a_{1n}(t)x_n + f_1(t) \\ \vdots \\ \frac{dx_n}{dt} = a_{n1}(t)x_1 + \dots + a_{nn}(t)x_n + f_n(t) \end{cases}$$

- We can write the linear system in matrix form as

$$\frac{d}{dt} \begin{bmatrix} x_1(t) \\ \vdots \\ x_n(t) \end{bmatrix} = \begin{bmatrix} a_{11}(t) & \dots & a_{1n}(t) \\ \vdots & & \vdots \\ a_{n1}(t) & \dots & a_{nn}(t) \end{bmatrix} \begin{bmatrix} x_1(t) \\ \vdots \\ x_n(t) \end{bmatrix} + \begin{bmatrix} f_1(t) \\ \vdots \\ f_n(t) \end{bmatrix}$$

and we call it an **inhomogeneous system of linear, first-order differential equations**.

- We can write our inhomogeneous system of linear, first-order differential equations as follows

$$\frac{d\mathbf{x}}{dt} = A(t)\mathbf{x} + \mathbf{f}(t)$$

- We are mainly interested in the case when $\mathbf{f}(t) = \mathbf{0}$, that is,

$$\frac{d\mathbf{x}}{dt} = A(t)\mathbf{x},$$

an **homogeneous** system of linear, first-order differential equations.


- Finally, we will study the case in which $A(t)$ does not depend on t

$$\frac{d\mathbf{x}}{dt} = A\mathbf{x},$$

an **homogeneous system of linear, first-order differential equations with constant coefficients.**

Example 1 (Problem #8, Exam 3, Spring 2013)

- (a) Verify that the functions $x(t) = e^{4t} + 5e^{-t}$ and $y(t) = 4e^{4t} - 5e^{-t}$ (whose graphs are given below) are solutions of the system of DEs



$$\begin{cases} \frac{dx}{dt} = & y \\ \frac{dy}{dt} = 4x + 3y \end{cases}$$

with $x(0) = 6$ and $y(0) = -1$.

- (b) Rewrite the given system of DEs and its solutions in the form

$$\underbrace{\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}}_{\text{system of differential equations}}$$

$$\underbrace{\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix} e^{4t} + 5 \begin{bmatrix} \gamma \\ \delta \end{bmatrix} e^{-t}}_{\text{solutions}}$$

for appropriate choices of the constants $a, b, c, d, \alpha, \beta, \gamma,$ and δ .

Specific Solutions of a Linear System of DEs

- Consider the system $\frac{d\mathbf{x}}{dt} = A\mathbf{x}$.

- We claim that the vector-valued function

$$\mathbf{x}(t) = \begin{bmatrix} v_1 e^{\lambda t} \\ v_2 e^{\lambda t} \end{bmatrix} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} e^{\lambda t}$$

where λ , v_1 and v_2 are constants, is a solution of the given system of DEs, for an appropriate choice of values for λ , v_1 , and v_2 .

- More precisely, $\begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ is an eigenvector of the matrix A corresponding to the eigenvalue λ of A .

The Superposition Principle

Principle

Suppose that

$$\begin{bmatrix} \frac{dx_1}{dt} \\ \frac{dx_2}{dt} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix}.$$

If $\mathbf{y}(t) = \begin{bmatrix} y_1(t) \\ y_2(t) \end{bmatrix}$ and $\mathbf{z}(t) = \begin{bmatrix} z_1(t) \\ z_2(t) \end{bmatrix}$

are solutions of the given system of DEs, THEN

$$\mathbf{x}(t) = c_1\mathbf{y}(t) + c_2\mathbf{z}(t)$$

is also a solution of the given system of DEs for any constants c_1 and c_2 .

The General Solution

Theorem

Let

$$\frac{dx}{dt} = Ax$$

where A is a 2×2 matrix with **two real and distinct eigenvalues** λ_1 and λ_2 with corresponding eigenvectors \mathbf{v}_1 and \mathbf{v}_2 .

THEN

$$x(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t}$$

is the general solution of the given system of DEs.

The constants c_1 and c_2 depend on the initial condition.