

MA 138 – Calculus 2 with Life Science Applications  
**Linear Systems: Theory**  
(Section 11.1)

**Alberto Corso**

⟨alberto.corso@uky.edu⟩

Department of Mathematics  
University of Kentucky

Wednesday, April 19, 2017

# Equilibria and Stability

- In Section 8.2 we already encountered the concepts of equilibria and stability, when we discussed ordinary DEs. Both concepts can be extended to systems of DEs.

- We now restrict ourselves to the case

$$\frac{d\mathbf{x}}{dt} = A\mathbf{x} \quad \text{with} \quad A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix}.$$

- We say that a point  $\hat{\mathbf{x}} = \begin{bmatrix} \hat{x}_1 \\ \hat{x}_2 \end{bmatrix}$  is an **equilibrium point** of our given system of linear DEs whenever  $A\hat{\mathbf{x}} = \mathbf{0}$ .

- It follows from results in Section 9.2 that if  $\det A \neq 0$ , then  $(0, 0)$  is the only equilibrium of our given system of linear DEs. If  $\det A = 0$ , then there will be other equilibria.

- If we start a system of DEs at an equilibrium, it remains there at all later times.
- This does not mean that if the system is in equilibrium and is perturbed by a small amount, it will return to the equilibrium.
- Whether or not a solution will return to an equilibrium after a small perturbation is addressed by the **stability** of the equilibrium.
- In the case when the matrix  $A$  has two real and distinct eigenvalues, the solution of our given system of linear DEs is given by

$$x(t) = c_1 e^{\lambda_1 t} \mathbf{v}_1 + c_2 e^{\lambda_2 t} \mathbf{v}_2,$$

where  $\mathbf{v}_1$  and  $\mathbf{v}_2$  are the eigenvectors corresponding to the eigenvalues  $\lambda_1$  and  $\lambda_2$  of  $A$  and the constants  $c_1$  and  $c_2$  depend on the initial condition.

- Knowing the solution will allow us to study the behavior of the solutions as  $t \rightarrow \infty$  and thus address the question of stability, at least when the eigenvalues are distinct.

# Classification of Equilibria

**Case 1:** A has **two distinct real nonzero eigenvalues**  $\lambda_1$  and  $\lambda_2$

- 1. Both eigenvalues are negative:** The equilibrium  $(0, 0)$  is **globally stable**, since the solution will approach the equilibrium  $(0, 0)$  regardless of the starting point. We call  $(0, 0)$  a **sink** or a **stable node**.
- 2. The eigenvalues have opposite signs:** Unless we start in the direction of the eigenvector associated with the negative eigenvalue, the solution will not converge to the equilibrium  $(0, 0)$ . We say that the equilibrium  $(0, 0)$  is **unstable** and call  $(0, 0)$  a **saddle point**.
- 3. Both eigenvalues are positive:** The solution will not converge to  $(0, 0)$  unless we start at  $(0, 0)$ . We say that the equilibrium  $(0, 0)$  is **unstable**, and we call  $(0, 0)$  a **source** or an **unstable node**.

**Case 2:** A has **two complex conjugate eigenvalues**  $\lambda_{1,2} = \alpha \pm i\beta$


1.  $\alpha < 0$ : Starting from any point other than  $(0, 0)$ , solutions spiral into the equilibrium  $(0, 0)$ . For this reason, the equilibrium  $(0, 0)$  is called a **stable spiral**. When we plot solutions as functions of time, they show oscillations. The amplitude of the oscillations decreases over time. We therefore call the oscillations **damped**.
2.  $\alpha > 0$ : Starting from any point other than  $(0, 0)$ , the solutions spiral out from the equilibrium  $(0, 0)$ . For this reason, we call the equilibrium  $(0, 0)$  an **unstable spiral**. When we plot solutions as functions of time, we see that the solutions show oscillations as before, but this time their amplitudes are increasing.
3.  $\alpha = 0$ : Solutions spiral around the equilibrium  $(0, 0)$ , but neither approach nor move away from the equilibrium (since the amplitude of the solutions does not change). The equilibrium  $(0, 0)$  is called a **neutral spiral** or a **center**. The solutions form closed curves.

## Example 8

$(0, 0)$  globally stable equilibrium; sink or a stable node

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$


$$\lambda_1 = -1 \quad \lambda_2 = -4$$



**Example 9**  $(0, 0)$  unstable equilibrium; saddle point

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 & -2 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$


$$\lambda_1 = 1 \quad \lambda_2 = -2$$



## Example 10 $(0, 0)$ unstable node; source

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$\lambda_1 = 1 \quad \lambda_2 = 4$$







# Example 11 $(0, 0)$ stable spiral; negative real part

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -1 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$



$$\lambda_{1,2} = -\frac{1}{2} \pm i\frac{\sqrt{3}}{2}$$



## Example 12 $(0, 0)$ unstable spiral; positive real part

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$



$$\lambda_{1,2} = \frac{1}{2} \pm i \frac{\sqrt{3}}{2}$$



# Example 13 $(0, 0)$ neutral spiral or center; no real part

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ -5 & -3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$\lambda_{1,2} = \pm i$$



## Summary of Stability at $(0, 0)$

- The relationships between the characteristic polynomial, eigenvalues, trace, and determinant of a  $2 \times 2$  matrix  $A$  are given by

$$\det(A - \lambda I_2) = \lambda^2 - \text{trace}(A)\lambda + \det(A) = 0$$

$$\text{trace}(A) = \text{Re}(\lambda_1) + \text{Re}(\lambda_2) \qquad \det(A) = \lambda_1 \lambda_2.$$

- Also, the discriminant of the quadratic equation  $\det(A - \lambda I_2) = 0$  is

$$\Delta = [\text{trace}(A)]^2 - 4 \det(A)$$

Thus the condition  $\Delta = 0$  ( $\equiv$  repeated eigenvalue) describes the parabola  $\det(A) = 1/4[\text{trace}(A)]^2$  in the trace-det plane.

- **Theorem:** The origin  $(0, 0)$  of a system of two linear, homogeneous DEs with constant coefficients is a stable equilibrium  $\Leftrightarrow$  the real parts of both eigenvalues are negative  $\Leftrightarrow \det(A) > 0$  and  $\text{trace}(A) < 0$ .

The stability properties of the equilibrium at the origin can be summarized graphically in terms of the determinant and the trace of the matrix  $A$  in the trace-det plane:

