

MA 138 – Calculus 2 with Life Science Applications

The Substitution Rule

(Section 7.1)

Alberto Corso

(alberto.corso@uky.edu)

Department of Mathematics
University of Kentucky

Wednesday, January 17, 2017

<http://www.ms.uky.edu/~ma138>

Lecture 3

The Substitution Rule

Reversing these steps and integrating along the way, we get

$$\int 3(6x^2 + 3)^2(12x) dx = \int 3u^2 du = u^3 + C = (6x^2 + 3)^3 + C.$$

In the first step, we substituted u for $6x^2 + 3$ and used $du = 12x dx$.

This substitution simplified the integrand.

At the end, we substitute back $6x^2 + 3$ for u to get the final answer in terms of x .We summarize this discussion, by stating the following **general principle**:

Substitution Rule for Indefinite Integrals

If $u = g(x)$, then
$$\int f(\underbrace{g(x)}_u) \underbrace{g'(x) dx}_{du} = \int f(u) du.$$

<http://www.ms.uky.edu/~ma138>

Lecture 3

Section 7.1: The Substitution Rule

The substitution rule is the chain rule in integral form.

We therefore begin by recalling the chain rule.

Suppose that we wish to differentiate

$$f(x) = (6x^2 + 3)^3.$$

This is clearly a situation in which we need to use the chain rule.

We set $u = 6x^2 + 3$ so that $f(u) = u^3$.

The chain rule, using Leibniz notation, tells us that

$$\frac{df}{dx} = \frac{df}{du} \frac{du}{dx} = 3u^2 \cdot (6 \cdot 2x) = 3(6x^2 + 3)^2(12x).$$

<http://www.ms.uky.edu/~ma138>

Lecture 3

The Substitution Rule

Example 1

Evaluate the indefinite integral $\int \cos x \sin x dx$

- by using the substitution $u = \cos x$;
- by using the substitution $u = \sin x$;
- by using the trigonometric identity $\sin(2x) = 2 \sin x \cos x$.

Compare your answers.

<http://www.ms.uky.edu/~ma138>

Lecture 3

Example 2

Evaluate the indefinite integral $\int (2x + 1)e^{x^2+x} dx$.

Substitution Rule for Definite Integrals

The first method (1) is the more common one, and we summarize the procedure as follows:

Substitution Rule for Definite Integrals

If $u = g(x)$, then
$$\int_a^b \underbrace{f(g(x))}_u \underbrace{g'(x) dx}_{du} = \int_{g(a)}^{g(b)} f(u) du.$$

Substitution Rule for Definite Integrals

Part II of the FTC says that when we evaluate a definite integral, we must find an antiderivative of the integrand and then evaluate the antiderivative at the limits of integration.

When we use the substitution $u = g(x)$ to find an antiderivative of an integrand, the antiderivative will be given in terms of u at first.

To complete the calculation, we can proceed in either of two ways:

- (1) we can leave the antiderivative in terms of u and change the limits of integration according to $u = g(x)$;
- (2) we can substitute $g(x)$ for u in the antiderivative and then evaluate the antiderivative at the limits of integration in terms of x .

Example 3 (Online Homework # 6)

Evaluate the definite integral $\int_1^{e^5} \frac{dx}{x(1 + \ln x)}$.

Example 4 (Online Homework # 8)

Consider the indefinite integral $\int \frac{3}{3+e^x} dx$.

- The most appropriate substitution to simplify this integral is $u = f(x)$ where $f(x) = \underline{\hspace{2cm}}$.
We then have $dx = g(u)du$ where $g(u) = \underline{\hspace{2cm}}$.
(**Hint:** you need to back substitute for x in terms of u for this part.)
- After substituting into the original integral we obtain $\int h(u) du$ where $h(u) = \underline{\hspace{2cm}}$.
- To evaluate this integral rewrite the numerator as $3 = u - (u - 3)$.
Simplify, then integrate, thus obtaining $\int h(u) du = H(u)$ where $H(u) = \underline{\hspace{2cm}} + C$.

Example 5 (Online Homework # 9)

Consider the definite integral $\int_0^1 x^2 \sqrt{5x+6} dx$.

- Then the most appropriate substitution to simplify this integral is $u = \underline{\hspace{2cm}}$. Then $dx = f(x)du$ where $f(x) = \underline{\hspace{2cm}}$.
- After making the substitution and simplifying we obtain the integral $\int_a^b g(u) du$ where $g(u) = \underline{\hspace{2cm}}$, $a = \underline{\hspace{1cm}}$ and $b = \underline{\hspace{1cm}}$.
- This definite integral has value = $\underline{\hspace{2cm}}$.

Example 4, cont.ed (Online Homework # 8)

- After substituting back for u we obtain our final answer $\int \frac{3}{3+e^x} dx = \underline{\hspace{2cm}} + C$.

Example 6 (similar to Example 5)

Consider the definite integral $\int_1^2 x^5 \sqrt{x^3+2} dx$.

- Then the most appropriate substitution to simplify this integral is $u = \underline{\hspace{2cm}}$. Then $dx = f(x)du$ where $f(x) = \underline{\hspace{2cm}}$.
- After making the substitution and simplifying we obtain the integral $\int_a^b g(u) du$ where $g(u) = \underline{\hspace{2cm}}$, $a = \underline{\hspace{1cm}}$ and $b = \underline{\hspace{1cm}}$.
- This definite integral has value = $\underline{\hspace{2cm}}$.

Example 7 (Online Homework # 11)

Consider the indefinite integral $\int \frac{1}{3x + 7\sqrt{x}} dx$.

- Then the most appropriate substitution to simplify this integral is $u = \underline{\hspace{2cm}}$. Then $dx = f(x)du$ where $f(x) = \underline{\hspace{2cm}}$.
- After making the substitution and simplifying we obtain the integral

$$\int g(u) du$$

where $g(u) = \underline{\hspace{2cm}}$.

- This last integral is: $= \underline{\hspace{2cm}} + C$.
(Leave out constant of integration from your answer.)
- After substituting back for u we obtain the following final form of the answer: $= \underline{\hspace{2cm}} + C$.