



## Multiple Choice Questions

1. If  $\mathbf{v} = \langle 1, 0, -1 \rangle$ ,  $\mathbf{w} = \langle 1, 2, 3 \rangle$ , and  $\mathbf{p} = \langle 0, 2, 1 \rangle$ , then  $(\mathbf{v} \times \mathbf{w}) - 3\mathbf{p}$  is
  - A.  $\langle 0, 0, 0 \rangle$
  - B.  $\langle 2, -6, 1 \rangle$
  - C.  $\langle 2, -4, 2 \rangle$
  - D.  $\langle 2, -2, -1 \rangle$
  - E.  $\langle 2, -10, -1 \rangle$
  
2. Find the equation of the line through  $(2, 1, 0)$  and perpendicular to the vectors  $\mathbf{i} + \mathbf{j}$  and  $\mathbf{j} + \mathbf{k}$ .
  - A.  $\mathbf{r}(t) = \langle 2 - t, 1 + 2t, t \rangle$
  - B.  $\mathbf{r}(t) = \langle 2 + t, 1 - t, t \rangle$
  - C.  $\mathbf{r}(t) = \langle 2 + t, 1 + t, 0 \rangle$
  - D.  $\mathbf{r}(t) = \langle 2, 1 + t, 1 + 2t \rangle$
  - E.  $\mathbf{r}(t) = \langle 2 + t, 1, -t \rangle$
  
3. The tangent line to the space curve  $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$  at  $t = 1$  meets the  $xy$  plane at the point:
  - A.  $(1, 2, 3)$
  - B.  $(2/3, -1/3, 0)$
  - C.  $(-2/3, 1/3, 0)$
  - D.  $(0, 0, 0)$
  - E.  $(2/3, 1/3, 0)$

4. Let  $\mathbf{F} = \langle xy^2, yz, zx^2 \rangle$ . Then  $\text{curl}(\mathbf{F})$  is equal to:
- A.  $\langle x^2 - 2zx, -2xy + y, z^2 - z \rangle$
  - B.  $-y - x^2 - z^2$
  - C.  $\langle -y, -2xz, -2xy \rangle$
  - D.  $x^2 - 2zx - 2xy + y + z^2 - z$
  - E.  $\langle 2xy, z, 2zx \rangle$
5. The surface  $xyz + y^2 + 4z = 6$  has a normal line  $L$  at  $P = (1, 1, 1)$ . Then  $L$  meets the  $xy$  plane at point  $Q$  which is:
- A.  $(4/5, 2/5, 0)$
  - B.  $(4, 2, 0)$
  - C.  $(-4, 2, 0)$
  - D.  $(-4/5, 2/5, 0)$
  - E.  $(4, -2, 0)$
6. The integral  $\int_0^1 \int_0^x \int_0^y (6xy + 4yz) dz dy dx$  is equal to:
- A.  $19/30$
  - B.  $7/30$
  - C.  $1$
  - D.  $1/2$
  - E. None of the above

7. Let  $x(u, v) = u^2 + uv$  and  $y(u, v) = uv^2$ . Then the Jacobian determinant

$$J = \left| \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix} \right|$$

is:

- A.  $u^2v + 4uv^2$
  - B.  $2u^2v + 2uv^2$
  - C.  $4u^2v + uv^2$
  - D.  $4u^2v$
  - E.  $4uv^2$
8. Find  $\int_C xy^4 ds$  if  $C$  is the right half of the circle  $x^2 + y^2 = 4$ .
- A.  $128/5$
  - B.  $64/5$
  - C.  $32/5$
  - D.  $64\pi/5$
  - E.  $32\pi/5$
9. Find a scalar function  $f$  so that  $\mathbf{F} = \nabla f$  if

$$\mathbf{F}(x, y, z) = yz\mathbf{i} + xz\mathbf{j} + (xy + 2z)\mathbf{k}.$$

- A.  $f(x, y, z) = xyz + \frac{1}{2}z^2$
  - B.  $f(x, y, z) = xyz$
  - C.  $f(x, y, z) = xyz + z^2$
  - D.  $f(x, y, z) = xy + z^2$
  - E. There is no such scalar function
10. Let  $\mathbf{F}$  denote a vector field and let  $f$  define a scalar function of three variables. Which of the following expression *is* a meaningful expression?
- A.  $\text{div}(\text{grad } f)$
  - B.  $\text{div}(\text{div } \mathbf{F})$ .
  - C.  $\text{curl}(\text{div } \mathbf{F})$
  - D.  $\text{grad}(\text{grad } \mathbf{F})$
  - E.  $\text{grad}(\text{grad } f)$