

DEPARTMENT OF MATHEMATICS

Ma213 - FINAL EXAM Spring 2013

DO NOT TURN THIS PAGE UNTIL YOU ARE INSTRUCTED TO DO SO

There are 9 problems and a total of 10 pages including this one. **You are permitted to use:** two sheets of your own notes (front and back). **These will not be shared.**

Calculators are permitted, so long as they are not capable of wireless communication.

Problem	Maximum Score	Earned Score
1	20	
2	15	
3	15	
4	10	
5	10	
6	10	
7	10	
8	10	
9	10	
WHS	10	
Total	120	

NAME: _____

SECTION No: _____

LAST 4 digits of Student No.: _____

1. 20 points

(a) Find all vectors $w = \langle a, b, c \rangle$ which are perpendicular to both

$$v_1 = \langle 2, -2, 3 \rangle, v_2 = \langle 3, 1, -7 \rangle$$

and such that $c^2 = 64$.

Answer:

(b) Find the equation of a plane through the point $P(2, 3, 5)$ and parallel to both $v_1 = \langle 2, -2, 3 \rangle$ and $v_2 = \langle 3, 1, -7 \rangle$

Answer:

(c) Find the equation of the plane passing through $P(2, 3, 5)$, $Q(4, 1, 8)$ and $R(5, 4, -2)$.

Answer:

(d) What is the area of the triangle formed by $P(2, 3, 5)$, $Q(4, 1, 8)$ and $R(5, 4, -2)$?

Answer:

(e) Find the equation(s) of the line joining $Q(3, 1, 10)$ and $R(5, 5, 4)$. What is the midpoint of the line segment QR ?

Answer:

2. 15 points

Consider the space curve C given by the parametric form:

$$\mathbf{r}(t) = \langle \sin(2t), -t, \cos(2t) \rangle .$$

Answer the following questions.

- (a) Determine the unit tangent \mathbf{T} , and the unit bitangent \mathbf{B} to the curve C at $t = \pi$.

Answer:

- (b) Calculate $\int_C (x + y + z) ds$ where C is the space curve described above as t varies from 0 to 2π .

Answer:

- (c) Calculate the integral $\int_C F \cdot dr$ where C is the space curve described above as t varies from 0 to 2π and $F = \langle 0, y, z \rangle$.

Answer:

3. 10 points

(a) Given the function

$$f(x, y, z) = 5xz - 3xy - 4yz$$

and the substitution:

$$x = uv, y = u^2 + v^2, z = u - v,$$

calculate $\frac{\partial f}{\partial u}$. **You may** leave the answer in terms of x, y, z, u, v .

Answer:

(b) Evaluate (and simplify) the above answers when $u = 1, v = -1$. Here, the final answer **must be** a number.

Answer:

(c) Consider the function $F(x, y) = x^2 + txy + ty^2$, where t is a constant. The point $(0, 0)$ is clearly a critical point for the function $F(x, y)$, for all values of t .

Answer the following. **Hint:** You may ignore the values of t for which the second derivative test is inconclusive.

i. Determine all values of t (if any) for which, $(0, 0)$ is a saddle point. Explain your reasoning. **Hint:** Calculate the D at $(0, 0)$.

Answer:

ii. Determine all points (if any) for which $(0, 0)$ is a local **minimum**. Explain your reasoning.

Answer:

iii. Determine all points (if any) for which $(0, 0)$ is a local **maximum**. Explain your reasoning. **Answer:**

4. 10 points

Consider the function

$$f(x, y, z) = \frac{x}{y} + 2\frac{y}{z} - 2\frac{z}{x}.$$

Answer the following questions:

- (a) Calculate $\nabla(f)$ and the Laplacian $\nabla^2(f) = \nabla \cdot \nabla(f)$.

Answer:

- (b) Calculate $\nabla(f)(1, 1, -1)$ and $\nabla \cdot \nabla(f)(1, 1, -1)$.

Answer:

- (c) Find the directional derivative of f in the direction $\langle 3, -1, -1 \rangle$ at the point $P(1, 1, -1)$.

Answer:

- (d) Find the direction u in which the directional derivative of f at the point $P(1, 1, -1)$ (i.e. $D_u f(1, 1, -1)$) is minimized. What is the value of this minimum derivative?

Answer:

5. 10 points

Consider the part of the paraboloid

$$z = 7 - 3x^2 - 3y^2 \text{ lying above the plane } z = 4.$$

Answer the following questions:

(a) **Carefully sketch** the surface and clearly mark the region of integration.

Answer:

(b) **First set up** the double integral for the surface area in terms of x, y . Describe the region of integration, but it is not necessary to identify the limits here.

Answer:

(c) **Now evaluate** the integral **using an appropriate change of variables**. The limits must be clearly stated here.

Answer:

6. 10 points

- (a) **Carefully sketch** the prism lying above the triangle with corners $A(0, 0, 0)$, $B(2, 0, 0)$, $C(0, 1, 0)$ and bounded by the planes $z = 1$ and $z = 5$. Call the resulting solid R .

- (b) Suppose that the solid R has density given by $\rho(x, y, z) = xyz$. Suppose we calculate the mass of the solid R using the integral

$$\int_{z=a}^b \int_{y=c}^d \int_{x=e}^f (xyz) dx dy dz.$$

Determine the correct limits for this integral and write out the correct expression of the integral.

Answer:

- (c) Determine **the mass** of the solid region R . Be sure to simplify the final answer.

Answer:

- (d) Set up the formula for \bar{z} , the z -coordinate for the center of mass. Your answer should involve an explicit integral and may use the mass found above. **It is not necessary** to evaluate the integral.

Answer:

7. 10 points

(a) Given the vector field

$$F = \langle x^2 - xy, y^2 - 2xy, 2xyz \rangle$$

calculate $\text{curl } F$ and $\text{div } F$.

Answer:

(b) Is F a conservative field? Why?

If it is a conservative field, find a function f such that $\nabla f = F$.

Answer:

(c) Given the function $g(x, y, z) = e^z(x + 2y)$ compute its gradient $\nabla(g)$ and its Laplacian $\nabla \cdot \nabla(g)$.

Answer:

8. 10 points

Use **Green's Theorem** to evaluate the line integral

$$\int_C (\exp(x) \sin(2x) + 2y) dx + (2x + \cos(3y) + y^2) dy$$

where C the positively oriented boundary of the region bounded by the two parabolas:

$$y = 9 - x^2 \quad \text{and} \quad y = -9 + x^2.$$

Carry out the following instructions. Each part will be graded.

- Carefully sketch the region.
- Be sure to show the boundary curve and mark the orientation.
- Explain why the Green's theorem is applicable.

Answer:

9. 10 points

Consider the integral

$$\int_C (3x^2y^3 + 4x^2)dx + (3x^3y^2 + 5y)dy$$

where C is any simple continuous curve in the plane.

Answer the following:

- (a) In your own words, briefly explain what it means to say the the above line integral is independent of path. Be sure to write complete correct sentences.

- (b) Prove that the above integral is independent of path.

Hint: Find a function $F(x, y)$ such that

$$(3x^2y^3 + 4x^2)dx + (3x^3y^2 + 5y)dy = dF = F_x dx + F_y dy.$$

- (c) Use this to find the value of the integral when C is some path from $P(0, 0)$ to $Q(3, 4)$.

Answer: