


Loci

MA 341 - Topics in Geometry
Lecture 25

UK

It's loci with a soft "c" otherwise you get



28-Oct-2011

MA 341

2

Locus

Set of points satisfying a given property is called the locus of that property.

Examples:

1. Circle of radius r centered at O : locus of points at a distance r from O .
2. Locus of points equidistant from 2 points is perpendicular bisector of segment joining the points.


28-Oct-2011

MA 341

3

Locus

3. The locus of points that are equidistant from the sides of a given angle is the angle bisector.
4. Given points A and B and real number r , $0 < r < \pi$, locus of all points X so that $\angle AXB = r$ is the union of two circular arcs.




28-Oct-2011

MA 341

4

Locus

5. The locus of points that are equidistant from a line segment is:




28-Oct-2011

MA 341

5

Locus

6. Given points A and B , consider 2 circles on one side of AB , tangent to AB at A and at B and externally tangent to each other at X . Find the locus of X .




28-Oct-2011

MA 341

6

Locus

Let l be the radical axis through X .
It intersects AB at O .
Claim: $OA = OB = OX$.



28-Oct-2011


MA 341

7

Locus

Locus of points X with desired property lies
on semicircle centered at O with radius
 $AB/2$.

As circles change, we get entire semicircle



28-Oct-2011

MA 341

8

Classical Loci

Ellipse

Hyperbola

Parabola

How many have seen the description of
these as

- conic sections?
- quadratic equations?
- loci of points?

28-Oct-2011

MA 341

9

Conic Sections



28-Oct-2011

MA 341


10

Ellipse

Given two points F_1 and F_2 and $s > 0$.

Find the locus of points, C , so that

$$CF_1 + CF_2 = s$$



28-Oct-2011

MA 341

11

Ellipse

Called an ellipse with foci F_1 and F_2 .
 CF_1 and CF_2 = focal radii

Three cases:

- i) $s < F_1F_2$
- ii) $s = F_1F_2$
- iii) $s > F_1F_2$

i) If $s < F_1F_2$, then by triangle inequality
 $s = CF_1 + CF_2 \geq F_1F_2$. Thus, it is empty.

ii) If $s = F_1F_2$, then the only points will be
those on F_1F_2 . The ellipse is the segment

28-Oct-2011

MA 341

12


Ellipse

Let C be point on ray F_1F_2 so that

$$CF_2 = \frac{1}{2}(s - F_1F_2)$$

$$CF_1 + CF_2 = (CF_2 + F_1F_2) + CF_2 = 2CF_2 + F_1F_2 = s$$

Similar point C' on other side of F_1 .



28-Oct-2011

MA 341

13

Ellipse


M = midpoint of F_1F_2

$\frac{1}{2}s > \frac{1}{2}F_1F_2$ so $\exists C$ on perp bisector so that

$$CM = \sqrt{\left(\frac{s}{2}\right)^2 - \left(\frac{F_1F_2}{2}\right)^2}$$

$$(CF_1)^2 = (F_1M)^2 + \left(\sqrt{\left(\frac{s}{2}\right)^2 - \left(\frac{F_1F_2}{2}\right)^2}\right)^2 = \left(\frac{s}{2}\right)^2$$

$$CF_1 = \frac{s}{2} = CF_2$$




28-Oct-2011

MA 341

14

Ellipse

Use reflective symmetry to complete the curve.



What is the interior of the ellipse?

$$CF_1 + CF_2 < s$$

28-Oct-2011

MA 341


15

Ellipse

Major axis is the line containing F_1 and F_2
 Minor axis is the perpendicular bisector of F_1F_2
 Eccentricity: e or ϵ is the ratio of F_1F_2 to V_1V_2 .

$$\epsilon = \frac{F_1F_2}{V_1V_2}$$

$$0 < \epsilon < 1$$



28-Oct-2011

MA 341

16

Ellipse

What do we get if $\epsilon = 0$?


What do we get if $\epsilon = 1$?

28-Oct-2011

MA 341

17


Ellipse



28-Oct-2011

MA 341

18



Ellipse

$$\sqrt{(x-1)^2 + (y-1)^2} + \sqrt{(x-4)^2 + (y-5)^2} = 7$$

$$\left(\sqrt{(x-1)^2 + (y-1)^2} + \sqrt{(x-4)^2 + (y-5)^2} \right)^2 = 49$$

$$(x-1)^2 + (y-1)^2 + (x-4)^2 + (y-5)^2 + 2\sqrt{(x-1)^2 + (y-1)^2} \sqrt{(x-4)^2 + (y-5)^2} = 49$$

$$2\sqrt{(x-1)^2 + (y-1)^2} \sqrt{(x-4)^2 + (y-5)^2} = 49 - ((x-1)^2 + (y-1)^2 + (x-4)^2 + (y-5)^2)$$

$$2\sqrt{(x-1)^2 + (y-1)^2} \sqrt{(x-4)^2 + (y-5)^2} = 49 - (2x^2 + 2y^2 - 10x - 12y + 43)$$

$$= 6 - (2x^2 + 2y^2 - 10x - 12y)$$

$$\sqrt{(x-1)^2 + (y-1)^2} \sqrt{(x-4)^2 + (y-5)^2} = 3 - (x^2 + y^2 - 5x - 6y)$$

$$\left(\sqrt{(x-1)^2 + (y-1)^2} \sqrt{(x-4)^2 + (y-5)^2} \right)^2 = (3 - (x^2 + y^2 - 5x - 6y))^2$$

28-Oct-2011 MA 341 20

Ellipse

$$((x-1)^2 + (y-1)^2)((x-4)^2 + (y-5)^2) = (3 - (x^2 + y^2 - 5x - 6y))^2$$

$$x^2 + y^2 - 2x - 2y + 2 \mid x^2 + y^2 - 8x - 10y + 41 =$$


$$x^4 + 2x^2y^2 + y^4 - 10x^3 - 12x^2y - 10xy^2 - 12y^3 + 19x^2 + 60xy + 30y^2 + 30x + 36y + 9$$


$$x^4 + 2x^2y^2 + y^4 - 10x^3 - 12x^2y - 10xy^2 - 12y^3 + 59x^2 + 36xy + 63y^2 - 98x - 102y + 82 =$$

$$x^4 + 2x^2y^2 + y^4 - 10x^3 - 12x^2y - 10xy^2 - 12y^3 + 19x^2 + 60xy + 30y^2 + 30x + 36y + 9$$

$$40x^2 - 24xy + 33y^2 - 128x - 138y + 73 = 0$$

28-Oct-2011 MA 341 21





Ellipse

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$$

$$x = x' \cos(\theta) - y' \sin(\theta)$$

$$y = x' \sin(\theta) + y' \cos(\theta)$$

28-Oct-2011 MA 341 24
