
Quadrilateral Geometry


MA 341 - Topics in Geometry Lecture 19





Varignon's Theorem I

The quadrilateral formed by joining the midpoints of consecutive sides of any quadrilateral is a parallelogram.









Starting with any quadrilateral gives us a parallelogram

What type of quadrilateral will give us a square?
a rhombus?
a rectangle?

12-Oct-2011 MA 341

The quadrilateral formed by joining the midpoints of consecutive sides of a quadrilateral whose diagonals are perpendicular is a rectangle.

PQRS is a parallelogram
Each side is parallel to one of
the diagonals
Diagonals perpendicular ⇒
sides of parallelogram are
perpendicular

 \Rightarrow parallelogram is a rectangle.




12-Oct-2011

MA 341

Varignon's Corollary: Rhombus

The quadrilateral formed by joining the midpoints of consecutive sides of a quadrilateral whose diagonals are congruent is a rhombus.

PQRS is a parallelogram
Each side is half of one of the diagonals



12-Oct-2011

MA 341

Varignon's Corollary: Square

The quadrilateral formed by joining the midpoints of consecutive sides of a quadrilateral whose diagonals are congruent and perpendicular is a square.

12-Oct-2011


MA 341

Quadri	lateral	Cente	rs
--------	---------	-------	----

Each quadrilateral gives rise to 4 triangles using the diagonals.

P and Q = centroids of $\triangle ABD$ and $\triangle CDB$ R and S = centroids of $\triangle ABC$ and $\triangle ADC$ The point of intersection of the segments PQ and RS is the centroid of ABCD


12-Oct-2011 MA 341 10





Quadrilateral Centers

The centerpoint of a quadrilateral is the point of intersection of the two segments joining the midpoints of opposite sides of the quadrilateral. Let us call this point O.

12-Oct-2011 MA 341 12








Theorem

The segment joining the midpoints of the diagonals of a quadrilateral is bisected by the centerpoint.



Proof

Need to show that PMRN a parallelogram

In $\triangle ADC$, PN a midline and PN||DC and PN= $\frac{1}{2}DC$

In $\triangle BDC$, MR a midline and MR||DC and MR= $\frac{1}{2}DC$


- \Rightarrow MR||PN and MR=PN
- ⇒PMRN a parallelogram

Diagonals bisect one another. Then MN intersects PR at its midpoint, which we know is O.

12-Oct-2011

12-Oct-2011


MA 341



Theorem

Consider a quadrilateral ABCD and let E, F, G, H be the centroids of the triangles \triangle ABC, \triangle BCD, \triangle ACD, and \triangle ABD.

- 1. EF||AD, FG||AB, GH||BC, and EH||CD;
- 2. $K_{ABCD} = 9 K_{EFGH}$.



M_{BC} = midpoint of BC of $\triangle ABC$ and E lies 2 and M_{BC} , EM_{BC} = 1/3 A	OCB and DF:FM _B ,=2:1 ve EF AD and EF=	E B
12-Oct-2011	MA 341	19

Theorem

The sum of the squares of the lengths of the sides of a parallelogram equals the sum of the squares of the lengths of the diagonals.

AB2+BC2+CD2+AD2=AC2+BD2

12-Oct-2011 MA 341

Proof

By Law of Cosines $\triangle ABE$ $AB^2 = BE^2 + AE^2 - 2AE \cdot BE \cos(BEA)$ Note that $\cos BEA = FE/BE$, so $AB^2 = BE^2 + AE^2 - 2AE \cdot FE$ Apply Stewart's Theorem to $\triangle EBC$ we have $BC^2 = BE^2 + EC^2 + 2EC \cdot FE$ ABCD parallelogram \Rightarrow diagonals bisect each other

Thus AE = EC. Adding the first two equations we get $AB^2 + BC^2 = 2BE^2 + 2AE^2$ Apply this same process to $\triangle CAD$ and we have $CD^2 + AD^2 = 2DE^2 + 2CE^2$.

20

12-Oct-2011 MA 341

Proof

Now, add these equations and recall that AE=EC and BE=ED.

 $AB^2+BC^2+CD^2+AD^2 = BE^2+2AE^2+2DE^2+2CE^2$

 $= 4AE^2 + 4BE^2$

 $= (2AE)^2 + (2BE)^2$

 $= AC^2 + BD^2$

12-Oct-2011 MA 341

Varignon's Theorem II

The area of the Varignon parallelogram is half that of the corresponding quadrilateral, and the perimeter of the parallelogram is equal to the sum of the diagonals of the original quadrilateral.

12-Oct-2011 MA 341

Proof

Recall SP = midline of $\triangle ABD$ and $K_{ASP} = \frac{1}{4} K_{ABD}$

 $K_{DSR} = \frac{1}{4} K_{DAC}$

 $K_{CQR} = \frac{1}{4} K_{CBD}$

 $K_{BPQ} = \frac{1}{4} K_{BAC}$

Therefore,

 $K_{ASP} + K_{DSR} + K_{CQR} + K_{BPQ} = \frac{1}{4} (K_{ABD} + K_{CBD}) + \frac{1}{4} (K_{DAC} + K_{BAC})$

 $= \frac{1}{4} K_{ABCD} + \frac{1}{4} K_{ABCD}$

 $=\frac{1}{2}K_{ABCD}$

12-Oct-2011 MA 341 2

Proof

Then,

$$K_{PQRS} = K_{ABCD} - (K_{ASP} + K_{DSR} + K_{CQR} + K_{BPQ})$$

$$= K_{ABCD} - \frac{1}{2} K_{ABCD}$$

$$= \frac{1}{2} K_{ABCD}$$

Also PQ = $\frac{1}{2}$ AC = SR and SP = $\frac{1}{2}$ BD = QR Easy to see that the perimeter of the Varignon parallelogram is the sum of the diagonals.

12-Oct-2011

MA 341

Wittenbauer's Theorem

Given a quadrilateral ABCD a parallelogram is formed by dividing the sides of a quadrilateral into three equal parts, and connecting and extending adjacent points on either side of each vertex. Its area is 8/9 of the quadrilateral. The centroid of ABCD is the center of Wittenbauer's parallelogram (intersection of the diagonals).

12-Oct-2011

MA 341

26

