

Coordinatization of the Plane

MA 341 - Topics in Geometry
Lecture 29



Background

Used by Egyptians & Romans for surveying

Used by Greeks for mapmaking

Nicole Oresme graphed one variable against another in 14th century

Credit goes to Rene Descartes and Pierre Fermat for formalizing the process of analytic geometry

7-Nov-2011

MA 341

2

The Line

Ruler Postulate guarantees that each line looks like the real line

There is a one-to-one correspondence between each line and the real line

We can pick a point that corresponds to 0, then each point will be given some value - say A gets coordinate a, B gets coordinate b, and $d(A,B) = |b - a|$

7-Nov-2011

MA 341

3

Line

Given two points A and B and $k > 0$ there is at least one point C so that

$$\frac{AC}{CB} = k$$

Assume $a < b$. Find C so that

$$\frac{AC}{CB} = \frac{|c-a|}{|c-b|} = k$$

7-Nov-2011 MA 341 4

Line

$$\frac{AC}{CB} = \frac{|c-a|}{|c-b|} = k \Leftrightarrow \begin{cases} \frac{c-a}{b-c} = k & \text{if } a < c < b \\ \frac{c-a}{c-b} = k & \text{if } c < a \text{ or } c > b \end{cases}$$

$$\Leftrightarrow \begin{cases} c-a = kb - kc & \text{if } a < c < b \\ c-a = kc - kb & \text{if } c < a \text{ or } c > b \end{cases}$$

$$\Leftrightarrow \begin{cases} c = \frac{kb+a}{k+1} & \text{if } a < c < b \\ c = \frac{kb-a}{k-1} & \text{if } c < a \text{ or } c > b, k \neq 1 \end{cases}$$

7-Nov-2011 MA 341 5

Line

Unless $k=1$
There are 2 points c to satisfy the ratio:
one between A and B
other not between A and B

7-Nov-2011 MA 341 6

Line

Note $k=1$ is a special point - the midpoint which can be shown **GEOMETRICALLY** to be unique

a $\frac{a+b}{2}$ b
A M B

7-Nov-2011 MA 341 7

Plane

Let O, X, Y be three non-collinear points in plane.

$OX = x$ -axis, the abscissa
 $OY = y$ -axis, the ordinate
 $O =$ origin

7-Nov-2011 MA 341 8

Plane

Lines run parallel to the axes

$P=(a,b)$
 b
 a
 O

7-Nov-2011 MA 341 9

Plane

Regular assumptions:
 OY comes from OX by a counterclockwise rotation of OX about O. (orientation)

$P_y = (0, b)$
 - projection of P on OY

$P_x = (a, 0)$
 - projection of P on OX

7-Nov-2011 MA 341 10

Plane

There is a one-to-one correspondence between the points in the plane and the set of ordered pairs of real numbers

$$\{ (x, y) \mid x, y \in \mathbb{R} \}$$

7-Nov-2011 MA 341 11

Plane

Given two points A and B and $k > 0$ there is at least one point C that divides the line segment AB in the ratio $k:1$.

Assume $x_A \leq x_B$. Find coordinates of point (x_C, y_C) that internally divides AB in a ratio $k:1$.

7-Nov-2011 MA 341 12

Plane

$AA' \parallel BB' \parallel CC'$

Then $\frac{AC}{CB} = \frac{A'C'}{C'B'}$

thus $x_C = \frac{kx_B + x_A}{k+1}$

7-Nov-2011 MA 341 13

Plane

$AA'' \parallel BB'' \parallel CC'' \parallel OY$

Then $\frac{AC}{CB} = \frac{A''C''}{C''B''}$ thus $y_C = \frac{ky_B + y_A}{k+1}$

7-Nov-2011 MA 341 14

Plane


Thus, $C = \left(\frac{kx_B + x_A}{k+1}, \frac{ky_B + y_A}{k+1} \right)$

Midpoint: $M = \left(\frac{x_B + x_A}{2}, \frac{y_B + y_A}{2} \right)$

7-Nov-2011 MA 341 15

Plane

Normally take OX and OY perpendicular!!




7-Nov-2011 MA 341 16

Distance Formula

$A = (x_A, y_A)$ and $B = (x_B, y_B)$
AB = hypotenuse of right triangle

x-leg = $x_B - x_A$
y-leg = $y_B - y_A$

Pythagorean Theorem


$$AB = \sqrt{(x_B - x_A)^2 + (y_B - y_A)^2}$$


7-Nov-2011 MA 341 17

Lines

$A = (x_A, y_A)$ and $B = (x_B, y_B)$

Slope = rise/run = abscissal/ordinatal

Why is this well-defined?


$$m_{AB} = \frac{y_B - y_A}{x_B - x_A}$$

7-Nov-2011 MA 341 18

Line Theorem

Let l be a line with slope m and let A be a point on l . A point P lies on l if and only if its coordinates satisfy the equation

$$y - y_A = m(x - x_A)$$

Proof: Let $P = (x_P, y_P)$ lie on l , $P \neq A$.

Then $l = AP$, so

$$m = \frac{y_P - y_A}{x_P - x_A}$$

Therefore, (x_P, y_P) satisfies the equation

$$y - y_A = m(x - x_A)$$

7-Nov-2011

MA 341

19

Line Theorem

Proof: Assume that $P = (x_P, y_P)$ satisfies the equation

$$y - y_A = m(x - x_A)$$

Therefore

$$y_P - y_A = m(x_P - x_A)$$

Let P' lie on l and let P' have the same x -coordinate as P : $x_{P'} = x_P$

Then

$$\begin{aligned} y_{P'} - y_A &= m(x_{P'} - x_A) \\ &= m(x_P - x_A) \\ &= y_P - y_A \end{aligned}$$

$$y_{P'} = y_P \quad \text{and } P = P'$$

7-Nov-2011

MA 341

20

Equations of Lines

$y - y_0 = m(x - x_0)$ - point-slope form
 $y = mx + b$ - slope-intercept form

$$\frac{x}{a} + \frac{y}{b} = 1 \quad \text{- xy intercept form}$$

$ax + by + c = 0$ - standard form

Corollary: Lines are parallel if and only if they have the same slope or no slope.

7-Nov-2011

MA 341

21

Inclination of Lines

For any non-horizontal line define its inclination to be the smallest positive measure, θ , of angle measured counterclockwise from the positive x-axis.

If horizontal define its inclination to be 0.

Note: $0^\circ \leq \theta \leq 180^\circ$

$$m = \tan \theta$$

7-Nov-2011

MA 341

22

Intersection of Lines

Let α be angle between intersecting non-vertical lines l_1 and l_2 having slopes m_1 and m_2 . Then $\alpha = 0$ iff $m_1 m_2 = -1$. Otherwise

$$\tan(\alpha) = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

7-Nov-2011

MA 341

23

Intersection of Lines

$$m_1 = \tan(\theta_1) \text{ and } m_2 = \tan(\theta_2)$$

$$\theta_1 = \theta_2 + (180 - \alpha)$$

$$\theta_1 - \theta_2 = (180 - \alpha)$$

$$\tan(\alpha) = |\tan(180 - \alpha)|$$


$$= |\tan(\theta_1 - \theta_2)|$$

$$= \left| \frac{\tan(\theta_1) - \tan(\theta_2)}{1 + \tan(\theta_1)\tan(\theta_2)} \right|$$

7-Nov-2011

MA 341

24



Distance from a Point to a Line

If a vertical or horizontal line, easy.

$P = (x_p, y_p)$ and line $l: ax + by + c = 0$

Assume $ab \neq 0$. Then $m = -a/b$, so line through P perpendicular to l is $y - y_p = b/a(x - x_p)$

Find point of intersection:

$$x_q = \frac{b^2 x_p - ac - aby_p}{a^2 + b^2} \quad y_q = \frac{a^2 y_p - bc - abx_p}{a^2 + b^2}$$

$$x_p - x_q = \frac{a(ax_p + by_p + c)}{a^2 + b^2} \quad y_p - y_q = \frac{b(ax_p + by_p + c)}{a^2 + b^2}$$

7-Nov-2011

MA 341

25

Distance from a Point to a Line

$$d(P, l) = d(P, Q) = \sqrt{(x_p - x_q)^2 + (y_p - y_q)^2}$$

$$= \sqrt{\left(\frac{a(ax_p + by_p + c)}{a^2 + b^2}\right)^2 + \left(\frac{b(ax_p + by_p + c)}{a^2 + b^2}\right)^2}$$

$$= \sqrt{\frac{(a^2 + b^2)(ax_p + by_p + c)^2}{(a^2 + b^2)^2}}$$

$$= \frac{|ax_p + by_p + c|}{\sqrt{a^2 + b^2}}$$

7-Nov-2011

MA 341

26

Applications

Medians are concurrent and centroid divides each in a 2:1 ratio.

- a) Brute force method - find midpoint of each side, equation of each median, points of intersection of each pair of lines - algebraically tedious

$$G = \left(\frac{x_a + x_b + x_c}{3}, \frac{y_a + y_b + y_c}{3} \right)$$

7-Nov-2011

MA 341

27

Applications

b) Create coordinate system so that median of AC is origin, $A=(a,0)$ and $B=(0,1)$ and $C=(-a,0)$

Gives simple equations for each median

c) Don't find equations of lines at all, but use earlier result on ratios. D =midpoint of BC

so

$$D = \left(\frac{x_b + x_c}{2}, \frac{y_b + y_c}{2} \right)$$

Let P =point on AD so that $AP/PD=2/1$.

$$P = \left(\frac{x_a + 2x_d}{3}, \frac{y_a + 2y_d}{3} \right) = \left(\frac{x_a + x_b + x_c}{3}, \frac{y_a + y_b + y_c}{3} \right)$$

7-Nov-2011

MA 341

28

Applications

Given two intersecting lines and a positive number k , find the locus of all points the sum of whose distances to the lines is k .

Choose OX and OY so that they are the angle bisectors of the angles formed by the two lines.

CLAIM: OX and OY are perpendicular.

Line l_1 has equation $y = mx$, $m > 0$, line l_2 has equation $y = -mx$.

7-Nov-2011

MA 341

29

Applications

(x,y) on the locus if and only if

$$d((x,y),l_1) + d((x,y),l_2) = k$$

$$\frac{|mx - y|}{\sqrt{m^2 + 1}} + \frac{|-mx - y|}{\sqrt{(-m)^2 + 1}} = k$$

$$\text{Let } K = k\sqrt{m^2 + 1}$$

We have equation: $|mx - y| + |mx + y| = K$

Note: If (x,y) satisfies this then so do $(-x,y)$, $(-x,-y)$ and $(x,-y)$

We only need work in first quadrant

7-Nov-2011

MA 341

30

Applications

In first quadrant $mx + y \geq 0$, so
 $|mx - y| + mx + y = K$
 Reduces to 2 cases: $mx - y \geq 0$ or $mx - y \leq 0$

If $0 \leq y \leq mx$, then
 $|mx - y| + mx + y = mx - y + mx + y = K$
 $x = K/(2m)$
 and the graph is a vertical segment

If $0 \leq mx < y$, then
 $|mx - y| + mx + y = -mx + y + mx + y = K$
 $y = K/2$ and the graph is a horizontal segment

7-Nov-2011 MA 341 31

Applications

7-Nov-2011 MA 341 32
