

Math 654 - Algebraic Topology

Homework 4

Fall 2015

1. Recall the functor $\mathrm{Gl}_n : \mathbf{Comm} \rightarrow \mathbf{Gp}$ from HW1. When $n = 1$, this gives the functor $(-)^{\times} : \mathbf{Comm} \rightarrow \mathbf{Gp}$ which takes a commutative ring R and gives R^{\times} , the units (invertible elements) in R . Show that the determinant yields a natural transformation $\det : \mathrm{Gl}_n \rightarrow (-)^{\times}$.
2. Let G be a group. Define a category \star_G which has a single object, \star , and such that $\mathrm{Hom}(\star, \star) = G$. The identity morphism and composition of morphisms are defined to be the identity element of the group and the group multiplication, respectively.
 - (a) Show that a G -set X is the same data as a functor $\mathcal{X} : \star_G \rightarrow \mathbf{Set}$.
 - (b) Show that if X and Y are G -sets, then a G -equivariant function $f : X \rightarrow Y$ corresponds precisely to a natural transformation of functors $\mathcal{X} \rightarrow \mathcal{Y}$.
3. Let $\mathcal{I} = \{\bullet \rightarrow \bullet\}$ be the category with two objects and a single non-identity morphism. Describe the data involved in a natural transformation $\eta : F \Rightarrow G : \mathcal{I} \rightarrow \mathcal{C}$.
4. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Let $G : \mathrm{Ob}(\mathcal{C}) \rightarrow \mathrm{Ob}(\mathcal{D})$ be a function, and suppose given an isomorphism $\eta_C : F(C) \cong G(C)$ for each $C \in \mathcal{C}$. Show that there is a unique way to define G on morphisms of \mathcal{C} that makes $\{\eta_C\}$ a natural isomorphism.