## Math 654 - Algebraic Topology Homework 5 Fall 2016

- 1. Recall the functor  $Gl_n : \mathbf{Comm} \to \mathbf{Gp}$  from HW1. When n = 1, this gives the functor  $(-)^\times : \mathbf{Comm} \to \mathbf{Gp}$  which takes a commutative ring R and gives  $R^\times$ , the units (invertible elements) in R. Show that the determinant yields a natural transformation  $\det : Gl_n \to (-)^\times$ .
- 2. Let G be a group. Define a category  $\star_G$  which has a single object,  $\star$ , and such that  $\operatorname{Hom}(\star,\star)=G$ . The identity morphism and composition of morphisms are defined to be the identity element of the group and the group multiplication, respectively.
  - (a) Show that a *G*-set *X* is the same data as a functor  $\mathcal{X} : \star_G \longrightarrow \mathbf{Set}$ .
  - (b) Show that if X and Y are G-sets, then a G-equivariant function  $f: X \longrightarrow Y$  corresponds precisely to a natural transformation of functors  $\mathcal{X} \longrightarrow \mathcal{Y}$ .
- 3. Let  $\mathscr{I} = \{ \bullet \longrightarrow \bullet \}$  be the category with two objects and a single non-identity morphism. Describe the data involved in a natural transformation  $\eta : F \Rightarrow G : \mathscr{I} \longrightarrow \mathscr{C}$ .
- 4. Let  $F: \mathscr{C} \longrightarrow \mathscr{D}$  be a functor. Let  $G: \mathrm{Ob}(\mathscr{C}) \longrightarrow \mathrm{Ob}(\mathscr{D})$  be a function, and suppose given an isomorphism  $\eta_C: F(C) \cong G(C)$  for each  $C \in \mathscr{C}$ . Show that there is a unique way to define G on morphisms of  $\mathscr{C}$  that makes  $\{\eta_C\}$  a natural isomorphism.