

Math 654 - Algebraic Topology

Homework 8

Fall 2019

1. (Adjoint functors) A pair of functors $L: \mathcal{C} \rightleftarrows \mathcal{D}: R$ is said to be an **adjoint pair** if there is a bijection

$$\text{Hom}_{\mathcal{D}}(L(X), Y) \cong \text{Hom}_{\mathcal{C}}(X, R(Y))$$

that is natural in both $X \in \mathcal{C}$ and $Y \in \mathcal{D}$. The functor L is called the **left adjoint**, while R is the **right adjoint**.

- (a) Show that any left adjoint automatically preserves coproducts. That is, show that there is an isomorphism $L(\coprod_i X_i) \cong \coprod_i L(X_i)$.
- (b) Show that any left adjoint preserves pushouts. That is, the diagram

$$\begin{array}{ccc} L(A) & \longrightarrow & L(Y) \\ \downarrow & & \downarrow \\ L(X) & \longrightarrow & L(X \amalg_A Y) \end{array}$$

is automatically a pushout diagram.

2. (a) Show that abelianization $(-)_{ab} : \mathbf{Gp} \rightarrow \mathbf{Ab}$ is left adjoint to the inclusion $\mathbf{Ab} \hookrightarrow \mathbf{Gp}$.
- (b) Use this to show that if $N \trianglelefteq G$, then $(G/N)_{ab} \cong G_{ab}/N_{ab}$.
3. (a) Show that the free abelian group construction $\mathbb{Z}\{-\} : \mathbf{Set} \rightarrow \mathbf{Ab}$ is left adjoint to the “forgetful functor” $U : \mathbf{Ab} \rightarrow \mathbf{Set}$ that forgets the group structure and remembers only the underlying set.
- (b) Conclude that $\mathbb{Z}\{A \amalg B\} \cong \mathbb{Z}\{A\} \oplus \mathbb{Z}\{B\}$.
- (c) Show that $\mathbb{Z}\{A \times B\} \cong \mathbb{Z}\{A\} \otimes \mathbb{Z}\{B\}$. (Hint: $U(\text{Hom}(\mathbb{Z}\{Y\}, A)) \cong A^Y$.)