Name:	Section and/or TA:
Name.	Section and of TA.

Do not remove this answer page. You will return the whole exam. You will be allowed two hours to complete this test. No books or notes may be used. You may use a graphing calculator during the exam, but NO calculator with a Computer Algebra System (CAS) or communication capabilities is permitted. You may not use a phone or other communication device during the exam.

The exam consists of 12 multiple choice questions that are worth 5 points each and 4 free response questions that are worth 10 points each. You should work the multiple choice questions on the question page. After you have checked your work carefully, record your answers by completely filling in the circle below that corresponds to your answer. If you must change your answer, make a note on the front of the exam. Be sure to check carefully when you transfer your answers to the cover sheet.

Show all work to receive full credit on the free response problems. You do not need to compute a decimal approximation to your answer. For example, the answer  $4\pi$  is preferred to 12.57.

## Multiple Choice Questions

1	A B C D E	7 A B C D E
2	A B C D E	8 A B C D E
3	A B C D E	9 A B C D E
4	A B C D E	<b>10</b> (A) (B) (C) (D) (E)
5	A B C D E	<b>11</b> (A) (B) (C) (D) (E)
6	A B C D E	<b>12</b> (A) (B) (C) (D) (E)

## SCORE

Multiple					Total
Choice	13	14	15	16	Score
60	10	10	10	10	100

This page is left blank.

## Multiple Choice Questions

1. (5 points) Let

$$f(x) = \begin{cases} 1 + x - cx^2 & x < 1\\ \frac{cx}{1+x} & x \ge 1 \end{cases}$$

For what value(s) of c is this function continuous?

- A. c = 3/4
- B. c = 4/3
- C. c = 2
- D. c = 1/3
- E. There is no value of c for which f is continuous

2. (5 points) A function f satisfies  $-x^2 + 3x + 14 \le f(x) \le x^2 + 7x + 16$  for all real numbers x. There is exactly one real number c where we may use the Squeeze Theorem to compute the limit  $\lim_{x\to c} f(x) = L$ . Find c and L.

- A. c = 0 and L = 14
- B. c = 1 and L = 16
- C. c = -2 and L = 4
- D. c = -1 and L = 10
- E. c = 2 and L = 34

- 3. (5 points) Consider the limit  $\lim_{h\to 0} \frac{\sec(\pi+h)-\sec(\pi)}{h} = L$ . Select the correct statement.
  - A. The value L is the derivative of sec(x) at x = 0.
  - B. The value L is the derivative of  $sec(\pi + x)$  at  $x = \pi$ .
  - C. The value L is the derivative of sec(x + h) at  $x = \pi$ .
  - D. The value L is the derivative of sec(x) at  $x = \pi$ .
  - E. The value L is the derivative of sec(x) at x = h.


- 4. (5 points) Consider the ellipse defined by  $x^2 xy + 2y^2 = 4$ . Find the tangent line to this curve at the point (x, y) = (2, 0).
  - A. y = 2x + 4
  - B. y = 2x 4
  - C. y = 2x 2
  - D. y = -2x 4
  - E. y = -2x + 4

- 5. (5 points) The height (in meters) of a bullet fired in the air vertically from ground level is  $s(t) = -5t^2 + 120t$ . Find the bullet's maximum height.
  - A. 720 meters
  - B. 100 meters
  - C. 125 meters
  - D. 5 meters
  - E. 120 meters

- 6. (5 points) The *derivative* of f is  $f'(x) = \frac{x(x^2 81)}{\sin^2(x) + 1}$ . Find the interval or intervals where f is increasing.
  - A. (-9,0) and  $(9,\infty)$
  - B.  $(-\infty, -9)$  and (0, 9)
  - C. (-9,9)
  - D.  $(-\infty, 0)$  and  $(9, \infty)$
  - E.  $(0, \infty)$

- 7. (5 points) Consider the function
- $f(x) = \frac{1}{x}$
- defined on the interval [1, 4].

Estimate  $\int_1^4 \frac{1}{x} dx$  using <u>left</u> endpoints for n = 3 approximating rectangles all having bases of the same length, as shown in the picture.



- A. ln(4)
- B. 5/6
- C. 1/2
- D. 13/12
- E. 11/6
- 8. (5 points) Determine which of the following integrals is equal to the given limit without evaluating the limit:

$$\lim_{n \to \infty} \sum_{i=1}^{n} \frac{10}{n} \sin\left(3 + \frac{5i}{n}\right)$$

- A.  $\int_{3}^{5} 10 \sin(x) dx$
- $B. \int_3^8 2\sin(x) \, dx$
- C.  $\int_{3}^{8} \sin(x) dx$
- D.  $\int_{3}^{5} \sin(x+3) \, dx$
- $E. \int_{5}^{10} \sin(x) \, dx$

9. (5 points) Use L'Hôpital's rule and the Fundamental Theorem of Calculus to evaluate the following limit

$$\lim_{x\to 0} \frac{\int_0^x (7-7\cos(t))\,dt}{x^3}$$

- A. 7/3
- B. 7/6
- C. 0
- D. 1
- E. The limit does not exist

10. (5 points) Assuming that  $\int_{1}^{6} f(x) dx = \frac{5}{2}$  and  $\int_{6}^{4} f(x) dx = \frac{3}{2}$ , find  $\int_{1}^{4} f(x) dx$ .

- A. 1
- B. 8
- C. 5
- D. 1/2
- E. 4

11. (5 points) Use the Fundamental Theorem of Calculus to find the function f that verifies the following equation

$$\int_0^x \frac{f(t)}{\cos(t)} dt = \sin(x)$$

- A.  $f(x) = \ln(\cos(x))$
- B.  $f(x) = \sin(x)\cos(x)$
- C.  $f(x) = \frac{(\sin(x))^2}{2}$
- D.  $f(x) = \cos^2(x)$
- $E. f(x) = \sin^2(x)$

12. (5 points) An object is moving along a line so that its velocity at time t is  $v(t) = -t^2 + 3$  meters/second. Find the change in position between t = 0 and t = 6 seconds. Assume that displacement to the right is positive.

- A. 6 meters to the right
- B. 54 meters to the left
- C. 6 meters to the left
- D. 3 meters to the left
- E. 54 meters to the right

Free response questions: Show work clearly with proper notation.

- 13. (10 points) Compute the derivatives, you do not need to simplify your answers
  - (a)  $\frac{d}{dx}(\cos(x)\ln(x^2))$
  - (b)  $\frac{d}{dx} \left( \frac{x+5}{x+7} \right)$
  - (c)  $\frac{d}{dt}\sqrt{e^{3t}+1}$

MA 113 Exam 4 Fall 2024

14. (10 points) Let 
$$f(x) = \int_0^x \frac{(16-t^2)}{\sin^2(t) + 1} dt$$
.

- (a) Find f'.
- (b) Find the intervals where f is increasing and decreasing.
- (c) Find the locations of the local maxima or minima, if any.

- 15. (10 points) Evaluate the following integrals using substitution. You must clearly show steps of the substitution to receive full credit.
  - (a)  $\int \frac{\cos(8x)}{\sin(8x)} \, dx$
  - (b)  $\int \frac{2}{x} \ln(x) \, dx$

- 16. (10 points) Below are the graphs of f(x) = 2x + 2 and  $g(x) = 2x^2 + 2x$ .
  - (a) Set up an integral whose value is the area of the shaded region.
  - (b) Evaluate your integral to find the area of the region.

