

Name: \_\_\_\_\_ Section: \_\_\_\_\_

Do not remove this answer page — you will return the whole exam. You will be allowed two hours to complete this test. No books or notes may be used. You may use a graphing calculator during the exam, but NO calculator with a Computer Algebra System (CAS) or a QWERTY keyboard is permitted. Absolutely no cell phone use during the exam is allowed.

The exam consists of 10 multiple choice questions and 5 free response questions. Record your answers to the multiple choice questions on this page by filling in the circle corresponding to the correct answer.

Show all work to receive full credit on the free response problems. You should also show your work on the multiple choice questions as it will make it easier for you to check your work. You should give exact answers, rather than a decimal approximation unless the problem asks for a decimal answer. Thus, if the answer is  $2\pi$ , you should not give a decimal approximation such as 6.283 as your final answer.

## Multiple Choice Questions

**1**     A     B     C     D     E**2**     A     B     C     D     E**3**     A     B     C     D     E**4**     A     B     C     D     E**5**     A     B     C     D     E**6**     A     B     C     D     E**7**     A     B     C     D     E**8**     A     B     C     D     E**9**     A     B     C     D     E**10**     A     B     C     D     E

Multiple Choice	11	12	13	14	15	Total Score
50	10	10	10	10	10	100

This page is intentionally left blank.

## Multiple Choice Questions

1. (5 points) Which of the following is equal to  $\int x f(x) dx$ ?

A.  $f(x) + \int x^2 f(x) dx$

B.  $\frac{x^2}{2} f(x) + \frac{1}{2} \int x^2 f'(x) dx$

C.  $\frac{x^2}{2} f(x) - \frac{1}{2} \int x^2 f'(x) dx$

D.  $x f(x) - \frac{1}{2} \int x^2 f'(x) dx$

E.  $x f(x) + \frac{1}{2} \int x^2 f'(x) dx$

2. (5 points) Find  $\int x \sin(2x) dx$ .

A.  $\frac{1}{2} x^2 \sin(2x) + \frac{1}{4} \cos(2x) + C$

B.  $-\frac{1}{2} x \cos(2x) + \frac{1}{4} \sin(2x) + C$

C.  $\frac{1}{2} x \cos(2x) - \frac{1}{4} \sin(2x) + C$

D.  $\frac{1}{2} x \sin(2x) - \frac{1}{4} \cos(2x) + C$

E.  $2x \sin(2x) - \cos(2x) + C$

3. (5 points) Suppose that  $x = 2 \tan(u)$  and  $-\pi/2 < u < \pi/2$ . Express  $\cos(u)$  in terms of  $x$ .

- A.  $\frac{2}{\sqrt{4+x^2}}$   
B.  $\frac{1}{\sqrt{4+x^2}}$   
C.  $\sqrt{4+x^2}$   
D.  $\frac{\sqrt{1+x^2}}{2}$   
E.  $\frac{x}{\sqrt{4+x^2}}$

4. (5 points) Choose the best substitution to evaluate the integral  $\int \frac{x}{\sqrt{4-x^2}} dx$ .

- A.  $u = 4 - x^2$   
B.  $x = 4 - u^2$   
C.  $x = 2 \tan(u)$   
D.  $u = 2 \tan(x)$   
E.  $u = 2 \sin(x)$

5. (5 points) Give the partial fractions decomposition for the function

$$\frac{x+4}{x^2-4}.$$

- A.  $\frac{3}{x-2} - \frac{1}{x+2}$   
B.  $\frac{3}{2(x-2)} + \frac{1}{2(x+2)}$   
C.  $\frac{3}{2(x-2)} - \frac{1}{2(x+2)}$   
D.  $\frac{1}{2(x-2)} - \frac{1}{2(x+2)}$   
E.  $\frac{1}{2(x-2)} - \frac{3}{2(x+2)}$

6. (5 points) Consider the function  $f(x) = \frac{1}{(x+2)^2(x^2+1)^2}$ . Which of the following terms does not appear in the partial fractions decomposition of  $f$ ?

A.  $\frac{A}{(x+2)^3}$

B.  $\frac{B}{(x+2)^2}$

C.  $\frac{C}{x+2}$

D.  $\frac{D_1x + D_2}{x^2 + 1}$

E.  $\frac{E_1x + E_2}{(x^2 + 1)^2}$

7. (5 points) Use the following information and Simpson's rule  $S_4$  to find an approximate value for the integral  $\int_1^3 f(x) dx$ .

$x$	1	1.5	2	2.5	3
$f(x)$	3	6	12	9	6

A.  $29/2$


B.  $31/2$

C.  $33/2$

D. 15

E. 16

8. (5 points) We consider  $R_n$ ,  $L_n$ ,  $T_n$  and the value of the integral  $I = \int_0^2 f(x)$  for the function whose graph is pictured. Which of the following statements is correct?



- A.  $I \leq L_n \leq R_n \leq T_n$   
 B.  $L_n \leq T_n \leq R_n \leq I$   
 C.  $R_n \leq I \leq T_n \leq L_n$   
 D.  $L_n \leq T_n \leq I \leq R_n$   
 E.  $L_n \leq I \leq T_n \leq R_n$

9. (5 points) Determine if the improper integral  $\int_1^{\infty} \frac{1}{x^2} dx$  exists and, if it exists, give its value.

- A. 1  
 B. 2  
 C. 1/2  
 D. -1  
 E. The integral does not exist.

10. (5 points) Determine if the improper integral  $\int_{-1}^1 \frac{1}{x} dx$  exists and, if it exists, give its value.

- A. -1  
 B. 1  
 C. 0  
 D.  $2e$   
 E. The integral does not exist.

## Free Response Questions

11. (a) (8 points) Find  $\int \cos^4(x) dx$ .

(b) (2 points) Compute the definite integral  $\int_0^{\pi/2} \cos^4(x) dx$ .

12. (10 points) Find the anti-derivative  $\int \frac{1}{(4-x^2)^{3/2}} dx$ . Simplify to give an answer that does not include trigonometric functions.

13. (a) (7 points) Find the partial fractions decomposition of  $\frac{x^2 + 3x + 1}{x(x + 1)^2}$ .
- (b) (3 points) Find  $\int \frac{x^2 + 3x + 1}{x(x + 1)^2} dx$ .

14. (a) (7 points) Find the anti-derivative  $\int x^2 e^{-x} dx$ .
- (b) (3 points) Determine if the improper integral  $\int_0^{\infty} x^2 e^{-x} dx$  exists and, if it exists, find its value.

15. Consider the integral  $\int_1^3 \frac{1}{x} dx$ .

- (a) (4 points) Use the Trapezoid rule with  $n = 4$  to find an approximate value for this integral.
- (b) (6 points) If we use the Trapezoid rule with  $n$ , then the error satisfies

$$|T_n - \int_a^b f(x) dx| \leq \frac{K(b-a)^3}{12n^2}$$

where  $K$  is a number that satisfies  $|f''(x)| \leq K$  for  $a \leq x \leq b$ .

Use this estimate to find a value for  $n$  for which we have  $|T_n - \int_1^3 \frac{1}{x} dx| \leq \frac{1}{500}$ .