

MA137 — Calculus 1 with Life Science Applications

Derivatives of Logarithmic Functions and Logarithmic Differentiation

(Section 4.10)

Department of Mathematics
University of Kentucky

The Derivative of the Natural Logarithmic Function

Theorem

The function $\ln x$ is differentiable for all $x > 0$, and $\frac{d}{dx} \ln x = \frac{1}{x}$.

In particular, if $g(x)$ is a differentiable function, it follows from the chain rule that

$$\frac{d}{dx} \ln g(x) = \frac{1}{g(x)} g'(x).$$

We can use the derivative of e^x and the relationship between the exponential and the natural logarithmic functions to find the derivative of the function $\ln x$. Namely, we start by taking the derivative with respect to x of both sides of $e^{\ln x} = x$. We obtain

$$\frac{d}{dx} e^{\ln x} = \frac{d}{dx} x \iff e^{\ln x} \frac{d}{dx} \ln x = 1 \iff \frac{d}{dx} \ln x = \frac{1}{e^{\ln x}} = \frac{1}{x}$$

Alternative Proof

We use the formal definition of the derivative and $e^x = \lim_{u \rightarrow \infty} \left(1 + \frac{x}{u}\right)^u$

$$\begin{aligned} \frac{d}{dx} \ln x &\stackrel{\text{def}}{=} \lim_{h \rightarrow 0} \frac{\ln(x+h) - \ln x}{h} \\ &\stackrel{\text{ln prop.}}{=} \lim_{h \rightarrow 0} \frac{1}{h} \ln \left(\frac{x+h}{x} \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \frac{x}{h} \ln \left(1 + \frac{1}{x/h} \right) \quad u = x/h \\ &\stackrel{\text{laws}}{=} \frac{1}{x} \lim_{u \rightarrow \infty} \ln \left(1 + \frac{1}{u} \right)^u \\ &\stackrel{\text{cont.}}{=} \frac{1}{x} \ln \left[\lim_{u \rightarrow \infty} \left(1 + \frac{1}{u} \right)^u \right] \\ &= \frac{1}{x} \ln e = \frac{1}{x} \end{aligned}$$

The Derivative of ANY Logarithmic Function

Theorem

The function $\log_a x$ is differentiable for $x > 0$, and $\frac{d}{dx} \log_a x = \frac{1}{(\ln a)x}$.

In particular, if $g(x)$ is a differentiable function, it follows from the chain rule that

$$\frac{d}{dx} \log_a g(x) = \frac{1}{(\ln a)g(x)} g'(x).$$

From the base change formula for logarithms we have that

$$\log_a x = \frac{\ln x}{\ln a}$$

Thus it is enough to find the derivative of $\ln x$. Hence the formula.

Example 1: (Nuehauser, Problems # 28/34/52, p. 204)Find $\frac{dy}{dx}$ when $y = \ln(1 - x^2)$.Find $\frac{dy}{dx}$ when $y = [\ln(1 - x^2)]^3$.Find $\frac{dy}{ds}$ when $y = \ln(\ln s)$.

(a) $y = \ln(1 - x^2)$

$$\frac{dy}{dx} = \frac{1}{1-x^2} \cdot (-2x) = \boxed{\frac{-2x}{1-x^2}}$$

chain rule

(b) $y = [\ln(1 - x^2)]^3$

$$\begin{aligned} \frac{dy}{dx} &= 3 [\ln(1 - x^2)]^{3-1} \cdot (\ln(1 - x^2))' \\ &= 3 [\ln(1 - x^2)]^2 \cdot \frac{1}{1-x^2} \cdot (-2x) = \boxed{\frac{-6x[\ln(1 - x^2)]^2}{1-x^2}} \end{aligned}$$

(c) $y = \ln(\ln s)$

$$\frac{dy}{ds} = \frac{1}{\ln s} \cdot (\ln s)' = \boxed{\frac{1}{\ln s} \cdot \frac{1}{s}}$$

$y = \log(3x^2 - x + 2)$

$$\frac{dy}{dx} = \frac{1}{\ln(10) \cdot (3x^2 - x + 2)} \cdot (6x - 1)$$

Recall the formula :

$$\frac{d}{dx} \left[\log_a g(x) \right] = \left[\frac{1}{\ln(a)} \cdot \frac{1}{g(x)} \right] \cdot g'(x)$$

Example 2: (Nuehauser, Problem # 56, p. 204)Find $\frac{dy}{dx}$ when $y = \log(3x^2 - x + 2)$.[Note: $\log = \log_{10}$]

Example 3: (Nuehauser, Problem # 62, p. 204)

Assume that $f(x)$ is differentiable with respect to x . Show that

$$\frac{d}{dx} \ln \left[\frac{f(x)}{x} \right] = \frac{f'(x)}{f(x)} - \frac{1}{x}$$

$$y = \ln \left[\frac{f(x)}{x} \right]$$

We want to compute $\frac{dy}{dx}$.

1st method

$$y' = \frac{1}{\frac{f(x)}{x}} \cdot \left[\frac{f(x)}{x} \right]' = \frac{x}{f(x)} \cdot \frac{f'(x) \cdot x - f(x)}{x^2}$$

$$= \frac{f'(x)x - f(x)}{xf(x)} = \frac{\cancel{f'(x)}}{\cancel{f(x)}} - \frac{1}{x}$$

2nd method

$$y = \ln \left[\frac{f(x)}{x} \right] = \underline{\ln f(x)} - \underline{\ln x}$$

$$\text{Hence } y' = \frac{1}{f(x)} \cdot f'(x) - \frac{1}{x} = \boxed{\frac{f'(x)}{f(x)} - \frac{1}{x}}$$

Logarithmic Differentiation

In 1695, Leibniz introduced logarithmic differentiation, following Johann Bernoulli's suggestion to find derivatives of functions of the form

$$y = [f(x)]^x.$$

Bernoulli generalized this method and published his results two years later.

The **basic idea** is to take logarithms on both sides and then to use implicit differentiation.

Example 4: (Neuhauer, Example # 12, p. 202)

Find $\frac{dy}{dx}$ when $y = x^x$.

What about $\frac{d}{dx} [(2x)^{2x}]$?

$$y = x^x$$

to find y' we can take \ln of both sides:

$$\ln y = \ln x^x \quad \text{OR} \quad \ln y = x \cdot \ln x$$

Hence, now take $\frac{d}{dx}$ of both sides and use the chain rule:

$$\frac{d}{dx} [\ln y] = \frac{d}{dx} [x \cdot \ln x]$$

$$\frac{1}{y} \cdot \frac{dy}{dx} = 1 \cdot \ln x + x \cdot \frac{1}{x}$$

$$\frac{dy}{dx} = y \left[\ln x + 1 \right] = \underline{\underline{x^x [\ln x + 1]}}$$

About $y = (2x)^{2x}$

1st method: $\ln y = \ln [(2x)^{2x}] \iff \ln y = 2x \cdot \ln(2x)$

Take the derivative: $\frac{d}{dx} [\ln y] = \frac{d}{dx} [2x \cdot \ln(2x)]$

$$\iff \frac{1}{y} \cdot \frac{dy}{dx} = 2 \cdot \ln(2x) + 2x \cdot \left[\frac{1}{2x} \cdot 2 \right]$$

$$\frac{dy}{dx} = y \left[2 \ln(2x) + 2 \right] = (2x)^{2x} \left[\underline{\underline{2 \ln(2x) + 2}} \right]$$

2nd method: $y = (2x)^{2x} = e^{\ln[(2x)^{2x}]} = e^{2x \cdot \ln(2x)}$

Hence $y' = e^{2x \cdot \ln(2x)} \cdot \left[2x \cdot \ln(2x) \right]' \\ = e^{\ln(2x)^{2x}} \cdot \left[2 \ln(2x) + 2x \cdot \left(\frac{1}{2x} \cdot 2 \right) \right] \\ = (2x)^{2x} \cdot \left[\underline{\underline{2 \ln(2x) + 2}} \right]$

Alternative method:

$$y = x^x = e^{\ln x^x} = e^{\underline{\underline{x \ln x}}}$$

$$\text{Hence } y' = e^{\underline{\underline{x \ln x}}} \cdot (x \ln x)'$$

$$= e^{\underline{\underline{x \ln x}}} \cdot \left(1 \cdot \ln x + x \cdot \frac{1}{x} \right)$$

$$= e^{\underline{\underline{x \ln x}}} \cdot (\ln x + 1)$$

$$= x^x \cdot (\ln x + 1)$$

same answer as before !!!

Derivatives of Logarithmic Functions
Logarithmic Differentiation

Theory
Examples

Example 5: (Neuhäuser, Problems # 68/75/76, p. 204)

Use logarithmic differentiation to find the first derivative of the functions

$$y = (\ln x)^{3x}$$

$$y = x^{\cos x}$$

$$y = (\cos x)^x$$

(a) $y = (\ln x)^{3x}$

Take $\ln y$ of both sides: $\ln y = \ln[(\ln x)^{3x}]$

so $\ln y = 3x \cdot \ln[\ln(x)]$. Take $\frac{d}{dx}$ of both sides:

$$\frac{d}{dx}[\ln y] = \frac{d}{dx}[3x \cdot \ln[\ln(x)]]$$

$$\frac{1}{y} \frac{dy}{dx} = 3 \cdot \ln(\ln(x)) + 3x \cdot \left(\frac{1}{\ln x} \cdot \frac{1}{x} \right)$$

$$\therefore \frac{dy}{dx} = y \left[3 \ln(\ln(x)) + \frac{3}{\ln x} \right]$$

$$= (\ln x)^{3x} \cdot \left[3 \ln(\ln(x)) + \frac{3}{\ln x} \right]$$

(b) Consider $y = x^{\cos x}$. Take: $\ln y = \ln[x^{\cos x}]$

$\Leftrightarrow \ln y = \cos x \cdot \ln x$

Take $\frac{d}{dx}$ of both sides: $\frac{d}{dx}[\ln y] = \frac{d}{dx}[\cos x \cdot \ln x]$

$$\underbrace{\frac{1}{y} \frac{dy}{dx}}_{\text{chain rule}} = -\sin x \cdot \ln x + \cos x \cdot \frac{1}{x}$$

$$\therefore \frac{dy}{dx} = y \left[-\sin x \ln x + \frac{\cos x}{x} \right]$$

OR
$$\frac{dy}{dx} = x^{\cos x} \cdot \left[-\sin x \ln x + \frac{\cos x}{x} \right]$$

(c) $y = (\cos x)^x \Leftrightarrow \ln y = x \cdot \ln(\cos x)$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = 1 \cdot \ln(\cos x) + x \cdot \frac{1}{\cos x} (-\sin x)$$

$$\Rightarrow \frac{dy}{dx} = (\cos x)^x \cdot [\ln(\cos x) - x \tan x]$$

$$y = \frac{e^{2x} \cdot (9x-2)^3}{\sqrt[4]{(x^2+1)(3x^3-7)}}$$

If we try to do the derivative with the product rule, it will be complicated.

Let's take "ln" of both sides:

$$\Leftrightarrow \ln y = \ln \left[\frac{e^{2x} (9x-2)^3}{\sqrt[4]{(x^2+1)(3x^3-7)}} \right]$$

$$\Leftrightarrow \ln y = \ln [e^{2x} (9x-2)^3] - \ln \left[(x^2+1)(3x^3-7) \right]^{\frac{1}{4}}$$

$$\Leftrightarrow \ln y = \ln(e^{2x}) + \ln[(9x-2)^3] - \frac{1}{4} \left[\ln[(x^2+1)(3x^3-7)] \right]$$

$$\Leftrightarrow \ln y = 2x + 3 \ln(9x-2) - \frac{1}{4} \ln(x^2+1) - \frac{1}{4} \ln(3x^3-7)$$

Derivatives of Logarithmic Functions
Logarithmic Differentiation

Theory Examples

Example 6: (Neuhauser, Problem # 77, p. 204)

Use logarithmic differentiation to find the first derivative of the function

$$y = \frac{e^{2x}(9x-2)^3}{\sqrt[4]{(x^2+1)(3x^3-7)}}$$

Hence, after expanding, we have that

$$y = \frac{e^{2x} \cdot (9x-2)^3}{\sqrt[4]{(x^2+1)(3x^3-7)}} \quad (\Leftrightarrow)$$

$$\ln y = 2x + 3 \ln(9x-2) - \frac{1}{4} \ln(x^2+1) - \frac{1}{4} \ln(3x^3-7)$$

Take the derivative w.r.t x of both sides:

$$\frac{1}{y} \cdot \frac{dy}{dx} = 2 + 3 \frac{1}{9x-2} \cdot (9) - \frac{1}{4} \frac{1}{x^2+1} \cdot (2x) - \frac{1}{4} \frac{1}{3x^3-7} \cdot (9x^2)$$

hence $\frac{dy}{dx} = y \left[2 + \frac{27}{9x-2} - \frac{x}{2(x^2+1)} - \frac{9x^2}{4(3x^3-7)} \right]$

or $\frac{dy}{dx} = \frac{e^{2x}(9x-2)^3}{\sqrt[4]{(x^2+1)(3x^3-7)}} \cdot \left[2 + \frac{27}{9x-2} - \frac{x}{2(x^2+1)} - \frac{9x^2}{4(3x^3-7)} \right]$

Power Rule (General Form)

Theorem

Let $f(x) = x^r$, where r is any real number. Then

$$\frac{d}{dx} x^r = r x^{r-1}$$

Proof: We set $y = x^r$ and use logarithmic differentiation to obtain

$$\begin{aligned}\frac{d}{dx} [\ln y] &= \frac{d}{dx} [\ln x^r] \\ \frac{1}{y} \frac{dy}{dx} &= \frac{d}{dx} [r \ln x] \\ \frac{1}{y} \frac{dy}{dx} &= r \frac{1}{x}\end{aligned}$$

Solving for dy/dx yields

$$\frac{dy}{dx} = r \frac{1}{x} y = r \frac{1}{x} x^r = r x^{r-1}$$