

Math 213 - Partial Derivatives

Peter A. Perry

University of Kentucky

September 26, 2018

Homework

- Re-read section 14.3
- Start working on practice problems in section 14.3, 15-31 (odd), 43, 47, 49, 51, 52, 53, 55, 63-69 (odd), 75, 77
- Be ready to work in recitation tomorrow on section 14.3
- Read section 14.4 for Wednesday's lecture


Unit II: Differential Calculus of Several Variables

- Lecture 13 Functions of Several Variables
- Lecture 14 Limits and Continuity
- Lecture 15 **Partial Derivatives**
- Lecture 16 Tangent Planes and Linear Approximation, I
- Lecture 17 Tangent Planes and Linear Approximation, II
- Lecture 18 The Chain Rule
- Lecture 19 Directional Derivatives and the Gradient
- Lecture 20 Maximum and Minimum Values, I
- Lecture 21 Maximum and Minimum Values, II
- Lecture 22 Lagrange Multipliers
- Lecture 23 Review for Exam 2


Goals of the Day

- Learn how to compute partial derivatives and know various different notations for them
- Understand the geometric interpretation of partial derivatives
- Know how to compute higher partial derivatives
- Understand their connection with partial differential equations

Derivatives - One Variable



Derivatives - One Variable




The derivative of f at a is the limit

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

if it exists.

Derivatives - One Variable




The derivative of f at a is the limit

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

if it exists.

$f'(a)$ is the slope of the tangent line to the graph of f at the point $(a, f(a))$.

Derivatives - One Variable



The derivative of f at a is the limit

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

if it exists.

$f'(a)$ is the slope of the tangent line to the graph of f at the point $(a, f(a))$.

$f'(a)$ is also the instantaneous rate of change of $y = f(x)$ at $x = a$

Partial Derivatives - Two Variables

A function of two variables has two very natural rates of change:

- The rate of change of $z = f(x, y)$ with respect to x when y is fixed
- The rate of change of $z = f(x, y)$ when respect to y when x is fixed

The first of these is called the *partial derivative of f with respect to x* , denoted $\partial f / \partial x$ or f_x


$$f_x(a, b) = \lim_{h \rightarrow 0} \frac{f(a + h, b) - f(a, b)}{h}$$

the second is called the *partial derivative of f with respect to y* , denoted $\partial f / \partial y$ or f_y


$$f_y(a, b) = \lim_{h \rightarrow 0} \frac{f(a, b + h) - f(a, b)}{h}$$

Geometric Interpretation

Given a function $f(x, y) \dots$



Geometric Interpretation




Given a function $f(x, y) \dots$

Compute $f_x(a, b)$ by setting $y = b$ and varying x :

$$f_x(a, b) = \lim_{h \rightarrow 0} \frac{f(a+h, b) - f(a, b)}{h}$$

Geometric Interpretation



Given a function $f(x, y) \dots$

Compute $f_x(a, b)$ by setting $y = b$ and varying x :

$$f_x(a, b) = \lim_{h \rightarrow 0} \frac{f(a+h, b) - f(a, b)}{h}$$

Compute $f_y(a, b)$ by setting $x = a$ and varying y :

$$f_y(a, b) = \lim_{h \rightarrow 0} \frac{f(a, b+h) - f(a, b)}{h}$$

Partial Derivatives

Rules for Finding Partial Derivatives of $z = f(x, y)$

1. To find f_x , regard y as a constant and differentiate $f(x, y)$ with respect to x
2. To find f_y , regard x as a constant and differentiate $f(x, y)$ with respect to y

Find both partial derivatives of the following functions:

1. $f(x, y) = x^4 + 5xy^3$

2. $f(x, t) = t^2 e^{-x}$

3. $g(u, v) = (u^2 + v^2)^3$

4. $f(x, y) = \sin(xy)$

5. $f(\text{George}, \text{Fran}) = (\text{George})^5 + (\text{Fran})^3$

More Partial Derivatives

Sometimes it's useful to remember that, to compute a partial derivative like $f_x(x, 1)$, you can set $y = 1$ before you start computing.

Find the following partial derivatives.

1. $f_x(x, 1)$ if $f(x, y) = x^{y^y} \sin(x)$
2. $f_y(3, y)$ if $f(x, y) = (x - 3) \sin(\cos(\log(y))) + xy$

Higher Partials

We can compute higher-order partial derivatives just by repeating operations. We'll find out what these partials actually mean later on!

Example Find the second partial derivatives of $f(x, y) = x^2y^2$

$$\frac{\partial f}{\partial x} = f_x(x, y) = 2xy^2, \quad \frac{\partial f}{\partial y} = 2x^2y$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 f}{\partial x \partial y} =$$

$$\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial^2 f}{\partial y^2} =$$

Notations:

$$\frac{\partial^2 f}{\partial y \partial x} = f_{xy} = (f_x)_y, \quad \frac{\partial^2 f}{\partial x \partial y} = f_{yx} = (f_y)_x$$

Clairaut's Theorem

Suppose f is defined on a disk D that contains the point (a, b) . If the functions f_{xy} and f_{yx} are both continuous on D , then

$$f_{xy}(a, b) = f_{yx}(a, b)$$

Check Clairaut's theorem for the function $f(x, y) = x^3y^2 - \sin(xy)$

Implicit Differentiation

You can find partial derivatives by implicit differentiation.

1. Find $\partial z/\partial x$ and $\partial z/\partial y$ if $x^2 + y^2 + z^2 = 1$
2. Find $\partial z/\partial x$ and $\partial z/\partial y$ if $e^z = xyz$

Partial Differential Equations

Partial Differential Equations describe many physical phenomena. The unknown function is a function of two or more variables.

The *wave equation* for $u(x, t)$, a function which, for each t gives a 'snapshot' of a one-dimensional travelling wave:

$$\frac{\partial^2 u}{\partial t^2} = \frac{1}{c^2} \frac{\partial^2 u}{\partial x^2}$$


The *heat equation* for $u(x, y, t)$, the temperature of a thin sheet at position (x, y) at time t :

$$\frac{\partial u}{\partial t}(x, y, t) = K \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) u(x, y, t)$$

Laplace's Equation for the electrostatic potential of a charge distribution ρ :

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) u(x, y, z) = 4\pi\rho(x, y, z)$$

The Wave Equation




$u(x, t)$ gives the height of a wave moving down a channel as a function of distance x and time t

For each fixed t , we get a “snapshot” of the wave

For each fixed x , we get the height of the wave, at that point, as a function of time

The Heat Equation

$$u(x, t) = (4\pi t)^{-1/2} e^{-x^2/4t}$$



For each t we get a “snapshot” of the distribution of heat—at first heat concentrates near $x = 0$, but then diffuses and cools as time moves forward