

Math 213 - Line Integrals I

Peter A. Perry

University of Kentucky

November 16, 2018

Homework

- Re-Read Section 16.2 for Monday
- Work on Stewart problems for 16.2: 1-21 (odd), 33-41 (odd), 49, 50
- Begin Webwork D1

Unit IV: Vector Calculus

Lecture 35	Vector Fields
Lecture 36	Line Integrals
Lecture 37	Line Integrals
Lecture 38	Fundamental Theorem
Lecture 39	Green's Theorem
Lecture 40	Curl and Divergence

Goals of the Day

- Know how to compute line integrals of a scalar function in the plane
- Know how to compute line integrals of a scalar function in space

Preview: Line Integrals



Our next topic will be integrals of *scalar functions* and *vector functions* over curves in the plane and in space. If C is a curve in the plane or in space, we'll learn how to compute:

- $\int_C f(x, y) ds$, the integral of a scalar function over a plane curve C
- $\int_C f(x, y, z) ds$, the integral of a scalar function over a space curve C
- $\int_C \mathbf{F} \cdot d\mathbf{r}$, the integral of a vector function $\mathbf{F}(x, y)$ over a plane curve C
- $\int_C \mathbf{F} \cdot d\mathbf{r}$, the integral of a vector function $\mathbf{F}(x, y, z)$ over a space curve C

In all cases, we'll reduce these to Calculus I and II type integrals by parameterizing the curve C . We'll also learn how to compute integrals like

- $\int_C f(x, y) dx$
- $\int_C f(x, y) dy$

Parameterizing Paths



Parameterize the following paths:

1. The first planar path shown on the left
2. The second planar path shown on the left
3. The path connecting $(0, 0, 0)$ to $(1, 0, 1)$
4. The path connecting $(1, 0, 1)$ to $(1, 2, 0)$

The Integral of a Scalar Function over a Plane Curve

If C is a plane curve, the **line integral of f along C** is

$$\int_C f(x, y) ds = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*, y_i^*) \Delta s_i$$

where we approximate the curve by n line segments of length Δs_i

As a practical matter, if C is parameterized by $(x(t), y(t))$ for $a \leq t \leq b$,

$$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

so

$$\int_C f(x, y) ds = \int_a^b f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

The Integral of a Scalar Function over a Plane Curve

if C is parameterized by $(x(t), y(t))$ for $a \leq t \leq b$, then

$$\int_C f(x, y) ds = \int_a^b f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

1. Find $\int_C (x/y) ds$ if C is the curve $x = t^2$, $y = 2t$ for $0 \leq t \leq 3$
2. Find $\int_C xy^4 ds$ if C is the right half of the circle $x^2 + y^2 = 16$

Line Integrals over Piecewise Smooth Curves



A curve C is *piecewise smooth* if it is a union of smooth curves C_1, \dots, C_n . Some examples are shown at left.

If C consists of several smooth components, then

$$\int_C f(x, y) ds = \sum_{i=1}^n \int_{C_i} f(x, y) ds$$

Notice that each of these curves has an *orientation* that determines how the curve is parameterized—the parameterization should “follow the arrows.”

1. Find $\int_C xy ds$ if C is the first curve shown at left.



Another Kind of Line Integral

For later use, we'll also need the line integral of f with respect to x and the line integral of f with respect to y :

$$\int_C f(x, y) dx = \int_a^b f(x(t), y(t))x'(t) dt$$

$$\int_C f(x, y) dy = \int_a^b f(x(t), y(t))y'(t) dt$$

1. Find $\int_C e^x dx$ if C is the arc of the curve $x = y^3$ from $(-1, -1)$ to $(1, 1)$
2. Find $\int_C x^2 dx + y^2 dy$ if C is the arc of the circle $x^2 + y^2 = 4$ from $(2, 0)$ to $(0, 2)$ followed by the line segment from $(0, 2)$ to $(4, 3)$

Summary of Line Integrals in the Plane

If C is a parameterized curve $(x(t), y(t))$ where $a \leq t \leq b$:

$$\int_C f(x, y) dx = \int_a^b f(x(t), y(t))x'(t) dt$$

$$\int_C f(x, y) dy = \int_a^b f(x(t), y(t))y'(t) dt$$


$$\int_C f(x, y) ds = \int_a^b f(x(t), y(t))\sqrt{(x'(t))^2 + y'(t)^2} dt$$

Applications - Center of Mass

A wire of mass m and density $\rho(x, y)$ along a curve C has center of mass

$$\bar{x} = \frac{1}{m} \int_C x \rho(x, y) ds$$

$$\bar{y} = \frac{1}{m} \int_C y \rho(x, y) ds$$



A thin wire has the shape of the first quadrant part of a circle with center at the origin and radius a . If the density of the wire is

$$\rho(x, y) = kxy,$$

find the mass and center of mass of the wire.

Line Integrals in Space

If C is a space curve $(x(t), y(t), z(t))$ where $a \leq t \leq b$, then

$$\int_C f(x, y, z) \, ds = \int_a^b f(x(t), y(t), z(t)) \sqrt{(x'(t))^2 + (y'(t))^2 + (z'(t))^2} \, dt$$

1. Find $\int_C (x^2 + y^2 + z^2) \, ds$ if C is the space curve $(x(t), y(t), z(t)) = (t, \cos 2t, \sin 2t)$ for $0 \leq t \leq 2\pi$

More Line Integrals in Space

Can you guess how to define $\int_C f(x, y, z) dx$, $\int_C f(x, y, z) dy$, and $\int_C f(x, y, z) dz$?

1. Find $\int_C (x + z) dx + \int_C (x + z) dy + \int_C (x + y) dz$ if C consists of the line segments from $(0, 0, 0)$ to $(1, 0, 1)$ and from $(1, 0, 1)$ to $(0, 1, 2)$

