

# Math 213 - Parametric Surfaces

Peter A. Perry

University of Kentucky

April 15, 2019

# Homework

- Homework D2 is due Wednesday
- Work on Stewart problems for 16.6: 1-25 odd, 33, 39-49 odd
- Read section 16.7 for Wednesday, April 17
- Remember that Dr. Perry will be out of the office April 17-19. Your lecturer will be Mr. Shane Clark

## Unit IV: Vector Calculus

- Lecture 36    Curl and Divergence
- Lecture 37    **Parametric Surfaces**
- Lecture 38    Surface Integrals
- Lecture 39    Stokes' Theorem
- Lecture 40    The Divergence Theorem
  
- Lecture 41    Final Review, Part I
- Lecture 42    Final Review, Part II

# Goals of the Day

This lecture is about parametric surfaces. You'll learn:

- How to define and visualize parametric surfaces
- How to find the tangent plane to a parametric surface at a point
- How to compute the surface area of a parametric surface using double integrals

# Parametric Curves and Parametric Surfaces

## Parametric Curve

A parametric curve in  $\mathbb{R}^3$  is given by

$$\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}$$

where  $a \leq t \leq b$

There is *one parameter*, because a curve is a *one-dimensional* object

There are *three component functions*, because the curve lives in *three-dimensional* space.

## Parametric Surface

A parametric surface in  $\mathbb{R}^3$  is given by

$$\mathbf{r}(u, v) = x(u, v)\mathbf{i} + y(u, v)\mathbf{j} + z(u, v)\mathbf{k}$$

where  $(u, v)$  lie in a region  $D$  of the  $uv$  plane.

There are *two parameters*, because a surface is a *two-dimensional* object

There are *three component functions*, because the surface lives in *three-dimensional* space.

# You Are Living on a Parametric Surface



Let  $u$  be your latitude (in radians, for this course)

Let  $v$  be your longitude (in radians)


Let  $R$  be the surface of the Earth

Your position is

$$\mathbf{r}(u, v) = R \cos(v) \cos(u)\mathbf{i} + R \cos(v) \sin(u)\mathbf{j} + R \sin(v)\mathbf{k}$$




## More Parameterized Surfaces: Planes



Find a parametric representation for the plane through  $\langle 1, 0, 1 \rangle$  that contains the vectors  $\langle 2, 0, 1 \rangle$  and  $\langle 0, 2, 0 \rangle$

## More Parameterized Surfaces: Planes




Find a parametric representation for the plane through  $\langle 1, 0, 1 \rangle$  that contains the vectors  $\langle 2, 0, 1 \rangle$  and  $\langle 0, 2, 0 \rangle$

*Solution:* Let  $\mathbf{r}_0 = \langle 1, 0, 1 \rangle$ . Any point in the plane is given by

$$\mathbf{r}(s, t) = \langle 1, 0, 1 \rangle + s\langle 2, 0, 1 \rangle + t\langle 0, 2, 0 \rangle$$

## More Parameterized Surfaces: Planes



Find a parametric representation for the plane through  $\langle 1, 0, 1 \rangle$  that contains the vectors  $\langle 2, 0, 1 \rangle$  and  $\langle 0, 2, 0 \rangle$


*Solution:* Let  $\mathbf{r}_0 = \langle 1, 0, 1 \rangle$ . Any point in the plane is given by

$$\mathbf{r}(s, t) = \langle 1, 0, 1 \rangle + s\langle 2, 0, 1 \rangle + t\langle 0, 2, 0 \rangle$$

Now you try it:

Find a parametric representation for the plane through the point  $(0, -1, 5)$  that contains the vectors  $\langle 2, 1, 4 \rangle$  and  $\langle -3, 2, 5 \rangle$ .


## More Parameterized Surfaces: The Cylinder




$$\mathbf{r}(u, v) = r \cos(u) \mathbf{i} + r \sin(u) \mathbf{j} + v \mathbf{k}$$

$$D = \{(u, v) : 0 \leq u \leq 2\pi, 0 \leq v \leq h\}$$

parameterizes a cylinder of radius  $r$  and height  $h$




# More Parameterized Surfaces: The Cylinder



$$\mathbf{r}(u, v) = r \cos(u)\mathbf{i} + r \sin(u)\mathbf{j} + v\mathbf{k}$$


$$D = \{(u, v) : 0 \leq u \leq 2\pi, 0 \leq v \leq h\}$$

parameterizes a cylinder of radius  $r$  and height  $h$



If we fix  $v$  and vary  $u$  over the cylinder, we trace out a **circle**


# More Parameterized Surfaces: The Cylinder



$$\mathbf{r}(u, v) = r \cos(u)\mathbf{i} + r \sin(u)\mathbf{j} + v\mathbf{k}$$

$$D = \{(u, v) : 0 \leq u \leq 2\pi, 0 \leq v \leq h\}$$


parameterizes a cylinder of radius  $r$  and height  $h$



If we fix  $v$  and vary  $u$  over the cylinder, we trace out a **circle**

If we fix  $u$  and vary  $v$ , we trace out a **vertical line**


# More Parameterized Surfaces: The Cylinder



$$\mathbf{r}(u, v) = r \cos(u)\mathbf{i} + r \sin(u)\mathbf{j} + v\mathbf{k}$$

$$D = \{(u, v) : 0 \leq u \leq 2\pi, 0 \leq v \leq h\}$$

parameterizes a cylinder of radius  $r$  and height  $h$



If we fix  $v$  and vary  $u$  over the cylinder, we trace out a **circle**


If we fix  $u$  and vary  $v$ , we trace out a **vertical line**

Each of these curves has a *tangent vector*:

$$\mathbf{r}_u(u, v) = -r \sin(u)\mathbf{i} + r \cos(u)\mathbf{j}$$

$$\mathbf{r}_v(u, v) = \mathbf{k}$$

# More Parameterized Surfaces: The Cylinder



$$\mathbf{r}(u, v) = r \cos(u)\mathbf{i} + r \sin(u)\mathbf{j} + v\mathbf{k}$$

$$D = \{(u, v) : 0 \leq u \leq 2\pi, 0 \leq v \leq h\}$$


parameterizes a cylinder of radius  $r$  and height  $h$

The two tangent vectors

$$\mathbf{r}_u(u, v) = -r \sin(u)\mathbf{i} + r \cos(u)\mathbf{j}$$

$$\mathbf{r}_v(u, v) = \mathbf{k}$$

span the *tangent plane* to the cylinder at the given point



# The Tangent Vectors $\mathbf{r}_u$ and $\mathbf{r}_v$

Suppose

$$\begin{aligned}\mathbf{r}(u, v) &= x(u, v)\mathbf{i} + y(u, v)\mathbf{j} + z(u, v)\mathbf{k} \\ (u, v) &\in D\end{aligned}$$

is a parameterized surface.

At a point  $P_0 = \mathbf{r}(u_0, v_0)$ , the vectors

$$\mathbf{r}_u(u_0, v_0) = \frac{\partial x}{\partial u}(u_0, v_0)\mathbf{i} + \frac{\partial y}{\partial u}(u_0, v_0)\mathbf{j} + \frac{\partial z}{\partial u}(u_0, v_0)\mathbf{k}$$

$$\mathbf{r}_v(u_0, v_0) = \frac{\partial x}{\partial v}(u_0, v_0)\mathbf{i} + \frac{\partial y}{\partial v}(u_0, v_0)\mathbf{j} + \frac{\partial z}{\partial v}(u_0, v_0)\mathbf{k}$$

are *both* tangent to the surface.

# The Tangent Plane

$$\mathbf{r}_u(u_0, v_0) = \frac{\partial x}{\partial u}(u_0, v_0)\mathbf{i} + \frac{\partial y}{\partial u}(u_0, v_0)\mathbf{j} + \frac{\partial z}{\partial u}(u_0, v_0)\mathbf{k}$$

$$\mathbf{r}_v(u_0, v_0) = \frac{\partial x}{\partial v}(u_0, v_0)\mathbf{i} + \frac{\partial y}{\partial v}(u_0, v_0)\mathbf{j} + \frac{\partial z}{\partial v}(u_0, v_0)\mathbf{k}$$


The *tangent plane* to a parameterized surface at  $P_0 = \mathbf{r}(u_0, v_0)$  is the plane passing through  $P_0$  and perpendicular to  $\mathbf{r}_u(u_0, v_0) \times \mathbf{r}_v(u_0, v_0)$ .

Find the equation of the tangent plane to the surface


$$\mathbf{r}(u, v) = u^2\mathbf{i} + 2u \sin v\mathbf{j} + u \cos v\mathbf{k}$$

at  $u = 1, v = 0$ .

# The Tangent Plane



# The Tangent Plane




$$\mathbf{r}(u, v) = \langle u^2, 2u \sin v, u \cos v \rangle$$

$$\mathbf{r}_u(u, v) = \langle 2u, 2 \sin v, \cos v \rangle$$

$$\mathbf{r}_v(u, v) = \langle 0, 2u \cos v, -u \sin v \rangle$$

# The Tangent Plane



$$\mathbf{r}(u, v) = \langle u^2, 2u \sin v, u \cos v \rangle$$

$$\mathbf{r}_u(u, v) = \langle 2u, 2 \sin v, \cos v \rangle$$


$$\mathbf{r}_v(u, v) = \langle 0, 2u \cos v, -u \sin v \rangle$$

$$\mathbf{r}(1, 0) = \langle 1, 0, 1 \rangle$$

$$\mathbf{r}_u(1, 0) = \langle 2, 0, 1 \rangle$$

$$\mathbf{r}_v(1, 0) = \langle 0, 2, 0 \rangle$$

# The Tangent Plane



The normal to the plane is

$$\mathbf{r}_u \times \mathbf{r}_v = \langle -1, 0, 2 \rangle$$

$$\mathbf{r}(u, v) = \langle u^2, 2u \sin v, u \cos v \rangle$$

$$\mathbf{r}_u(u, v) = \langle 2u, 2 \sin v, \cos v \rangle$$


$$\mathbf{r}_v(u, v) = \langle 0, 2u \cos v, -u \sin v \rangle$$

$$\mathbf{r}(1, 0) = \langle 1, 0, 1 \rangle$$

$$\mathbf{r}_u(1, 0) = \langle 2, 0, 1 \rangle$$

$$\mathbf{r}_v(1, 0) = \langle 0, 2, 0 \rangle$$

# The Tangent Plane



$$\mathbf{r}(u, v) = \langle u^2, 2u \sin v, u \cos v \rangle$$

$$\mathbf{r}_u(u, v) = \langle 2u, 2 \sin v, \cos v \rangle$$

$$\mathbf{r}_v(u, v) = \langle 0, 2u \cos v, -u \sin v \rangle$$

$$\mathbf{r}(1, 0) = \langle 1, 0, 1 \rangle$$

$$\mathbf{r}_u(1, 0) = \langle 2, 0, 1 \rangle$$

$$\mathbf{r}_v(1, 0) = \langle 0, 2, 0 \rangle$$

The normal to the plane is

$$\mathbf{r}_u \times \mathbf{r}_v = \langle -1, 0, 2 \rangle$$


so the equation of the plane is

$$(-1)(x - 1) + 2(z - 1) = 0$$

The tangent plane to the surface at  $(1, 0, 1)$  is parameterized by

$$\langle 1 + 2s, 2t, 1 + s \rangle$$

# The Sphere Revisited




$$\begin{aligned}\mathbf{r}(u, v) = & \sin(v) \cos(u) \mathbf{i} \\ & + \sin(v) \sin(u) \mathbf{j} \\ & + \cos(v) \mathbf{k}\end{aligned}$$

$$0 \leq u \leq 2\pi, \quad 0 \leq v \leq \pi$$

$$\begin{aligned}\mathbf{r}_u = & -\sin(v) \sin(u) \mathbf{i} + \sin(v) \cos(u) \mathbf{j} \\ \mathbf{r}_v = & \cos(v) \cos(u) \mathbf{i} + \cos(v) \sin(u) \mathbf{j} \\ & - \sin(v) \mathbf{k}\end{aligned}$$

Find the tangent plane to the sphere at  $(u, v) = (\pi/4, \pi/4)$

# The Sphere Revisited



$$\begin{aligned}\mathbf{r}(u, v) &= \sin(v) \cos(u) \mathbf{i} \\ &\quad + \sin(v) \sin(u) \mathbf{j} \\ &\quad + \cos(v) \mathbf{k}\end{aligned}$$

$$0 \leq u \leq 2\pi, \quad 0 \leq v \leq \pi$$

$$\begin{aligned}\mathbf{r}_u &= -\sin(v) \sin(u) \mathbf{i} + \sin(v) \cos(u) \mathbf{j} \\ \mathbf{r}_v &= \cos(v) \cos(u) \mathbf{i} + \cos(v) \sin(u) \mathbf{j} \\ &\quad - \sin(v) \mathbf{k}\end{aligned}$$


Find the tangent plane to the sphere at  $(u, v) = (\pi/4, \pi/4)$

$$\mathbf{r}(\pi/4, \pi/4) = \frac{1}{2} \mathbf{i} + \frac{1}{2} \mathbf{j} + \frac{\sqrt{2}}{2} \mathbf{k}$$

$$\mathbf{r}_u(\pi/4, \pi/4) = -\frac{1}{2} \mathbf{i} + \frac{1}{2} \mathbf{j}$$

$$\mathbf{r}_v(\pi/4, \pi/4) = \frac{1}{2} \mathbf{i} + \frac{1}{2} \mathbf{j} - \frac{\sqrt{2}}{2} \mathbf{k}$$

# The Sphere Revisited



$$\begin{aligned}\mathbf{r}(u, v) &= \sin(v) \cos(u) \mathbf{i} \\ &\quad + \sin(v) \sin(u) \mathbf{j} \\ &\quad + \cos(v) \mathbf{k}\end{aligned}$$

$$0 \leq u \leq 2\pi, \quad 0 \leq v \leq \pi$$

$$\begin{aligned}\mathbf{r}_u &= -\sin(v) \sin(u) \mathbf{i} + \sin(v) \cos(u) \mathbf{j} \\ \mathbf{r}_v &= \cos(v) \cos(u) \mathbf{i} + \cos(v) \sin(u) \mathbf{j} \\ &\quad - \sin(v) \mathbf{k}\end{aligned}$$

Find the tangent plane to the sphere at  $(u, v) = (\pi/4, \pi/4)$

$$\mathbf{r}(\pi/4, \pi/4) = \frac{1}{2} \mathbf{i} + \frac{1}{2} \mathbf{j} + \frac{\sqrt{2}}{2} \mathbf{k}$$

$$\mathbf{r}_u(\pi/4, \pi/4) = -\frac{1}{2} \mathbf{i} + \frac{1}{2} \mathbf{j}$$

$$\mathbf{r}_v(\pi/4, \pi/4) = \frac{1}{2} \mathbf{i} + \frac{1}{2} \mathbf{j} - \frac{\sqrt{2}}{2} \mathbf{k}$$

$$\begin{aligned}\mathbf{n} = \mathbf{r}_u \times \mathbf{r}_v &= -\frac{1}{2} \left( \frac{1}{\sqrt{2}} \mathbf{i} + \frac{1}{\sqrt{2}} \mathbf{j} + \mathbf{k} \right) \\ 0 &= \frac{1}{\sqrt{2}} \left( x - \frac{1}{2} \right) + \frac{1}{\sqrt{2}} \left( y - \frac{1}{2} \right) \\ &\quad + \left( z - \frac{\sqrt{2}}{2} \right)\end{aligned}$$

# Sneak Preview

## Parametric Curves - Arc Length

$$\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}$$

$$\mathbf{r}'(t) = x'(t)\mathbf{i} + y'(t)\mathbf{j} + z'(t)\mathbf{k}$$

$$|\mathbf{r}'(t)| = \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2}$$

$$ds = |\mathbf{r}'(t)| dt$$

$$L = \int_a^b |\mathbf{r}'(t)| dt$$

## Parametric Surfaces - Area

$$\mathbf{r}(u, v) = x(u, v)\mathbf{i} + y(u, v)\mathbf{j} + z(u, v)\mathbf{k}$$



$$\mathbf{r}_u(u, v) = \frac{\partial \mathbf{r}}{\partial u}(u, v)$$

$$\mathbf{r}_v(u, v) = \frac{\partial \mathbf{r}}{\partial v}(u, v)$$

$$dA = |\mathbf{r}_u \times \mathbf{r}_v| du dv$$

$$S = \iint_D |\mathbf{r}_u \times \mathbf{r}_v| du dv$$

# Surface Area



Find the area  $\Delta A$  of a small patch of surface

The map  $(u, v) \mapsto \mathbf{r}(u, v)$  takes the square to a parallelogram with sides  $\mathbf{r}_u \Delta u$  and  $\mathbf{r}_v \Delta v$

The area of the parallelogram is

$$|\mathbf{r}_u \Delta u \times \mathbf{r}_v \Delta v| = |\mathbf{r}_u \times \mathbf{r}_v| \Delta u \Delta v$$


The area of the surface is approximately

$$A \approx \sum_{i,j} |\mathbf{r}_u(u_i, v_i) \times \mathbf{r}_v(u_i, v_i)| \Delta u \Delta v$$

and exactly

$$\iint_D |\mathbf{r}_u(u, v) \times \mathbf{r}_v(u, v)| \, du \, dv$$

# Surface Area of a Sphere



$$\begin{aligned}\mathbf{r}(u, v) &= a \sin(v) \cos(u) \mathbf{i} \\ &\quad + a \sin(v) \sin(u) \mathbf{j} \\ &\quad + a \cos(v) \mathbf{k}\end{aligned}$$

$$0 \leq u \leq 2\pi, \quad 0 \leq v \leq \pi$$

$$\begin{aligned}\mathbf{r}_u &= -a \sin(v) \sin(u) \mathbf{i} + a \sin(v) \cos(u) \mathbf{j} \\ \mathbf{r}_v &= a \cos(v) \cos(u) \mathbf{i} + a \cos(v) \sin(u) \mathbf{j} \\ &\quad - \sin(v) \mathbf{k}\end{aligned}$$

$$\begin{aligned}\mathbf{r}_u \times \mathbf{r}_v &= a^2 \sin^2(v) \cos(u) \mathbf{i} + a^2 \sin^2(v) \sin(u) \mathbf{j} - a^2 \cos(v) \sin(v) \mathbf{k} \\ |\mathbf{r}_u \times \mathbf{r}_v| &= a^2 \sin^2(v)\end{aligned}$$

Hence

$$S = \int_0^\pi \int_0^{2\pi} a^2 \sin^2 v \, du \, dv = 4\pi a^2$$

## Surfaces Area of a Graph

The graph of a function  $z = f(x, y)$  is also a parameterized surface:

$$\mathbf{r}(x, y) = x\mathbf{i} + y\mathbf{j} + f(x, y)\mathbf{k}$$

$$\mathbf{r}_x(x, y) = \mathbf{i} + \frac{\partial f}{\partial x}\mathbf{k}$$

$$\mathbf{r}_y(x, y) = \mathbf{j} + \frac{\partial f}{\partial y}\mathbf{k}$$

$$\mathbf{r}_x \times \mathbf{r}_y = -\frac{\partial f}{\partial x}\mathbf{i} + -\frac{\partial f}{\partial y}\mathbf{j} + \mathbf{k}$$

$$|\mathbf{r}_x \times \mathbf{r}_y| = \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2}$$

Hence, the surface area of the graph over a domain  $D$  in the  $xy$  plane is

$$S = \iint_D \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dA$$

# Surface Area of a Graph

The surface area of the graph over a domain  $D$  in the  $xy$  plane is

$$S = \iint_D \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dA$$

Find the area under the graph of  $z = x^2 + y^2$  that lies over the cylinder  $x^2 + y^2 = 4$

# Curves and Surfaces

## Curves

Parameterization

$$\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}$$

Tangent

$$\mathbf{r}'(t) = x'(t)\mathbf{i} + y'(t)\mathbf{j} + z'(t)\mathbf{k}$$

Tangent line at  $t = a$

$$\mathbf{L}(s) = \mathbf{r}(a) + s\mathbf{r}'(a)$$

Arc length differential

$$ds = \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2} dt$$

## Surfaces

Parameterization

$$\mathbf{r}(u, v) = x(u, v)\mathbf{i} + y(u, v)\mathbf{j} + z(u, v)\mathbf{k}$$

Tangents

$$\mathbf{r}_u(u, v) = \frac{\partial}{\partial u} \mathbf{r}(u, v)$$

$$\mathbf{r}_v(u, v) = \frac{\partial}{\partial v} \mathbf{r}(u, v)$$

Normal

$$\mathbf{n} = \mathbf{r}_u \times \mathbf{r}_v$$

Area Differential

$$dA = |\mathbf{r}_u \times \mathbf{r}_v| du dv$$