1 Lecture 04: The tangent and velocity problem, informal treatment of limits

- 1. Estimating the slope of a tangent line.
- 2. Instantaneous velocity
- 3. A limit that does not exist, one-sided limits

1.1 The tangent problem

It is a well-known fact from geometry that the tangent to a circle is line that is perpendicular to the radius. One of the fundamental ideas of calculus can be used to help find tangent lines to many curves. Suppose we have a function y = f(x) and want to define a tangent line to the graph.

We know several ways to write the equation of a line. The approach that is most useful in this problem is the "point-slope form of a line", the line through (x_0, y_0) with slope m is

$$y - y_0 = m(x - x_0).$$

If we want to find the tangent line to f(x) at $x = x_0$, then we know the line should pass through $(x_0, f(x_0))$. The only mystery is what is the appropriate value for the slope. The technique we will use is to pick a point (x, f(x)) that is near $(x_0, f(x_0))$ and compute the slope of the line joining (x, f(x)) and $(x_0, f(x_0))$. This line which meets the graph of f at least twice will be called a *secant line*. We try various values of x that are close to f(x) and hope that we can guess the value of the slope when the distance between x and x_0 vanishes.

Example. Consider the function $f(x) = e^x$. What is the slope of the tangent line to the graph of f at x = 0?

Solution. If x is a point near 0, the slope of the line joining (0, f(0)) to (x, f(x)) is

$$m = \frac{f(x) - f(0)}{x - 0} = \frac{e^x - 1}{x}.$$

If we compute this for several values of x, we obtain

Value of x	Slope of secant line
0.1	1.105
0.01	1.005
-0.002	0.999
$1/\pi^3$	1.0163

A moment's reflection might lead us to guess that the slope is 1. Thus the tangent line to the graph of e^x is y = x + 1. The graph below suggests that this correct.



1.2 Limits

The process we used in the previous section to find the tangent line is of fundamental importance. We give an informal definition.

Definition. Suppose f(x) is a function that is defined an interval containing a number a, except possibly at a. If the values f(x) become close to a number L when we let the distance between x and a approaches 0, then we call L the *limit of* f as x approaches a and write

$$\lim_{x \to a} f(x) = L.$$

In the previous example where we found the slope of the tangent line, we were trying to find:

$$\lim_{x \to 0} \frac{e^x - 1}{x}$$

The function $(e^x - 1)/x$ is defined near x = 0, but not at 0. We are interested in trying to determine the behavior near 0.

1.3 Velocity

To describe the motion of an object moving along a line, for example an object that is thrown straight up in the air, we use a position function. Suppose, we want to describe the velocity of the object. We recall the fundamental relation

distance = rate \times time.

In our case, we want to compute a velocity at $t = t_0$. Our solution will look similar (identical) to our solution of the tangent problem.

We fix a small interval $[t_0, t]$ with one endpoint t_0 and a nearby time t. The distance travelled in this interval is $h(t) - h(t_0)$ and the time it takes to travel this distance is $t - t_0$. Thus the *average velocity* on this interval is

$$\frac{h(t) - h(t_0)}{t - t_0}.$$

If we let t approach t_0 , and the average velocities cluster around one number, then we call this number the *instantaneous velocity* at t_0 . This instantaneous velocity is given by the limit

$$\lim_{t \to t_0} \frac{p(t) - p(t_0)}{t - t_0}.$$

Example. We give a simple numerical example. A ball thrown up in the air and its height in meters at time t seconds is given by $p(t) = -5t^2 + 20t$. Find the average velocity on the interval (3, 3 + h) and guess the instantaneous velocity at 3.

Solution. On the interval $3 \le t \le 3 + h$, the change in position is p(3 + h) - p(3) meters and the time interval is of length 3 + h - 3 = h seconds. Thus the average velocity is

$$\frac{p(3+h) - p(3)}{h}$$

We could again try numerical values of h, but this problem we can simplify algebraically:

$$\frac{p(3+h) - p(3)}{h} = \frac{-5(3+h)^2 + 60 + 20h - 15}{h} \tag{1}$$

$$=\frac{-10h+5h^2}{l}$$
 (2)

$$= -10 + 5h \tag{3}$$

Using the last expression it is easy to see that this expression approaches -10 as h gets close to zero.

Using our new notation, we would write

$$\lim_{h \to 0} \frac{p(3+h) - p(3)}{h} = -10.$$

and that the instantaneous velocity at 3 is -10 meters/second.

Exercise. Find the tangent line to the graph of $f(x) = \frac{1}{x}$ at x = 2.

Exercise. A ball is thrown so that its height at time t is

$$h(t) = -5t^2 + 20t$$

meters after t seconds. Find the instantaneous velocity at time t = 2 seconds. What are the units for this velocity?

Find the instantaneous velocity at an arbitrary time t = a.

1.4 One sided limits

Example. Can you find the tangent line to f(x) = |x - 1| at x = 1?

Solution. In this case, we would want to consider the slope

$$\frac{f(x) - f(1)}{x - 1} = \frac{|x - 1|}{x - 1}.$$

Let us call this a new function g(x) = |x - 1|/(x - 1) and consider the graph of g,



Examining the graph, we see that the function g does not have a limit. When x > 1, the value of g is 1 and when x < 1, the value of g is -1. As a result there is no single value which g approaches when x approaches 1.

Returning to the graph of f, we see that there is corner at x = 1 and there is no clear way to define a single tangent line. The graph includes several lines with touch the graph at one point.



The previous example serves to introduce one-sided limits.

Definition. Suppose f(x) is a function that is defined on an interval (a, b) for some b > a, except possibly at a. If the values f(x) become close to a number L when the distance between x and a approaches 0 and x > a, then we call L the *limit of* f as x approaches a from above and write

I

$$\lim_{x \to a^+} f(x) = L.$$

Definition. Suppose f(x) is a function that is defined on an interval containing (b, a) for some b < a. If the values f(x) become close to a number L when the distance between x and a approaches 0 and x < a, then we call L the *limit of* f as x approaches a from below and write

$$\lim_{x \to a^{-}} f(x) = L.$$

The following theorem gives the relation between one and two-sided limits.

Theorem 4 Suppose that f is a function defined on an open interval containing a, except possibly at a. Then we have $\lim_{x\to a} f(x)$ exists if and only if both of the one-sided limits $\lim_{x\to a^+} f(x)$ and $\lim_{x\to a^-} f(x)$ exist and are equal.

Example. Find the limits from the graph.

1.5 Limits that are infinite

Recall that if $\lim_{x\to a} f(x) = L$, the values of f become arbitrarily close to L, but we may never have f(x) = L. We want to describe the behavior of a function like $f(x) = 1/x^2$ near x = 0. As x small, the reciprocal $1/x^2$ becomes large and positive. We say that $\lim_{x\to 0} \frac{1}{x}^2 = +\infty$. But there is no number ∞ so that f never reaches ∞ .

We try to give a definition of this behaviour.

Definition. We say that the limit of f as x approaches a is $+\infty$ and write

$$\lim_{x \to a} f(x) = \infty$$

if the values of f become arbitrarily large and positive as the distance between x and a approaches 0.

We leave it to the reader to define what it means for a limit to be $-\infty$ and one-sided limits which approach $\pm\infty$.

Example. Discuss the limit $\lim_{x\to -4} x - 8x - 4$.

Solution. If we consider values of x > 4, then x - 4 > 0, but becomes small as x approaches 4. Thus the reciprocal 1/(x - 4) approaches $+\infty$ as x approaches 4 from the left. Also x - 8 < 0 for x near 4. Together we have

$$\lim_{x \to 4^+} \frac{x - 8}{x - 4} = -\infty.$$

Similar reasoning with x < 4, but close to 4 gives that

$$\lim_{x \to 4^{-}} \frac{x - 8}{x - 4} = +\infty.$$

Since the left and right limits are not the same, the limit does not exist and is not $+\infty$ or $-\infty$.