

# 1 Lecture 08: The squeeze theorem

- The squeeze theorem
- The limit of  $\sin(x)/x$
- Related trig limits

## 1.1 The squeeze theorem

*Example.* Is the function  $g$  defined by

$$g(x) = \begin{cases} x^2 \sin(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

continuous?

*Solution.* If  $x \neq 0$ , then  $\sin(1/x)$  is a composition of continuous function and thus  $x^2 \sin(1/x)$  is a product of continuous function and hence continuous.

If  $x = 0$ , we need to have that  $\lim_{x \rightarrow 0} g(x) = g(0) = 0$  in order for  $g$  to satisfy the definition of continuity. Recalling that  $\sin(1/x)$  oscillates between  $-1 \leq x \leq 1$ , we have that

$$-x^2 \leq g(x) \leq x^2$$

and since  $\lim_{x \rightarrow 0} x^2 = \lim_{x \rightarrow 0} -x^2 = 0$ , the theorem below tells us we have  $\lim_{x \rightarrow 0} g(x) = 0$ . ■

**Theorem 1 (The squeeze theorem)** *If  $f$ ,  $g$ , and  $h$  are functions and for all  $x$  in an open interval containing  $c$ , but perhaps not at  $c$ , we have*

$$f(x) \leq g(x) \leq h(x)$$

*and*

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} h(x) = L,$$

*then*

$$\lim_{x \rightarrow c} g(x) = L.$$

We will not give a proof but it should be intuitive that if  $g$  is trapped between two functions that approach the limit  $L$ , then  $g$  also approaches that limit.

## 1.2 The limit of $\sin(x)/x$

We consider the limit

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x}.$$

The quotient rule for limits does not apply since the limit of the denominator is 0. Unlike our previous limits, we cannot simplify to obtain a function where we can use the direct substitution rule or another rule. Instead, we will use the squeeze theorem.


### Theorem 2

$$\lim_{t \rightarrow 0} \frac{\sin(t)}{t}.$$

*Proof.* We start by observing that  $\sin(-t)/(-t) = \sin(t)/t$ , so it suffices to consider  $\lim_{t \rightarrow 0^+} \sin(t)/t$ .

In the figure below, we observe that we have the inequalities

$$\text{Area triangle } OAB \leq \text{Area sector } OAB \leq \text{Area triangle } OAC.$$



We have

$$\begin{aligned} \text{Area triangle } OAB &= \frac{1}{2} \sin(t) \\ \text{Area sector } OAB &= \frac{1}{2} t \\ \text{Area triangle } OAC &= \frac{1}{2} \tan(t) \end{aligned}$$

Thus we have

$$\frac{1}{2} \sin(t) \leq t/2 \leq \frac{1}{2} \tan(t).$$

Since  $t > 0$ , we can rearrange to obtain

$$\cos(t) \leq \frac{\sin(t)}{t} \leq 1$$

and then the squeeze theorem gives that

$$\lim_{t \rightarrow 0} \frac{\sin(t)}{t} = 1.$$

■

### 1.3 Some consequences

Using this limit, we can find several related limits.

The first one will be used in the next chapter.

*Example.* Find the limit

$$\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x}.$$

*Solution.* We note that since the limit of the denominator is zero, we cannot use the quotient rule for limits. However, if we multiply and divide by  $1 + \cos(x)$  and use the identity  $\sin^2(x) + \cos^2(x) = 1$ , we have

$$\frac{1 - \cos(x)}{x} = \frac{(1 - \cos(x))(1 + \cos(x))}{x(1 + \cos(x))} = \frac{\sin^2(x)}{x}.$$

Thus, we may use the rule for a limit of a product,

$$\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x} = \lim_{x \rightarrow 0} \frac{\sin^2(x)}{x} = \lim_{x \rightarrow 0} \sin(x) \lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 0.$$

■

Below are a few more to try

1.  $\lim_{t \rightarrow 0} \frac{\sin(2t)}{t}$
2.  $\lim_{t \rightarrow 0} \frac{\sin(2t)}{\sin(3t)}$
3.  $\lim_{t \rightarrow 0} \frac{1 - \cos(t)}{t^2}$

September 16, 2013