

1 Lecture 14: The product and quotient rule

1.1 Outline

- The product rule
- The reciprocal rule
- The quotient rule.

1.2 The derivative of a product

Theorem 1 Suppose that f and g are two functions which are differentiable at a point x , then fg is differentiable at x and

$$(fg)'(x) = f'(x)g(x) + f(x)g'(x).$$

Proof. The proof depends on rewriting the difference quotient for fg in terms of the difference quotients for f and g . This depends on the trick of adding and subtracting $f(x)g(x+h)$ as follows

$$\begin{aligned}\frac{f(x+h)g(x+h) - f(x)g(x)}{h} &= \frac{f(x+h)g(x+h) - f(x)g(x+h) + f(x)g(x+h) - f(x)g(x)}{h} \\ &= \frac{f(x+h) - f(x)}{h}g(x+h) + f(x)\frac{g(x+h) - g(x)}{h}.\end{aligned}$$

We know that the difference quotients for f and g have a limit as h tends to zero. Since g is differentiable at x , it is continuous and we have

$$\lim_{h \rightarrow 0} g(x+h) = g(x).$$

Thus we may use the rules for sums and products of limits to obtain that

$$\begin{aligned}(fg)'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \lim_{h \rightarrow 0} g(x+h) + f(x) \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &= f'(x)g(x) + f(x)g'(x)\end{aligned}$$


■

One way to understand this rule is to think of a rectangle whose length ℓ and width w are given by $\ell(t) = \ell + mt$ and $w(t) = w + nt$. Then the area will be given by

$$\ell(t)w(t) = (\ell + mt)(w + nt) = \ell w + (mw + \ell n)t + mnt^2.$$

At $t = 0$, the instantaneous rate of change of the area will be the coefficient of t , $mw + \ell n$. Since m is the rate of change of ℓ and n is the rate of change of w , the rate

of change of the product is exactly what we see in the product rule. The picture shows how the increase in area of a rectangle is related to the sidelengths of the rectangle.



Example. Compute the derivative $f(x) = (1 + 2x)^2 e^x$.

Solution. At the moment, we do not know how to differentiate the function $(1 + 2x)^2$. However, if we expand the square, we can write

$$(1 + 2x)^2 e^x = (1 + 4x + 4x^2) e^x.$$

We use the Leibniz notation,

$$\begin{aligned} \frac{d}{dx}((1 + 4x + 4x^2)e^x) &= \left(\frac{d}{dx}(1 + 4x + 4x^2)\right)e^x + (1 + 4x + 4x^2)\frac{d}{dx}e^x \\ &= (4 + 8x)e^x + (1 + 4x + 4x^2)e^x \\ &= (5 + 12x + 4x^2)e^x. \end{aligned}$$

■

Example. Find the derivative of x^5 by writing $x^5 = x^4 \cdot x$ and applying the product rule.

Solution. We write $x^5 = x^4 \cdot x$ and apply the product rule,

$$\frac{d}{dx}x^5 = \frac{d}{dx}(x^4 \cdot x) = x^4 \frac{d}{dx}x + \left(\frac{d}{dx}x^4\right)x.$$

Computing the derivatives gives

$$4x^3 \cdot x + x^4 \cdot 1 = 5x^4.$$

■

1.3 Reciprocals

We find the derivative of a reciprocal or the multiplicative inverse of a function.

Theorem 2 *If g is differentiable at x and $g(x) \neq 0$, then $1/g$ is differentiable at x and we have*

$$\left(\frac{1}{g}\right)'(x) = \frac{-g'(x)}{g(x)^2}.$$

Proof. We write out the difference quotient for $1/g$, obtain a common denominator and simplify to express it in terms of the difference quotient for g ,

$$\begin{aligned}\frac{1}{h}\left(\frac{1}{g(x+h)} - \frac{1}{g(x)}\right) &= \frac{1}{h} \frac{g(x)}{g(x)g(x+h)} - \frac{g(x+h)}{g(x)g(x+h)} \\ &= \frac{-1}{h}(g(x+h) - g(x)) \frac{1}{g(x+h)g(x)}.\end{aligned}$$

Now we may use the limit laws and that $1/g$ is continuous at x to write

$$\begin{aligned}\left(\frac{1}{g}\right)'(x) &= -\lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \cdot \lim_{h \rightarrow 0} \frac{1}{g(x)g(x+h)} \\ &= \frac{-g'(x)}{g(x)^2}.\end{aligned}$$

■

Example. Use this rule to find the derivative

$$\frac{d}{dx} \frac{1}{x^4}.$$

Solution. $\frac{d}{dx} \frac{1}{x^4} = \frac{-4x^3}{(x^4)^2} = \frac{-4}{x^5}.$

■

We can use the reciprocal rule to extend the power rule to negative exponents.

Example. Use the reciprocal rule to find the derivative

$$\frac{d}{dx} x^{-n}, \quad \text{for } n = 1, 2, 3, \dots$$

Solution. $\frac{d}{dx} \frac{1}{x^n} = \frac{-nx^{n-1}}{x^{2n}} = -nx^{-n-1}.$

■

1.4 Quotient rule

Finally we give the quotient rule. Note that it is often simpler to rewrite a quotient as a product and avoid the quotient rule.

Theorem 3 *If f and g are differentiable at x and $g(x) \neq 0$, then f/g is differentiable at x and*

$$\left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}.$$

Proof. We may prove this writing $f/g = f \cdot \frac{1}{g}$ and using the product and the reciprocal rule.

$$\left(\frac{f}{g}\right)'(x) = \left(f \frac{1}{g}\right)'(x) = f'(x) \frac{1}{g(x)} + f(x) \frac{-g'(x)}{g(x)^2}$$

We may simplify this last expression by obtaining a common denominator.

$$f'(x) \frac{1}{g(x)} + f(x) \frac{-g'(x)}{g(x)^2} = f'(x) \frac{g(x)}{g(x)^2} + f(x) \frac{-g'(x)}{g(x)^2} = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}.$$

■

Example. Find the tangent line to

$$f(x) = \frac{x^2 + 3}{x^2 - 3}$$

at $x = 1$.

Solution. The tangent line will pass through the point $(2, f(2)) = (2, 7)$. We need the derivative of f to compute the slope. We use the quotient rule to find the derivative of f ,

$$\begin{aligned} f'(x) &= \frac{\left(\frac{d}{dx}(x^2 + 3)\right)(x^2 - 3) - (x^2 + 3)\frac{d}{dx}(x^2 - 3)}{(x^2 - 3)^2} \\ &= \frac{2x(x^2 - 3) - (x^2 + 3)2x}{(x^2 - 3)^2} \\ &= \frac{-12x}{(x^2 - 3)^2}. \end{aligned}$$

At 2, we have $f'(2) = -24$. Thus the tangent line has the equation

$$y - 7 = -24(x - 2).$$

We simplify this to give $y = -24x + 55$ as the equation of the tangent line.

■

Exercise. Find the derivative of $f(x) = \frac{1+2x}{1-2x}$.

September 29, 2013