

1 Lecture 16: Higher derivatives

1.1 Outline

- Definition of higher-order derivatives
- Examples

1.2 Higher order derivatives

The acceleration is the derivative of the derivative of position. We call this a second derivative. We often write f'' for the derivative of f' and we call f'' the second derivative. We can define derivatives of any order by

$$f^{(0)} = f \quad \text{and} \quad f^{(n)} = f^{(n-1)'}$$

for the result of differentiating n -times. For derivatives up to order 2 or 3, we will add a prime (') but eventually we get tired of writing primes and will use one of the notations

$$f^{(n)} \quad \text{or} \quad \frac{d^n f}{dx^n}$$

Be careful to use the parentheses which help us to distinguish the n th-derivative $f^{(n)}$ from the n th power f^n .

1.3 Examples

Example. Find

$$\frac{d^2}{dx^2} x^2 e^x$$

If f is defined by $f(x) = e^x$, find $f^{(2012)}$.

Solution. The first derivative

$$\frac{d}{dx}(x^2 e^x) = x^2 e^x + 2x e^x = (x^2 + 2x)e^x$$

Differentiating again gives

$$\frac{d}{dx}(x^2 + 2x)e^x = (2x + 2)e^x + (x^2 + 2x)e^x = (x^2 + 4x + 2)e^x.$$

After spending all day computing 2012 derivatives, we find that $f^{(2012)}(x) = f^{(2011)}(x) = \dots = f(x) = e^x$.

■

Example. Can you find a formula for

$$\frac{d^n}{dx^n} x^n, \quad \frac{d^{2n}}{dx^{2n}} x^n \quad \frac{d^n}{dx^n} \frac{1}{x}?$$

Solution. Try a few examples and look for a pattern. ■

Example. Find a polynomial $p(x)$ of degree 2 so that with $f(x) = e^x$, $p(0) = f(0)$, $p'(0) = f'(0)$, and $p''(0) = f''(0)$.

Compare the values of e^x and $p(x)$.

Can you suggest a better way to approximate e^x ?

Solution. Since $f^{(k)}(x) = e^x$ for all k , we have $f^{(k)}(0) = 1$ for all $k = 0, 1, 2, \dots$. If $p(x) = ax^2 + bx + c$, then we have

$$p(0) = c, \quad p'(0) = b \quad p''(0) = 2a.$$

Since we want $p(0) = f(0) = 1$, we have $c = 1$. If $p'(0) = 1$, then we must have $b = 1$ and if $p''(0) = 1$, we must have $2a = 1$ or $a = 1/2$. Thus $p(x) = \frac{1}{2}x^2 + x + 1$.

Trying a few values we find

x	e^x	$p(x)$
0	1	1
0.01	1.010050167...	1.01005
0.2	1.22140...	1.22
-0.2	0.81873...	0.82

To do better we might look for a third degree polynomial with $p^{(k)}(0) = f^{(k)}(0)$ for $k = 0, 1, 2, 3$. ■

Example. Find the n th derivative of xe^x . Hint: Try a few and see if you can guess a pattern.

October 3, 2013